



- [GOAL Events](#)
- [Advocate Magazine](#)
- [Aquademia Podcast](#)
- [Blog](#)
- [Contact](#)
-
-
-
-
-
- [Log In](#)



- [About](#)
 - [Who We Are](#)
 - [Our History](#)
 - [Our Team](#)
 - [Sustainable Development Goals](#)
 - [Careers](#)
- [Membership](#)
 - [Overview](#)
 - [Our Members](#)
 - [Corporate Membership](#)
- [Resources](#)
- [Certification](#)
 - [Best Aquaculture Practices](#)
 - [Best Seafood Practices](#)

Search...



[Log In](#)

- [About](#)
 - [Who We Are](#)
 - [Our History](#)
 - [Our Team](#)
 - [Sustainable Development Goals](#)
 - [Careers](#)
- [Membership](#)
 - [Overview](#)
 - [Our Members](#)
 - [Corporate Membership](#)
- [Resources](#)
- [Certification](#)
 - [Best Aquaculture Practices](#)
 - [Best Seafood Practices](#)
- [GOAL Events](#)
- [Advocate Magazine](#)
- [Aquademia Podcast](#)
- [Blog](#)
- [Contact](#)



Amberjack culture progresses at Oceanic Institute



1 February 2004 Charles W. Laidley, Ph.D. Robin J. Shields, Ph.D. Anthony C. Ostrowski, Ph.D.



Longfin amberjack, or kahala, best suited to captivity



The natural spawning of wild-caught amberjacks at O.I. enabled the production of several thousand juveniles.

Amberjacks (*Seriola* sp.) have excellent aquaculture potential due to their adaptability to conditions of intensive culture, extremely fast growth, and high market value. Substantial commercial production already exists for yellowtail (*Seriola quinqueradiata*) and greater amberjack (*Seriola dumerili*) in Japan. The southern kingfish (*Seriola lalandi*) is raised in Australia. Seriolid culture is also being developed in the Mediterranean and Latin America.

However, a combination of difficulties in establishing reliable egg supply and commercially viable hatchery technologies has generally limited commercial amberjack production to the grow-out of wild-collected fingerlings. Even in Japan, where the industry is most developed, fingerling supplies continue to limit production and lead to large yearly shifts in market price.

Hatchery technology

With grant assistance from the National Oceanic and Atmospheric Administration and state of Hawaii, USA, the Oceanic Institute (O.I.) began working on hatchery technology for amberjack culture in the mid-1990s. It achieved natural spawns of wild-collected adults of both the greater amberjack and longfin or Almaco jack (*Seriola rivoliana*). Although both species have spawned at O.I., the longfin amberjack known locally as kahala has been much more adaptable to captivity. It is emerging as a prime candidate for the emerging offshore farming sector in Hawaii.

The natural spawning of wild-caught amberjacks in 1999 was a significant achievement that enabled the production of several thousand juveniles using intensive hatchery rearing techniques at O.I. The resulting F1 progeny were grown to maturation in 25-cubic-meter outdoor grow-out tanks using a commercial pelleted marine grower diet. The fish demonstrated excellent growth rates, achieving mean body weights of over 2 kg in the first year and near 5 kg at two years of age (Fig. 1).




Fig. 1: Growth of captive-spawned and -reared amberjack at Oceanic Institute. Fish were fed twice daily to satiation until 9 months of age and once daily thereafter. Water temperature was 26 degrees-C.

Although maintaining actively spawning stocks of wild-collected amberjacks has proven difficult due to the species' susceptibility to a number of parasitic and bacterial pathogens, the successful closing of the life cycle has allowed the use of domesticated F1 and F2 stocks as broodstock.

Domesticated broodstock

In captivity, male amberjack began maturing within the first year and actively spermiated at 21 to 22 months of age. Female amberjack mature slightly slower than males, with a rapid increase in gonadosomatic index and near-simultaneous onset of spawning at 24 months of age.

Under ambient photoperiod and temperature in Hawaii, F1 broodstock have spawned for over two years without interruption. They exhibit some seasonality in reproductive output, but remain in reproductive condition throughout the year (Fig. 2). Tanks containing 20 adults have yielded an average of 13 spawns per month with mean fecundity of 154,000 eggs per spawn and mean a fertility rate of 43 percent. Spawn size and fertility rates have both increased over time, indicating the value of long-term broodstock holding.

Fig. 2: Spawn frequency and egg production of captive-reared amberjack broodstock in outdoor spawning tanks.

In 2003, O.I. achieved spawning of the F2 populations at just under 2 years of age, indicating the ability of this species to be fully domesticated. Ongoing research at O.I. is directed at increasing stock biosecurity and improving fingerling production technologies to secure reliable year-round fingerling supplies at commercial scales of operation.

Amberjacks are being grown in submersible offshore cages in Hawaiian waters.

Offshore cage production

Pilot-scale production in offshore cages was initiated in January 2004. Several thousand amberjack fingerlings were stocked into a submersible offshore cage operated by Cates International to begin evaluating the performance of this species in the offshore environment.

(Editor's Note: This article was originally published in the February 2004 print edition of the Global Aquaculture Advocate.)

Now that you've finished reading the article ...


... we hope you'll consider supporting our mission to document the evolution of the global aquaculture industry and share our vast network of contributors' expansive knowledge every week.

By becoming a Global Seafood Alliance member, you're ensuring that all of the pre-competitive work we do through member benefits, resources and events can continue. Individual membership costs just \$50 a year.

Not a GSA member? Join us.

[Support GSA and Become a Member](#)


Authors

-  Charles W. Laidley, Ph.D.

Charles W. Laidley, Ph.D.


The Oceanic Institute
Makapuu Point
41-202 Kalaniana'ole Highway
Waimanalo, Hawaii 96795 USA

[103,114,111,46,101,116,117,116,105,116,115,110,105,99,105,110,97,101,99,111,64,121,101,108,100,105,97,108,99]

-  Robin J. Shields, Ph.D.

Robin J. Shields, Ph.D.





The Oceanic Institute
Makapuu Point
41-202 Kalanianaʻole Highway
Waimanalo, Hawaii 96795 USA

-  Anthony C. Ostrowski, Ph.D.

Anthony C. Ostrowski, Ph.D.

The Oceanic Institute
Makapuu Point
41-202 Kalanianaʻole Highway
Waimanalo, Hawaii 96795 USA

Share

-  [Share via Email](#)
-  [Share on Twitter](#)
-  [Share on Facebook](#)
-  [Share on LinkedIn](#)

Tagged With

[Robin J. Shields](#) [Oceanic Institute](#) [Anthony C. Ostrowski](#) [Charles W. Laidley](#) [Amberjack](#)

Related Posts

Health & Welfare

[Bivalent vaccine protects yellowtails, amberjack against pasteurellosis](#)

The authors developed an efficacious combination vaccine against *P. piscicida* and *L. garvieae* for yellowtail and amberjack. Field trials showed positive results, and the vaccine was approved for use in Japan.

Health & Welfare

[Novel soy proteins, oils replace fishmeal to achieve FIFO under 1:1 in amberjack](#)

In a nutrition trial, longfin amberjack received two diets composed of 40 percent soy protein concentrate and 10 percent fishmeal, and a control diet with 20 percent fishmeal.

Responsibility

[How aquaculture furthers the Blue Growth Initiative](#)

Blue Growth is as an integral part of sustainable global development and the Food and Agriculture Organization's Sustainable Development Goals as they relate to conservation of marine resources.

Health & Welfare

[Marine fish culture in Mexico](#)

Mexico has a wide diversity of areas and water temperatures that are suitable for a variety of aquaculture species.

About The Advocate

The Responsible Seafood Advocate supports the Global Seafood Alliance's (GSA) mission to advance responsible seafood practices through education, advocacy and third-party assurances.

[Learn More](#)

Search Responsible Seafood Advocate



Advertising Opportunities

[2022 Media & Events Kit](#)

Categories

[Aquafeeds](#) > [Health & Welfare](#) > [From Our Sponsors](#) > [Innovation & Investment](#) > [Intelligence Intelligence](#) > [Responsibility](#) > [Fisheries](#) > [Artículos en Español](#) >

Don't Miss an Article

Featured

- [Health & Welfare](#) [An update on vibriosis, the major bacterial disease shrimp farmers face](#)
- [Intelligence](#) [A seat at the table: Fed By Blue team says aquaculture needs a stronger voice](#)
- [Responsibility](#) [Quantifying habitat provisioning at macroalgae cultivation locations](#)

Popular Tags

All Tags ▼




Recent

- [Fisheries](#) [Second Test: Another filler for the fisheries category](#)
- [Fisheries](#) [Test: This is filler for the fisheries Category](#)
- [Aquafeeds](#) [Test Article](#)
- [Responsibility](#) [Study: Climate change will shuffle marine ecosystems in unexpected ways as ocean temperature warms](#)
- [Health & Welfare](#) [Indian shrimp researchers earn a patent for WSSV diagnostic tool](#)



- [About](#)
- [Membership](#)
- [Resources](#)
- [Best Aquaculture Practices \(BAP\)](#)
- [Best Seafood Practices \(BSP\)](#)
- [GOAL Events](#)
- [Advocate Magazine](#)
- [Aquademia Podcast](#)
- [Blog](#)
- [Contact](#)

Stay up to date with GSA

- 
- 
- 
- 
- 

Copyright © 2024 Global Seafood Alliance
All rights reserved.

[Privacy](#)

[Terms of Use](#)

[Glossary](#)