



- [GOAL Events](#)
- [Advocate Magazine](#)
- [Aquademia Podcast](#)
- [Blog](#)
- [Contact](#)
-
-
-
-
-
- [Log In](#)



- [About](#)
 - [Who We Are](#)
 - [Our History](#)
 - [Our Team](#)
 - [Sustainable Development Goals](#)
 - [Careers](#)
- [Membership](#)
 - [Overview](#)
 - [Our Members](#)
 - [Corporate Membership](#)
- [Resources](#)
- [Certification](#)
 - [Best Aquaculture Practices](#)
 - [Best Seafood Practices](#)

Search...



[Log In](#)

- [About](#)
 - [Who We Are](#)
 - [Our History](#)
 - [Our Team](#)
 - [Sustainable Development Goals](#)
 - [Careers](#)
- [Membership](#)
 - [Overview](#)
 - [Our Members](#)
 - [Corporate Membership](#)
- [Resources](#)
- [Certification](#)
 - [Best Aquaculture Practices](#)
 - [Best Seafood Practices](#)
- [GOAL Events](#)
- [Advocate Magazine](#)
- [Aquademia Podcast](#)
- [Blog](#)
- [Contact](#)



Aquaculture of Asian sea bass or barramundi



1 August 2000 Phil Boeing



Economics for recirculating culture are highly favorable



Aquaculture of barramundi is well established throughout Asia. Photo by Joerg Knoeckel.

The Asian sea bass (*Lates calcarifer*) is a euryhaline species that belongs to the Centropomidae (snook) family. It is widely distributed in the Indo-West pacific region, from the Arabian Gulf to China, Taiwan, the Philippines, the Indonesian archipelago, Papua New Guinea, and northern Australia. The species is known by various common names in different parts of its range, including “barramundi” in much of Asia, and “barramundi” (aboriginal for giant perch) in Australia and Papua New Guinea.

Barramundi supports important commercial and recreational fisheries throughout its range, and has developed into a mature aquaculture industry. Barramundi aquaculture industries are well established in Thailand, Malaysia, Singapore, Indonesia, Hong Kong, Taiwan, the Philippines and Australia.

Techniques for the propagation of barramundi were originally developed in Thailand in the 1970s and further advanced by research in the Philippines, Singapore, Hong Kong, and Australia (Grey 1987). With the exception of Australia, most barramundi are farmed in cages in freshwater or brackish water ponds, or in sea cages. In Australia, a number of indoor, closed-system barramundi farms have been established with controlled-environment buildings, using underground fresh or brackish water and recirculation technology.

Induced reproduction

Early work on the culture of barramundi relied on the use of fertilized eggs obtained from stripping males and females caught in estuary spawning grounds. However, hormonal treatments developed during the 1980s have become a standard procedure for many barramundi farms (Garcia 1990).

Barramundi broodfish may be kept in either fresh or saltwater, but must be placed in saltwater (28 to 35 ppt) prior to the breeding season to enable final gonadal maturation. In parts of Asia, barramundi are still induced to spawn by manipulation of environmental parameters, which simulate the migration of the species to the lower estuary and subsequent tidal regime at the time of natural spawning.

Salinity is gradually elevated in the brood tanks until full seawater is achieved. Once fish behavior indicates that spawning may be imminent, the tank temperature is raised to 31 to 32 degrees-C. The tank temperature is then dropped to 27 to 28 degrees-C to simulate tidal fluctuations in the lower estuary. Generally, most fish subjected to this technique will spawn.

More commonly, mature barramundi are spawned using a range of exogenous hormone preparations that are administered by injection, slowrelease cholesterol pellets, and osmotic pumps (Garcia 1989). The most common hormones are the luteinizing hormone-releasing hormone analogues (LHRHa) (Des-Gly10) D-Trp6, Pro9- LH-RH ethylamide, given at dosages of 3 to 5 micrograms per kilogram body weight for females and 40-100micrograms per kilogram for male barramundi (Almendras et al. 1988). Female barramundi broodfish generally average 2.5 to 6 kg and males from 2 to 3 kg. Females of this size can spawn up to five times after hormonal treatment and produce from three to 15 million eggs. Larger females produce considerably more eggs, with 35- to 40-kg females averaging around 45 million eggs per induced spawn. Hatching rates for barramundi spawned using hormonal manipulation range between 0.1 and 85 percent.



Barramundi culture in Australia is conducted in ponds, estuary cages, and intensive indoor facilities. Eagle Ranch Aquaculture in Beau Desert, Australia (pictured above), is designed to produce about ten tons of barramundi per year using recirculation and recycling. With daily backwashing of filters, it adds only 2 percent fresh water daily. Photos by Joerg Knoeckel.

Larval rearing

The requirements for larval rearing of barramundi are the same as for other finfish species. For hatchery rearing of barramundi larvae, initial stocking densities of ten to 40 fish per liter are common. To reduce cannibalism, it is recommended that the density be reduced to ten to 15 per liter on day 10, then to six per liter on day 21.

Most hatcheries have opted for a “green water” culture methodology, which maintains a microalgae concentration of *Nannochloropsis oculata* and/or *Tetraselmis* sp. in densities of 1 to 3 x 10⁵ cell/milligrams per liter of culture water. Larval barramundi begin feeding around 50 hours after hatching, at temperatures of 26 to 30 degrees-C and salinities of 28 to 31 ppt. It is best to have food available prior to first feeding (Parazo et al. 1990).

Daily feeding consists of 15 to 20 *Brachionus* sp. per ml on days 1 to 12, one half to two *Brachionus* sp. plus two to three artemia per milileter on days 12-15, and five to ten artemia per milileter on days 15 to 26 or until harvest.

The cladocerans *Daphnia*, *Moina* and *Diaphanosoma* have been used to supplement, or replace, brine shrimp as prey for intensively reared barramundi larvae. These species of zooplankton may be fed to barramundi from day 15, although they are more commonly used as a brine shrimp replacement from day 25.

A more recent development in larval rearing of barramundi involves extensive culture of day-old fish larvae in marine or brackish water ponds. Blooms of phytoplankton and zooplankton for feeding larvae are induced in ponds via the addition of organic and inorganic fertilizers (Rudledge and Rimmer 1991). Barramundi larvae are stocked in ponds at densities of 400,000 to 900,000 fish per hectare to coincide with peak densities of the smaller zooplankton-rotifers and copepods, which proliferate six to nine days after fertilization.

The survival of extensively reared barramundi is highly variable, ranging from 0 to 90 percent, with an average of 20 percent, compared to 15 to 40 percent for intensively hatchery reared larvae. However, the estimated production cost of extensively reared fingerlings is 40 to 64 percent that of intensively reared fingerlings, and this has resulted in greater adoption of extensive larval rearing techniques in Australia (Lobegeiger 1993).

Nursery techniques

Juvenile barramundi (1 to 2.5 cm total length) are transferred to nursery facilities after being harvested from rearing ponds or intensive culture tanks. If stocked directly in growout ponds, barramundi may be difficult to sizegrade. For this reason, cage culture in ponds or tank culture is more useful for the nursery phase, where cannibalism is a major cause of mortality. Cannibalism can be reduced by grading the fish every six to nine days, to ensure that the fish in the nursery system are similar in size.

Stocking densities of 80 to 100 fry per square meter are customary for cages for 30 to 45 days, or until the fingerling stage of 5 to 10 cm, where they can be graded and stocked in grow-out systems. Barramundi greater than 1 cm in length can be transferred from salt to freshwater in as little as 6 hours with no significant mortality.

Many barramundi farmers use fresh water for growout and therefore find it beneficial to use the same site for nursery culture. Barramundi can be weaned to artificial diets at 1 cm size. However, fish in the range of 1 to 2 cm show faster acceptance of artificial diets and better survival.

Grow-out procedures

Barramundi is cultured in southeast Asia using mostly extensive or semi-intensive technology, depending on the local market structure for sale of the fish, cost of feeds, and local infrastructure. Pond culture in these countries may use a stocking density of 2,000 to 3,000 fish per hectare, with harvest sizes ranging from 250 to 700 after a nine- to 12-month growout.

Cage culture farmers in Malaysia, the Philippines, and Thailand stock anywhere from 12 to 300 juvenile barramundi per square meter, depending on water quality and environmental conditions at the site. More fish are stocked in floating cages than in stationary cages because they are usually in sites with deeper water, lower salinity fluctuations, and better circulation of cleaner water.

Trash fish is the feed of choice for barramundi culture in most of southeast Asia, although some dry diets have been formulated (Aldon 1997). Seasonal variability of trash fish greatly impacts the growth rate achieved in this type of culture, with typical food conversion rates averaging 4:1 to 10:1. Average growth rates approximate 1 kg per year with survivals between 80 and 90 percent.

Recirculating culture in Australia

Australia has developed barramundi culture to a much more technical state through research and experimentation from both private and government institutions. There are three methods of growing barramundi in Australia.

The first is brackish water or freshwater pond culture, where usually the fish are maintained in cages to facilitate culling and harvesting. Some largescale cage production in estuarine waters is still practiced; however, difficulties with environmental issues and permitting have discouraged this type of culture. The third growing method is intensive production in recirculating indoor facilities.

Using this method, year-round steady-state production is possible and environmental concerns are avoided. Cage facilities stock as many as 100 kg of fish per square meter cage surface and cull to a harvest biomass of 30 to 50 kg per square meter. Recirculation facilities average 60 to 80 kilograms per square meter harvest density of fish using European recirculation technology initially developed for high-density culture of eels.

Most recirculation farms culture plate-sized barramundi (350 to 1,000 grams) obtainable after six to eight months in culture. Some cage farms culture larger fish (up to 2 kg) for a select fillet market.

The development of high-energy feeds has increased the productivity of the Australian barramundi industry (Catacutan and Coloso 1995). Fish growth is maximized with high-protein diets (>55 percent) and energy levels (digestible energy) of at least 18.0 megajoules per kilogram or higher (Williams et al. 1997). These diets have lowered the food conversion rate to 0.75: 1 through weight gains of 2 percent body weight per day.

Market and economics

The high market value of barramundi has fueled the aquaculture of this species in Australia to a greater degree than other producing countries. Ex-farm price of barramundi in Malaysia, Thailand, and the Philippines is generally under \$1.00 U.S. per kilogram (Buendia 1997). Only Hong Kong and Singapore markets for live barramundi command high prices, but these are relatively small niches (Surtida 1997). Australia possesses a very large fresh seafood demand, and this demand is chiefly responsible for the tremendous growth of barramundi farming. The wholesale market price for barramundi has stabilized at \$5.20 to 6.30 U.S. per kilogram for the last few years, having come down due to increased production from \$11.50 U.S. per kilogram in the early 1990s (Anderson 1998).

The principal growing areas of Australia produced 624 metric tons (MT) of barramundi during the 1996-1997 growing season, valued at \$3.7 million U.S. Gilled and gutted fish generally sell for \$2 per kilogram more than whole fish.

A profile of the economics of recirculating barramundi systems in Australia shows classic economy of scale returns. A 30-MT per year production facility will cost around \$460,000 in capital investment, and require a break-even selling price of \$4.90 per kilogram, whereas a 100-MT barramundi farm will cost around \$860,000 in capital investment but demonstrate a break-even price of \$2.89 per kilogram. Obviously, the larger farm merits attention in terms of risk management for any possible market price decline.

With over a third of the current increase in Australian production of barramundi coming from recirculating systems, it is clear that the economics for this type of culture are highly favorable. New developments in genetics and growth hormones for barramundi are already showing favorable results for lowering the production costs and further increasing profit margins (Anderson 1998). Note: Cited references upon request to the author.

(Editor's Note: This article was originally published in the August 2000 print edition of the Global Aquaculture Advocate.)

Now that you've finished reading the article ...


... we hope you'll consider supporting our mission to document the evolution of the global aquaculture industry and share our vast network of contributors' expansive knowledge every week.

By becoming a Global Seafood Alliance member, you're ensuring that all of the pre-competitive work we do through member benefits, resources and events can continue. Individual membership costs just \$50 a year.

Not a GSA member? Join us.

[Support GSA and Become a Member](#)

Author





-  Phil Boeing

Phil Boeing

Director of Research and Development
Aquafauna Bio-Marine, Inc.
P.O. Box 5
Hawthorne, California 90250 USA

[116,101,110,46,116,116,97,46,116,101,110,100,108,114,111,119,64,103,110,105,101,111,98,46,112]

Share

-  [Share via Email](#)
-  [Share on Twitter](#)
-  [Share on Facebook](#)
-  [Share on LinkedIn](#)

Tagged With

[barramundi](#) [Phil Boeing](#)

Related Posts

Intelligence

[Barramundi culture in Australia](#)

From recirculation units, earthen ponds and sea pens in Australia, barramundi are spreading to other continents due to the advent of sophisticated modular culture systems.

Intelligence

[Adding flavor complexity to farmed barramundi](#)

Organoleptic attributes such as flavor and aroma are among the most important factors that influence consumer acceptability and demand for fish products. Consumers have identified farmed fish as less complex and lacking “sealike” or “sea-fresh” flavors and aromas.

Health & Welfare

[Effects of poultry protein concentrate, phosphorus supplementation in barramundi juveniles](#)

This study characterized the nutritional value of a high-quality poultry protein concentrate against standard poultry meal in barramundi juveniles.

Intelligence

[Aquaculture key to increasing seafood supplies to Arab states](#)

Arab States have substantial natural resources to increase aquaculture production. Several types of systems are readily adaptable and can be implemented relatively quickly.

About The Advocate

The Responsible Seafood Advocate supports the Global Seafood Alliance’s (GSA) mission to advance responsible seafood practices through education, advocacy and third-party assurances.

[Learn More](#)

Search Responsible Seafood Advocate



Advertising Opportunities

[2022 Media & Events Kit](#)

Categories

[Aquafeeds](#) > [Health & Welfare](#) > [From Our Sponsors](#) > [Innovation & Investment](#) > [Intelligence Intelligence](#) > [Responsibility](#) > [Fisheries](#) > [Artículos en Español](#) >

Don't Miss an Article

Featured

- [Health & Welfare](#) [An update on vibriosis, the major bacterial disease shrimp farmers face](#)
- [Uncategorized](#) [A seat at the table: Fed By Blue team says aquaculture needs a stronger voice](#)
- [Responsibility](#) [Quantifying habitat provisioning at macroalgae cultivation locations](#)

Popular Tags

All Tags ▼




Recent

- [Fisheries](#) [Second Test: Another filler for the fisheries category](#)
- [Fisheries](#) [Test: This is filler for the fisheries Category](#)
- [Aquafeeds](#) [Test Article](#)
- [Responsibility](#) [Study: Climate change will shuffle marine ecosystems in unexpected ways as ocean temperature warms](#)
- [Health & Welfare](#) [Indian shrimp researchers earn a patent for WSSV diagnostic tool](#)



- [About](#)
- [Membership](#)
- [Resources](#)
- [Best Aquaculture Practices \(BAP\)](#)
- [Best Seafood Practices \(BSP\)](#)
- [GOAL Events](#)
- [Advocate Magazine](#)
- [Aquademia Podcast](#)
- [Blog](#)
- [Contact](#)

Stay up to date with GSA

- 
- 
- 
- 
- 

Copyright © 2024 Global Seafood Alliance
All rights reserved.

[Privacy](#)

[Terms of Use](#)

[Glossary](#)