Global Seafood Alliance Logo

- GOAL Events
- Advocate Magazine
- Aquademia Podcast
- <u>Blog</u>
- Contact
- 0
- X
- in
- Log In

- About
 - Who We Are
 - Our History
 - o Our Team
 - Sustainable Development Goals
 - Careers
- <u>Membership</u>
 - o <u>Overview</u>
 - Our Members
 - Corporate Membership
- Resources
- Certification
 - Best Aquaculture Practices
 - Best Seafood Practices

Search...

Q

Log In

- About
 - Who We Are
 - o Our History
 - o Our Team
 - Sustainable Development Goals
 - o <u>Careers</u>
- Membership
 - o <u>Overview</u>
 - o Our Members
 - Corporate Membership
- Resources
- Certification
 - Best Aquaculture Practices
 - Best Seafood Practices
- GOAL Events
- Advocate Magazine
- Aquademia Podcast
- **Blog**
- Contact

Health & Welfare



'Big picture' connects shrimp disease, inbreeding

Responsible Seafood Advocate logo

1 September 2014 Roger W. Doyle, Ph.D.



Inbreeding fundamentally a management issue controllable at farm level



Smaller shrimp aquaculture operations may face the greatest risks from problematic breeding practices.

The main effect of inbreeding among animals is to make a bad environment worse. As environments deteriorate, inbred animals die faster than noninbreds. The more inbred the stock, the greater the difference in survival between inbred and non-inbred animals.

Amplification of stress mortality has been shown time and time again in humans, fish, shrimp and many other animals and plants in laboratory, farm and natural environments. As far as we know, all non-accidental causes of mortality, including disease, nutritional and temperature stress and competition, are amplified by inbreeding.

It is very important to realize that inbreeding does not cause, but amplifies disease mortality. In this, it is like the way drunk driving amplifies the death rate from highway accidents. Sober drivers have accidents, too, for all kinds or reasons, but the mortality from accidents is higher when drivers are drunk. By analogy, mortality from all kinds of lethal stress, including disease, is higher when stocks are inbred.

Causal paths, EMS/AHPN

The environmental stress of most concern to shrimp aquaculture right now is a disease called early mortality syndrome (EMS) or acute hepatopancreatic necrosis (AHPN). The causal path diagram in Fig. 1 summarizes the current widely held "big picture" relationship between diseases, including EMS/AHPN, and the aquaculture industry.

Exposure To Pathogens Biosecurity Regulations Disease Mortality Fear

Fig. 1: Beneficial paths between biosecurity regulation and mortality from disease.

The disease mortality variable on the right side lumps together various measures of the seriousness of a disease, such as incidence, prevalence and morbidity. The biosecurity regulations variable on the left lumps together most, if not all, current and planned regulatory actions intended to diminish the disease problem, such as prohibition of stocks that are not officially free of listed pathogens, prohibitions on the transfer of animals among geographical regions, quarantines and "kill orders" imposed on infected farms and hatcheries.

The causal paths in Fig. 1 work in the following way. When something increases mortality in a farming region, like the emergence of a new pathogen, rising concern among farmers, hatcheries and breeders puts pressure on bureaucrats, scientists and consultants who, in turn, increase the level of regulatory activity. The path that connects disease to regulation is labeled "Fear" in Fig. 1. The fear pathway is positive because when the cause (mortality) increases, the effect (regulation) increases, too.

The second causal path in Fig. 1 is labeled "Exposure" because the essential purpose of most regulatory activity is to reduce exposure of animals in the farm ponds to pathogens. The exposure pathway is negative, because when regulatory activity goes up, mortality goes down.

Fig. 1 is obviously a simplified model of the disease/regulatory relationship. Whenever you simplify a model, you should also simplify the question you ask of it. In this case, we simply ask, "Is the system stable?"

Any engineer, statistician or audio technician will answer "yes," because Fig. 1 is a negative feedback loop. Something outside the loop drives mortality up, fear drives regulation up, increased regulation drives exposure down, and as exposure goes down, so does mortality. The model in Fig. 1 is selfdamping and predicts that when a new pathogen arises, the aquaculture system should "vibrate" around some stable level of regulation and disease mortality. If the big picture path diagram in Fig. 1 is essentially correct, then our handling of disease crises is on the right track.

Susceptibility

However, the path model in Fig. 1 is incomplete. A third causal path flows from biosecurity regulation to disease mortality, labeled "Susceptibility" in Fig. 2. Susceptibility to pathogens and other environmental stresses is where inbreeding enters the picture. As mentioned above, inbreeding amplifies the effects of environmental stress from poor pond management and exposure to pathogens.




Fig. 2: Detrimental (inbreeding paths) between biosecurity regulation and disease mortality.

The susceptibility path is positive for a very simple reason. More biosecurity means more postlarvae are produced by matings among close relatives and are therefore inbred. Or in genetic terms, increased regulation reduces the effective population size of the gene pool available to breeders and hatcheries in a farming region.

Closed borders inhibit the entry of new genetic diversity. Breeders respond to the increased value of their broodstock by selling postlarvae that generate inbred offspring when they are used as breeders. Good breeding companies never sell inbred postlarvae to hatcheries or farmers. The inbreeding problem arises from "copying."

Requirements for specific pathogen-free status also limit diversity. Both kinds of regulations reduce local availability of postlarvae because of limited production capacity and/or high prices. Small "copy hatcheries" that use few and often closely related breeders and do not follow good broodstock management practices increase production to fill the demand for postlarvae.

Increased mortality raises the level of fear and therefore regulation in a positive path. Regulation increases inbreeding and susceptibility to stress, also in a positive path. Susceptibility increases mortality. And so we go, round and round, because Fig. 2 is a self-amplifying feedback loop.

Greater effect on mortality?

All three causal paths are shown in Fig. 3. So we ask the simple question, "Which has the greater effect on mortality, the negative, stabilizing exposure path or the positive, destabilizing susceptibility path?"

Fig. 3: Estimated relative strengths of beneficial (blue) and detrimental (red) paths.

The best answer we have now, unfortunately, is that the susceptibility path is dominant, and the big picture of the current shrimp disease management system is unstable. In a technical paper by the author on shrimp inbreeding (available online at http://onlinelibrary.wiley.com), the author estimates that at least 70 percent of farm postlarvae come from hatcheries that enjoy none of the benefits of increased regulatory activity (lower exposure), but suffer the consequences of inbreeding (higher susceptibility). The path diagram in Fig. 3 is weighted accordingly.

How did this come about? Why does the current, widely held consensus resemble Fig. 1 rather than Fig. 3? It is partly because current proposals for disease management (increasing regulations to reduce exposure to pathogens) generally ignore inbreeding as a side effect. Partly because geneticists have focused on telling hatcheries how to manage their broodstock properly fully thinking through what happens in the poorly run hatcheries that produce 70 percent of the postlarvae. Partly because of a technical glitch: inbreeding at the farm level is often estimated incorrectly with microsatellite markers.

What happens next?

We know that self-amplifying feedback systems tend to end up either generating a sustained, unbearable howl at maximum amplitude or shake themselves apart. The component of shrimp aquaculture systems most likely to disappear during self-destruction is small-holder farming.

There is another possible outcome that retains the direct, beneficial effect of biosecurity regulation. The people who care about the industry could weaken the indirect (susceptibility) path that leads from biosecurity, through increased inbreeding, to mortality. But is it possible to do this while allowing the present diversity of hatchery and farming styles to persist?

Perhaps. Up to now, inbreeding has been considered solely a problem in broodstock management rather than farm management. Yet it is on the farm that mortality occurs from disease; from interactions among disease, oxygenation, competition and weather; and from inbreeding interactions - with all of these stresses alone and in combination.

Perspectives

It is my personal opinion that inbreeding is fundamentally a management issue that should be controlled at farm level by giving farmers access to verifiable information on the genetic quality of postlarvae, just as they need verifiable information about feed quality, SPF status and oxygen levels to manage ponds properly. Farmers can control inbreeding by choosing the postlarvae supplier, which can be either a big or a small company. If it is a small company, farmers should be able to tell which supplier produces postlarvae that are inbred, and which are not.

It might surprise many non-geneticists that obtaining and verifying this kind of information need not be technically difficult or expensive on a routine basis.

Editor's Note: The quantitative assertions made or implied in this article are documented in a pre-publication technical paper that is freely available for downloading from the journal Aquaculture Research. Search the website http://onlinelibrary.wiley.com for the phrase "Edwards inbreeding disease

(Editor's Note: This article was originally published in the September/October 2014 print edition of the Global Aquaculture Advocate.)

Author

Roger W. Doyle, Ph.D.

Roger W. Doyle, Ph.D.

Roger W. Doyle, Ph.D. Genetic Computation Ltd. 1-4630 Lochside Drive Victoria, British Columbia, Canada V8Y2T1

[109,111,99,46,112,109,111,99,101,110,101,103,64,101,108,121,111,100,114]

Share

- **Share via Email Share via Email**
- **У** Share on Twitter
- **f** Share on Facebook
- in Share on LinkedIn

Tagged With

disease EMS inbreeding mortality Roger W. Doyle Big Picture AHPN

Related Posts

Health & Welfare

Study: Inbreeding affects body weight, not survival, in white shrimp

Inbreeding commonly causes a decrease in the mean values of traits of productive interest in aquaculture. Results of a study, however, showed that inbreeding had no effect on grout survival of Pacific white shrimp from 65 to 130 days of age.

Intelligence

Bangladesh's tilapia aquaculture industry shows resilience

Tilapia aquaculture in Bangladesh has developed significantly since 1999, based on the Genetically Improved Farmed Tilapia (GIFT) strain of Nile tilapia (Oreochromis niloticus) introduced from Malaysia and on the significant genetic improvement research work by the Bangladesh Fisheries Research Institute (BFRI).

Health & Welfare

Selection for disease resistance in Indian white shrimp

A new breeding program for genetic improvement of Indian white shrimp (Fenneropenaeus indicus) in Egypt was established at a private shrimp farm at AL Dibah Triangle Zone (DTZ) to develop disease-resistant shrimp for the country's aquaculture industry.

Health & Welfare

A holistic management approach to EMS

Early Mortality Syndrome has devastated farmed shrimp in Asia and Latin America. With better understanding of the pathogen and the development and improvement of novel strategies, shrimp farmers are now able to better manage the disease.

About The Advocate

The Responsible Seafood Advocate supports the Global Seafood Alliance's (GSA) mission to advance responsible seafood practices through education, advocacy and third-party assurances.

Learn More

Search Responsible Seafood Advocate Search Search Q



Advertising Opportunities

2022 Media & Events Kit

Categories

Aquafeeds > Health & Welfare Health & Welfare > From Our Sponsors > Innovation & Investment > Intelligence > Responsibility > Fisheries > Artículos en Español >

Don't Miss an Article

Featured

- Health & Welfare An update on vibriosis, the major bacterial disease shrimp farmers face
- Intelligence A seat at the table: Fed By Blue team says aquaculture needs a stronger voice
- Responsibility Quantifying habitat provisioning at macroalgae cultivation locations

Popular Tags



Recent

- Fisheries Second Test: Another filler for the fisheries category
- Fisheries Test: This is filler for the fisheries Category
- Aquafeeds Test Article
- Responsibility Study: Climate change will shuffle marine ecosystems in unexpected ways as ocean temperature warms
- Health & Welfare Indian shrimp researchers earn a patent for WSSV diagnostic tool



- **About**
- <u>Membership</u>
- Resources
- Best Aquaculture Practices (BAP)
- Best Seafood Practices (BSP)
- GOAL Events
- Advocate Magazine
- Aquademia Podcast
- Blog
- Contact

Stay up to date with GSA

- 0

Copyright © 2024 Global Seafood Alliance All rights reserved.

Privacy Terms of Use <u>Glossary</u>