Global Seafood Alliance Logo

- GOAL Events
- Advocate Magazine
- Aquademia Podcast
- Blog
- Contact
- (O)

- <u>Log In</u>

- About
 - Who We Are
 - Our History
 - o Our Team
 - Sustainable Development Goals
 - <u>Careers</u>
- <u>Membership</u>
 - Overview
 - Our Members
 - Corporate Membership
- Resources
- **Certification**
 - Best Aquaculture Practices
 - Best Seafood Practices

Search...



Log In

- About
 - o Who We Are
 - Our History
 - o Our Team
 - Sustainable Development Goals
 - Careers
- <u>Membership</u>
 - o <u>Overview</u>
 - Our Members
 - o Corporate Membership
- Resources
- **Certification**
 - Best Aquaculture Practices
 - Best Seafood Practices
- GOAL Events
- Advocate Magazine

- <u>Aquademia Podcast</u>
- Blog
- Contact




Biofloc consumption by Pacific white shrimp postlarvae

Responsible Seafood Advocate logo

9 July 2016 Sabrina M. Suita Alessandro P. Cardozo Paulo Cesar Abreu Wilson Wasielesky Jr., Ph.D.



Using the stable isotope technique to determine relevance of food sources



Different food sources (A: Artemia; B: bioflocs; C: commercial aquafeed; D: ciliates; E, F: microalgae) and shrimp postlarvae (G, H) used in the trial.

In the early stages of marine shrimp culture, a diet including live food can be more efficient as a food source because these food items remain viable for a longer time in the water column and have high digestibility. However, live food is expensive to produce and maintain, and depends on highly-trained workers for their production.

The Biofloc Culture Systems – BFT (Biofloc Technology) can be a good source of live food in the grow-out phase of marine shrimp production. This technology is based on the addition of an organic carbon source that is used by bacteria to convert the excess nitrogen in the microbial biomass in the production tanks, and thus it can reduce water exchanges. In the early stages of larval shrimp rearing (from PL-1 phase until PL-30), shrimp production is generally performed in clear water systems with the addition of food items like phytoplankton, zooplankton

(Artemia sp.) and commercial feeds, and using high water exchange rates. An alternative feed can be the incorporation of BFT systems for these early life stages. However, information is needed on the consumption and assimilation of bioflocs by young, developing postlarvae of shrimp.



Production of live food for the study. A&B: Artemia culture; C&D: microalgae cultures; E: Harvesting decapsulated *Artemia* nauplii.

The stable isotope technique has been used as a complement to conventional techniques used in diet studies, because it can show the proportion of each prey/feed assimilated into the predator animal's diet. The principle of this technique is that stable isotopes of different chemical elements in the animal tissue reflect the ratio present in the food consumed, and this is registered as a signature in the new formed tissues of the consuming animal.

Study setup



Management of nursery system.

The experiment lasted for 30 days, starting with one-day-old (PL1) shrimp postlarvae, and used two different strategies to control ammonia when concentrations reached 0.5 mg/L. The first strategy was water exchange at a rate of approximately 70 percent of the total volume (the control treatment), and the second strategy included organic fertilization with dextrose (the BFT treatment) with no water exchange.

In both treatments, shrimp were fed with three different commercial feeds (CF), according to the animal stage development; shrimp were fed three times per day with commercial feeds, and once with frozen *Artemia* sp. nauplii. The microalgae *Chaetoceros muelleri* was inoculated into the experimental tanks in the first experimental day at a concentration of 5×10^4 cells/mL.



Weighing samples for stable isotopes analysis.

For water quality monitoring, dissolved oxygen, temperature and pH were measured daily. Ammonia (N- (NH3 + NH4 $^+$)) was analyzed every two days. Nitrite (N-NO $_2^-$) and phosphate (P-PO $_4^{3-}$) were measured weekly. Alkalinity (mg/L CaCO $_3$) and nitrate (N-NO $_3^-$) were determined in the initial and final samples. The growth performance of test animals was evaluated by size, weight and survival.


Samples of different food sources, bioflocs and shrimp postlarvae were collected on the 10^{th} , 20^{th} and 30^{th} days of the experiment to determine the carbon and nitrogen isotopic values. All of the samples were ground, weighed and placed into tin capsules. The analyses of carbon (13 C / 12 C) and nitrogen (15 N / 14 N) isotopic ratios were performed at the UC Davis Stable Isotope Facility, CA, USA. The Bayesian model of isotopic mixtures (package SIAR) was used to determine the relative contributions of food sources to the growth of shrimp in the different culture periods.

Results and perspectives

During the trial, no differences were determined in the performance parameters analyzed for the shrimp postlarvae (Table 1), or in the water quality parameters (Table 2), all of which remained within recommended values for *L. vannamei*. The commercial feeds tended to increase their contribution to the animals in both treatments (Figures 1 and 2). However, we observed that the biofloc contribution was around 50 percent (Figure 2). In the BFT treatment, the bioflocs appeared to have an important role in the performance of *L. vannamei* postlarvae. The microalgae presented a similar contribution when compared to the bioflocs in the BFT treatment, and in both treatments, microalgae were assimilated more than the *Artemia* sp. nauplii.

Treatments

Nitrate (mg L-1)






Fig. 1 (left): Average values (proportion x 100 percent) of the contribution of different food sources provided to Pacific white shrimp postlarvae during production in the CT (Control Treatment) on the 30th day. Fig. 2 (right): Average values (proportion x 100 percent) of the contribution of different food sources provided to Pacific white shrimp postlarvae during production in the BT (Biofloc Treatment) on the 30th day.

BT

 1.32 ± 1.17 1.9 ± 2

Survival (%)	93.2 ± 7.11	91.2 ± 8.53	
Initial biomass (g) $0.065 \pm 0.002 \ 0.065 \pm 0.002$			
Final biomass (g)	3.16 ± 0.7	3.14 ± 0.44	
Treatme	nts	\mathbf{CT}	BT
Temperature (degr	ees C)	27.8 ± 1.14	28.13 ± 1.33
Oxygen (mg L-1)		5.72 ± 0.4	5.73 ± 0.49
Salinity		$30.16 \pm 0.98a$	$31.33 \pm 0.52b$
pН		8.05 ± 0.24	8.04 ± 0.23
Alkalinity (mg L-1	l as CaCO3)	$248.75 \pm 2.5a$	$112.5 \pm 14.43b$
Phosphate (mg L-1	1)	0.99 ± 0.6	1.93 ± 1.21
Ammonia (mg L-1	1)	0.65 ± 0.15	0.65 ± 0.1
Nitrite (mg L-1)		$0.7 \pm 0.15a$	$1.34 \pm 0.3b$

 \mathbf{CT}

It was observed that *Artemia* sp. had a smaller contribution than other food sources in the growth of postlarvae (Figures 1 and 2). Despite being financially advantageous to reduce its use, it is unknown what proportions or what direct effect any reduction in the use of *Artemia* sp. could cause in shrimp postlarvae, because this food source provides nutrients that are necessary for the proper development of early life stages of penaeid shrimp.

During the pre-nursery phase, the influence of inert feed (or commercial feed) on the growth of the *L. vannamei* postlarvae increased over time, but young animals in the presence of bioflocs can obtain nutritional benefits, increasing their growth. In addition to this nutritional increase, fertilizations with carbon sources can reduce the amount of water used in the hatchery/pre-nursery phase.

Authors

• 📝 Sabrina M. Suita

Sabrina M. Suita

Marine Station of Aquaculture Institute of Oceanography Federal University of Rio Grande C.P.474-Rio Grande (RS)-CEP 96201-900- Brazil

[114,98,46,109,111,99,46,111,111,104,97,121,64,97,116,105,117,115,109,115]

• Alessandro P. Cardozo

Alessandro P. Cardozo

Marine Station of Aquaculture Institute of Oceanography Federal University of Rio Grande C.P.474-Rio Grande (RS)-CEP 96201-900- Brazil

 $\llbracket 109,111,99,46,108,105,97,109,116,111,104,64,111,114,100,110,97,115,115,101,108,97,99,111 \rrbracket$

• Paulo Cesar Abreu

Paulo Cesar Abreu

Marine Station of Aquaculture Institute of Oceanography Federal University of Rio Grande C.P.474-Rio Grande (RS)-CEP 96201-900- Brazil

[114,98,46,103,114,117,102,64,97,99,112,99,111,100]

• Wilson Wasielesky Jr., Ph.D.

Wilson Wasielesky Jr., Ph.D.

Marine Station of Aquaculture Institute of Oceanography Federal University of Rio Grande C.P.474-Rio Grande (RS)-CEP 96201-900- Brazil

[114,98,46,109,111,99,46,115,117,114,107,105,109,64,119,111,110,97,109]

Share

- Share via Email
- **Y** Share on Twitter
- **f** Share on Facebook
- in Share on LinkedIn

Tagged With

stable isotope technique biofloc consumption <u>Litopenaeus vannamei</u> <u>Artemia Wilson Wasielesky Paulo Cesar</u> Abreu Alessandro P. Cardozo Sabrina M. Suita

Related Posts

Innovation & Investment

Brazil study results encouraging for injector

Zero-exchange biofloc systems allow elevated stocking densities and production, but also require more dissolved oxygen and thorough water circulation. A new type of air injector uses only a centrifugal pump to recirculate water while naturally aspirating ambient air.

Health & Welfare

Shrimp biofloc production trials in Saudi Arabia

An R&D trial was conducted with shrimp biofloc technology at a commercial facility in Saudi Arabia. Results indicated that biofloc technology can be effectively applied, with some protocol adjustments, in the KSA's extreme desert environmental conditions.

Intelligence

Biofloc technology production promising in temperate zones

A study was conducted to assess the feasibility to grow Channel catfish (Ictalurus punctatus) in an outdoor biofloc system during winter in a temperate zone. High biomasses of market-size channel catfish were successfully maintained through the winter with high survival and in good condition in both treatments.

Aquafeeds

Biofloc systems viable for tilapia production

Well-designed and managed biofloc technology systems are a viable alternative for production of various species like tilapia, to increase feed efficiency by lowering aquafeed protein requirements and to help reduce or eliminate effluent discharges.

About The Advocate

The Responsible Seafood Advocate supports the Global Seafood Alliance's (GSA) mission to advance responsible seafood practices through education, advocacy and third-party assurances.

Learn More

Search Responsible Seafood Advocate Search Search



Advertising Opportunities

2022 Media & Events Kit

Categories

<u>Aquafeeds Aquafeeds</u> <u>Health & Welfare</u> <u>From Our Sponsors</u> <u>Innovation & Investment</u> <u>Intelligence</u> <u>Nesponsibility</u> <u>Fisheries</u> <u>Artículos en Español</u>

Don't Miss an Article

Featured

- Health & Welfare An update on vibriosis, the major bacterial disease shrimp farmers face
- Intelligence A seat at the table: Fed By Blue team says aquaculture needs a stronger voice
- Responsibility Quantifying habitat provisioning at macroalgae cultivation locations

Popular Tags

All Tags ✓

Recent

- Fisheries Second Test: Another filler for the fisheries category
- Fisheries Test: This is filler for the fisheries Category
- Aquafeeds Test Article
- Responsibility Study: Climate change will shuffle marine ecosystems in unexpected ways as ocean temperature warms
- Health & Welfare Indian shrimp researchers earn a patent for WSSV diagnostic tool



- About
- Membership
- Resources
- Best Aquaculture Practices (BAP)
- Best Seafood Practices (BSP)
- GOAL Events
- Advocate Magazine
- Aquademia Podcast
- Blog
- Contact

Stay up to date with GSA

Copyright © 2024 Global Seafood Alliance All rights reserved.

Privacy Terms of Use Glossary