## Global Seafood Alliance Logo

- GOAL Events
- Advocate Magazine
- Aquademia Podcast
- Blog
- Contact
- 0
- . 4
- X
- in
- 🔼
- Log In

- About
  - Who We Are
  - Our History
  - Our Team
  - Sustainable Development Goals
  - <u>Careers</u>
- <u>Membership</u>
  - Overview
  - Our Members
  - Corporate Membership
- Resources
- Certification
  - Best Aquaculture Practices
  - Best Seafood Practices

Search...



# Log In

- About
  - Who We Are
  - Our History
  - o Our Team
  - Sustainable Development Goals
  - Careers
- Membership
  - o <u>Overview</u>
  - Our Members
  - o Corporate Membership
- Resources
- Certification
  - Best Aquaculture Practices
  - Best Seafood Practices
- GOAL Events
- Advocate Magazine

- Aquademia Podcast
- Blog
- Contact



# Brazil study results encouraging for injector

### Responsible Seafood Advocate logo

29 September 2015 Dariano Krummenauer, Ph.D. Bob Advent Carlos Gaona, Ph.D.; Geraldo Foes, Ph.D.; Aline Bezerra, M.S.; Alessandro Cardozo, Ph.D.; Luis Poersch, Ph.D.; Wilson Wasielesky Jr., Ph.D.



# Aeration in super-intensive shrimp culture



Shrimp stocked in raceways equipped with air injector aeration for a growout study achieved weekly growth rates similar to those reported in super-intensive systems in southern Brazil.

Biofloc technology systems represent a new alternative for shrimp culture. These systems utilize minimal to zero water exchange and improve water quality through the formation of a microbial community consisting of microorganisms that include bacteria, microalgae, protozoa and zooplankton. In addition to improved water quality, bioflocs also supplement shrimp diets.

The formation of biofloc is stimulated by the addition of a carbon source to maintain a carbon:nitrogen ratio ranging from 12:1 to 20:1, depending upon physical, chemical and biological interactions. These interactions are stimulated by increased rates of mixing and oxygenation.

In early stages of culture, the abundance of bacteria that use the dissolved organic matter in the environment for growth increases. Conversely, low oxygen concentrations can negatively influence the floc-forming microbial community and lead to poor water quality and reduced performance of the cultured animals.

## **Aeration in biofloc systems**

Zero-exchange biofloc systems allow elevated stocking densities and production, but also require more dissolved oxygen, degassing and thorough mixing and circulation of the water column. Several aeration devices have been used in the systems. The most common aerators in super-intensive raceways are venturis and diffusers (air stones and tubes driven by blowers), while paddlewheel aerators are more common in lined ponds for intensive culture.

Recently, a new type of air injector has been used in raceway systems. These injectors use only a centrifugal pump to recirculate the water while naturally aspirating ambient air, sheering it into the water column at an air:water ratio of 3:1. This sheering produces fine microbubbles and high transfer rates, and eliminates the need for pure oxygen supplementation.

Mounted on the bottoms of raceways, the injectors mix the water vertically while creating a horizontal current to keep solids in suspension and assist in the distribution of feeds. The injectors' air intakes or "snorkels" can also be built into a manifold for the ability to select and adjust the source and location of the air being injected.

Utilizing the hot air accumulated near the top of a greenhouse allows for increased heating of the water, while drawing fresh air from outside aids in the reduction of carbon dioxide build-up in the greenhouse. By performing multiple tasks efficiently, these injectors can be a good alternative for super-intensive, zero-exchange biofloc systems.

# **Study setup**

To compare the performance of the air injectors with that of the diffuser tubes and paddlewheel aerators typically used in Southern Brazil, nursery and growout phase studies were performed at the Marine Station of Aquaculture of the Federal University of Rio Grande in southern Brazil.

Shrimp were fed commercial diets with 38 and 40 percent crude protein, with feeding rates adjusted according to weekly growth. For both studies, organic fertilization employed molasses from sugar cane as a carbon source. Following daily measurements, pH and alkalinity were corrected with calcium hydroxide whenever the pH was below 7.0 and alkalinity was equal to or less than 100 mg calcium carbonate/L.

## **Nursery phase study**

Pacific white shrimp (*Litopenaeus vannamei*) postlarvae from a commercial hatchery in Canguaretama, Rio Grande do Norte State, Brazil, were stocked at 2,500 shrimp/m3 in two, 70-m3 tanks operated with zero water exchange and raised for 30 days until each individual reached 1.0-g weight. The tanks were filled with a mixture of sand-filtered seawater with 28-ppt salinity and 5-ppt groundwater, and inoculated with biofloc-rich water from a previous growout study at 10 percent of the total volume.

Table I. Water quality parameters in a nursery phase biofloc system raising white shrimp.

Parameter	Injectors	Air Hoses
Temperature (°C)	28.9	27.2
Dissolved oxygen (mg/L)	5.77	6.05
pH	7.74	7.63
Salinity (ppt)	23.1	23.8
Total ammonia-nitrogen (mg/L)	0.70	0.18
Nitrite-nitrogen (mg/L)	1.90	0.72
Nitrate-nitrogen (mg/L)	4.83	5.92
Phosphate (mg/L)	1.40	1.80
Alkalinity (mg/L)	198.00	101.67
Total suspended solids (mg/L)	253.00	328.33

Table 1

This study compared the performance of the air injectors, driven by a 2-hp centrifugal pump, with a commercial microperforated air diffuser hose driven by a 2-hp blower. In the injector treatment, the injectors were positioned parallel to the direction of flow along the bottom of each tank wall. The diffuser hoses were cut into 10-cm lengths, connected by polyvinyl chloride pipe and placed every 1.5 m2 in the bottoms of the tanks.

Table 2. Zootechnical parameters of L. vannamei in a nursery phase biofloc system.

Parameter	Injectors	Air Hoses
Initial weight (g)	0.05	0.03
Final weight (g)	0.96	0.71
Survival (%)	98.8	80.7
Feed-conversion ratio	1.22	1.46

Table 2

The treatments were performed in duplicate. Water quality parameters were maintained within nor mal ranges for the culture of *L. vannamei* in both treatments (Table 1). The better growth and survival were observed in the injector treatment, with 0.96 g and 98.8 percent, respectively (Table 2).

# **Growout study**

An 86-day growout study was performed in two lined raceways to compare the air injectors to paddlewheel aerators. In the air injector system, 14 air injectors were positioned parallel to the direction of flow along the bottom of each raceway and driven by a 3-hp centrifugal pump. In the paddlewheel system, a 2-hp paddlewheel aerator was used.

Table 3. Water quality parameters in a growout phase biofloc system raising white shrimp.

Parameter	Injectors	Paddlewheel
Temperature (°C)	25.4	23.4
Dissolved oxygen (mg/L)	7.21	7.30
pH	7.78	7.84
Salinity	20.3	21.3
Total ammonia-nitrogen (mg/L)	0.24	0.28
Nitrite-nitrogen (mg/L)	4.12	1.88
Nitrate-nitrogen (mg/L)	11.60	18.00
Total suspended solids (mg/L)	458.00	662.00

Table 3

The raceways were stocked at 591 shrimp/m3 with 0.82-g L. vannamei juveniles. The raceways were each filled to 237 m3 with a mixture of sand-filtered seawater (185 m3, 28-ppt salinity), groundwater (52 m3, 5 ppt) and bioflocrich water (2 percent of the total volume) from a nursery study. Water quality parameters were maintained within normal ranges for the culture of L. vannamei (Table 3).

### Table 4

The shrimp in the injector treatment showed final average weights of 9.42 g, while those whose raceways received paddlewheel aeration reflected weights of 8.55 g. At 1.94, feed-conversion ratios were much better than the 3.95 recorded for the paddlewheel treatment.

However, the survival rate in the air injector trial was 73 percent versus 51 percent in the paddlewheel trial. Apparently the paddlewheel physically killed the shrimp, as during the trial, dead shrimp were found close to the aerator. This significantly affected final productivity and biomass values.

In this first trial with air injectors in biofloc systems, the authors compared the aeration performance of the injectors with that of the diffuser tubes and paddlewheel aerators typically used in southern Brazil.

At 0.71 and 0.64 g, respectively, the weekly growth rates for the injector and paddlewheel treatments were similar to those reported in super-intensive systems in Southern Brazil, but it is important to highlight that at the end of culture, low temperatures near 20 degrees-C were recorded.

# **Perspectives**

In this first trial with air injectors and bioflocs, the results were encouraging, as productivity exceeded 4 kg/m3. On the other hand, paddlewheel aerators may be a better choice for intensive systems in ponds and apparently are not intended for super-intensive raceway systems. In general, the detected temperatures were 2 degrees-C higher in the air injector system. This can be explained by the difference in injecting air into the bottom of the tanks versus casting water from the surface into the air.

#### **Authors**

• Dariano Krummenauer, Ph.D.

Dariano Krummenauer, Ph.D.

Laboratório de Carcinocultura Instituto de Oceanografia Universidade Federal do Rio Grande Rio Grande (R.S.), 96201-900, Brazil

[109,111,99,46,108,105,97,109,103,64,107,111,110,97,105,114,97,100]

Bob Advent

**Bob Advent** 

Advent Environmental Systems LLC Lake Mary, Florida, USA Carlos Gaona, Ph.D.; Geraldo Foes, Ph.D.; Aline Bezerra, M.S.; Alessandro Cardozo, Ph.D.; Luis
• Poersch, Ph.D.; Wilson Wasielesky Jr., Ph.D.

Carlos Gaona, Ph.D.; Geraldo Foes, Ph.D.; Aline Bezerra, M.S.; Alessandro Cardozo, Ph.D.; Luis Poersch, Ph.D.; Wilson Wasielesky Jr., Ph.D.

Laboratório de Carcinocultura Instituto de Oceanografia Universidade Federal do Rio Grande

#### Share

- Share via Email
- **Y** Share on Twitter
- **f** Share on Facebook
- in Share on LinkedIn

### **Tagged With**

biofloc air injector raceway centrifugal pump water quality Dariano Krummenauer Bob Advent

#### **Related Posts**

Aquafeeds

## The Bottom Line: Integrating feed manufacturing into fish or shrimp production operations

Incorporating feed manufacturing into an aquaculture business offers greater and more efficient product access, control and logistics, as well as additional profit potential. A feasibility study that plans for growth and includes both a business plan and economic analysis is a must.

**Innovation & Investment** 

## **Estimating biofilter size for RAS systems**

Controlling total ammonia-nitrogen (TAN) concentrations is the primary concern when sizing a biofilter for use in a recirculating aquaculture system. Sizing decisions are best based on previous experience with a given biofilter media in a specific biofilter configuration.

Responsibility

## Calcium and magnesium use in aquaculture

Aquatic plants and animals get the essential nutrients calcium and magnesium from water and food. Calcium concentrations impact the hydration and development of eggs in a hatchery, where calcium carbonate precipitation can be troublesome.

Health & Welfare

## **Building a better shrimp nursery, part 2**

Search **Q** 

Shrimp nursery systems offer the shrimp industry an important opportunity to increase profits, by helping produce strong, healthy and uniform juveniles with significant potential for compensatory growth after their transfer for final pond grow-out. Biofloc technology and water quality are critical components of shrimp nursery systems.

### **About The Advocate**

The Responsible Seafood Advocate supports the Global Seafood Alliance's (GSA) mission to advance responsible seafood practices through education, advocacy and third-party assurances.

### **Learn More**



### **Advertising Opportunities**

2022 Media & Events Kit

### Categories

Aquafeeds > Health & Welfare > From Our Sponsors > Innovation & Investment Innovation & Investment > Intelligence > Responsibility > Fisheries > Artículos en Español >

#### Don't Miss an Article

#### Featured

- Health & Welfare An update on vibriosis, the major bacterial disease shrimp farmers face
- Intelligence A seat at the table: Fed By Blue team says aquaculture needs a stronger voice
- Responsibility Quantifying habitat provisioning at macroalgae cultivation locations

### **Popular Tags**

All Tags	~
,agc	

#### Recent

- Fisheries Second Test: Another filler for the fisheries category
- Fisheries Test: This is filler for the fisheries Category
- Aquafeeds Test Article
- Responsibility Study: Climate change will shuffle marine ecosystems in unexpected ways as ocean temperature warms
- Health & Welfare Indian shrimp researchers earn a patent for WSSV diagnostic tool



About

- <u>Membership</u>
- Resources
- Best Aquaculture Practices (BAP)
- Best Seafood Practices (BSP)
- GOAL Events
- Advocate Magazine
- Aquademia Podcast
- Blog
- Contact

## Stay up to date with GSA

- 0

Copyright © 2024 Global Seafood Alliance All rights reserved.

**Privacy** 

Terms of Use

Glossary