



ALLIANCE™

[.https://gsa.rakadev.com](https://gsa.rakadev.com)**Responsible
Seafood**
ADVOCATEHealth &
Welfare

By-the-numbers nutritional bioenergetics for optimal feeding

1 May 2008

By Dr. Ingrid Lupatsch

Feed formulation, feeding regimes based on nutritional bioenergetics



The author developed a formulaic approach to determining energy and protein requirements for gilthead sea bream (left), white grouper (right) and two other species.

The successful culture of many fish species depends on formulated feeds containing sufficient levels of energy and protein to sustain animal growth. In addition, it is important to minimize feed waste that is costly for farmers and impacts the environment.

Aquaculture involves a great variety of species, and a general approach is needed to define energy and protein requirements so that each targeted species can realize its full growth and economic potential. Feed formulation and feeding regimes can be effectively based upon principles of nutritional bioenergetics.

The author recently carried out a study to provide a better understanding of growth and the underlying principles involved for several fish species. It focused on commercial species common to the Mediterranean area: gilthead sea bream (*Sparus aurata*), European sea bass (*Dicentrarchus labrax*), white grouper, (*Epinephelus aeneus*) and Asian sea bass or barramundi (*Lates calcarifer*).

Assumptions, calculations

A general assumption is that the energy and protein requirements of growing fish are the sum of the needs for maintenance plus growth. The requirement for maintenance, mainly a function of the size of the fish and water temperature, is proportional to the metabolic body weight in the following equation.

$$\text{Maintenance Needs} = a \times \text{Body Weight (kg)}^b$$

Where *a* is a constant, characteristic of a certain fish species at a set temperature, and *b* is the exponent of the metabolic body weight, which in fish is 0.80.

The requirement for growth is dependent on the amount and composition of the weight gain, including the cost of energy to deposit the new growth. The following equation shows how daily energy requirements can be quantified.

$$\text{Digestible Energy Needs (kJ)} = a \times \text{Body Weight (kg)}^{0.80} + c \times \text{Energy Gain (kJ)}$$

Where *c* is the cost of production in units of dietary energy to deposit energy as growth.

The same approach is used for the quantification of protein, except for the use of a different *b* exponent of 0.70 for metabolic body weight.

$$\text{Digestible Protein Needs (g)} = a \times \text{body weight (kg)}^{0.70} + c \times \text{Protein Gain (g)}$$

Where *c* is the cost of production in units of dietary protein to deposit protein as growth.

The significance of this approach is that protein and energy needs are expressed primarily in terms of absolute demand per fish body mass and anticipated weight gain.

Trial: maintenance requirements

To determine the maintenance requirement and efficiency for growth, fish were fed increasing levels of a feed with a known digestible energy (D.E.) and digestible protein (D.P.) content. Feeding levels included groups that received no feed up to maximum voluntary intake. Energy and protein gain were

determined by comparative carcass analyses. All the trials were done at an average temperature of 27 degrees-C.

Fig. 1 shows that the relationship between daily D.E. consumed and energy retained are linear. During starvation, the fish would lose energy as expected. The D.E. requirement for maintenance with no energy gain or loss can be determined for each species where the y axis value is 0.




Fig. 1: Energy requirements of several fish species for maintenance and growth at 27 degrees C.

The slope of the lines in Fig. 1 (approximately 0.67), regarded as the efficiency of energy utilization for growth deposition, is very similar for all species. The reciprocal value $c = 1/0.67$ is a measure for the “cost of production” in units of D.E. (kJ) to deposit 1 kJ as growth. The maintenance requirement for protein can be obtained in a similar manner by examining the relationship between the digestible protein consumed and protein retained.

Trial: growth requirements

The missing part of the equation – growth – is dependent on the amount of the daily weight gain and the composition of this gain. To assess the growth potential of a species over a full production cycle, the author monitored the growth rates of fish from 1 to 450 g at different water temperatures. Fish were fed to apparent satiation, and daily weight gain was determined by periodic weighing of the animals every two weeks. All the trials were done in full-strength seawater at ambient temperatures ranging 20 to 28 degrees-C.

The following equations were developed to express daily weight gain as a function of body weight and water temperature for the study fish.

$$\text{Grouper Weight Gain (g/day)} = 0.062 \times \text{Body Weight (g)}^{0.558} \times e^{0.030 \times \text{Temp}}$$

$$\text{Barramundi Weight Gain (g/day)} = 0.0051 \times \text{Body Weight (g)}^{0.508} \times e^{0.135 \times \text{Temp}}$$

$$\text{Seabream Weight Gain (g/day)} = 0.024 \times \text{Body Weight (g)}^{0.514} \times e^{0.060 \times \text{Temp}}$$

$$\text{Seabass Weight Gain (g/day)} = 0.020 \times \text{Body Weight (g)}^{0.517} \times e^{0.065 \times \text{Temp}}$$

Fig. 2 shows the daily weight gain in relation to body size at an average temperature of 27 degrees-C.




Fig. 2: Daily weight gain of fish at an average temperature of 27 degrees-C.

Composition of weight gain

The composition of the weight gain is an additional factor in determining energy and protein requirements. When measuring the whole body composition of fish at increasing sizes, each unit of weight gain is assumed to equal the body composition at a certain size. As pictured in Fig. 3, energy concentrations change with increasing fish weights and show different values typical for a species.

Fig. 3: Energy content of fish in relation to fish weight.

The whole body protein content is relatively constant regardless of size and fish species – on average, 169, 170, 175 and 171 mg per gram for grouper, barramundi, sea bream and sea bass, respectively. The following equations describe the change of energy density with fish size.

Practical applications

Using the parameters obtained, the daily energy and protein requirements can thus be determined. The following example is a comparison between grouper and seabream calculated for a water temperature of 27 degrees-C.

According to the growth prediction, a 300-g grouper has the potential to grow 3.36 grams per day. The energy content of the fish is 6.9 kJ per gram and protein content 169 mg per gram. Thus the anticipated gain would be 23.2 kJ and 0.57 grams in units of energy and protein, respectively.

Including the cost to produce this gain and the maintenance requirement, the total demand for D.E. is 53.8 kJ/day and for D.P. is 1.29 grams per day. As a result, the 300-gram grouper would have to consume 3.16 grams feed per day containing 17 MJ D.E. per kg and 409 g D.P. per kg feed, resulting in a feed-conversion ratio of 0.94.

On the other hand, a 300-gram sea bream would only grow 2.28 grams per day. The energy content of the gain is considerably higher (10.3 kJ per gram), as are maintenance requirements. Therefore, total requirements can be calculated for D.E. as 62.70 kJ per day and for D.P. as 1.18 grams per day.

As the ratio of dietary D.E. to D.P. is higher for sea-bream than grouper, the sea bream would have to consume 3.69 g feed containing 17 MJ D.E. and 320 g D.P. per kg feed, resulting in a feed-conversion ratio of 1.62.

Energy, protein calculations

Energy and protein requirements for every fish species can be calculated and adapted to changing conditions for the duration of a growth period (Table 1). Using this approach to quantify energy and protein demands, it is thus possible to estimate the biological and economical efficiency of different feeds and culture systems.

(Editor's Note: This article was originally published in the May/June 2008 print edition of the Global Aquaculture Advocate.)

Author



DR. INGRID LUPATSCH

Aquatic Animal Nutrition
Centre for Sustainable Aquaculture Research
Swansea University, Singleton Park
Swansea SA2 8PP, United Kingdom

i.lupatsch@swansea.ac.uk (<mailto:i.lupatsch@swansea.ac.uk>)

Copyright © 2023 Global Seafood Alliance

All rights reserved.