



- [GOAL Events](#)
- [Advocate Magazine](#)
- [Aquademia Podcast](#)
- [Blog](#)
- [Contact](#)
-
-
-
-
-
- [Log In](#)



- [About](#)
 - [Who We Are](#)
 - [Our History](#)
 - [Our Team](#)
 - [Sustainable Development Goals](#)
 - [Careers](#)
- [Membership](#)
 - [Overview](#)
 - [Our Members](#)
 - [Corporate Membership](#)
- [Resources](#)
- [Certification](#)
 - [Best Aquaculture Practices](#)
 - [Best Seafood Practices](#)

Search...



[Log In](#)

- [About](#)
 - [Who We Are](#)
 - [Our History](#)
 - [Our Team](#)
 - [Sustainable Development Goals](#)
 - [Careers](#)
- [Membership](#)
 - [Overview](#)
 - [Our Members](#)
 - [Corporate Membership](#)
- [Resources](#)
- [Certification](#)
 - [Best Aquaculture Practices](#)
 - [Best Seafood Practices](#)
- [GOAL Events](#)
- [Advocate Magazine](#)
- [Aquademia Podcast](#)
- [Blog](#)
- [Contact](#)



Health & Welfare
Health & Welfare

Captive reproduction, larviculture of Florida pompano



1 June 2005 Charles R. Weirich, Ph.D. Kenneth Riley, M.S. Megan Davis, Ph.D.



Trials indicate potential for mass production of feedstock



Florida pompano (*Trachinotus carolinus*).

Considered one of the finest marine table fish, the Florida pompano (*Trachinotus carolinus*) commands a significantly higher price than many other United States marine and fresh water finfish species. In 2003, the average wholesale price for whole pampano was \$7.50 per kilogram, and depending on the time of year and availability, pompano fillets can fetch \$35 per kilogram.

Recreational and commercial pompano fisheries exist along the south Atlantic and Gulf coasts of the United States, with 83 to 92 percent of landings occurring in Florida. Because pompano landings have never been large, demand has consistently exceeded supply.

Early research

Significant interest in the culture of pompano developed in the United States during the 1960s and 1970s. Initial research established that pampano exhibit rapid growth and tolerance to various culture conditions and systems. However, attempts to culture the fish on a commercial scale were largely unsuccessful.

As with other marine finfish, the availability of a sustainable supply of pompano seedstock limited culture operations. Aside from pioneering work conducted by Frank Hoff and associates, who bred pampano using hormonal induction and reared larvae with limited success in the early 1970s, no published research exists with respect to pompano reproduction and larviculture.

To further address this topic, the authors conducted preliminary trials in 2004 to evaluate captive reproduction of pampano via hormonal induction and to develop basic procedures for larval production.

Broodstock acquisition, feed training

Broodstock were acquired from local commercial fishermen during the fall and winter of 2003 to 2004. Fish were collected in the Indian River Lagoon on the east coast of Florida, USA. Three groups of 20 fish each were transported to the United States Department of Agriculture Agricultural Research Service (USDA-ARS) and Harbor Branch Oceanographic Institute (HBOI) aquaculture facilities in Fort Pierce, Florida, USA.

The fish were placed with an approximate sex ratio of 1:1 in 12,000-liter recirculating tanks with a natural photoperiod and water at 19 to 25 degrees-C and 28 to 30 grams per liter salinity. Over the course of two months, the fish were given treatments to eradicate protozoan and trematode parasites.

During the quarantine period, the fish were feed-trained. Peeled shrimp and cut squid were offered for the first two weeks after capture, then a moist diet mixed with peeled shrimp was fed for one week. At this point, the fish were not fed for five days. The moist diet was fed alone for one week, followed by

the moist diet in combination with a sinking pelleted diet for a week, until the pelleted diet was fed alone.

Spawning trials

In April 2004, the weight and sex of all fish were determined and transponder tags were injected for future identification. The mean weights of the males and females were 0.7 and 1.3 kilogram, respectively. Each group of 20 fish was placed back into its respective holding tank and chemical treatments were discontinued.

In May, fish were sampled to determine gonadal condition. The males were subjected to abdominal compression to detect the presence of milt, while oocyte samples were obtained from females through catheterization and examined under a microscope. Fourteen males possessed running milt, and the mean oocyte diameter of 16 females was 400 µ or greater.

Each fish was injected with a 75-µg gonadotropin-releasing hormone analog (GnRHa) pellet. Eight females and seven males were then placed into each of two 12,000-liter recirculating tanks equipped with side-mounted egg collectors to initiate spawning trial 1.

Subsequent trials were conducted in June and July. In trial 2, four of the eight females and all of the males in one tank were injected with GnRHa. In trial 3, six of the eight females and all of the males in the other tank were subjected to the hormonal treatment.

In each trial, spawning occurred in the early morning hours approximately 36 hours after GnRHa injection. Both floating and sinking eggs were enumerated by volume, and the fertilization rate of the floating eggs was determined after each spawning event. Fertilized eggs were approximately 1 mm in diameter with a single oil globule.

In the first and second trials, spawning occurred on five of six consecutive days with a total of 2.8 million and 0.9 million eggs collected, respectively (Table 1). In the third trial, spawning occurred on two of six consecutive days with a total of 0.8 million eggs collected.

Weirich, Results from pompano spawning trials, Table 1

Trial	Total Eggs	Floating Eggs	Fertilization	Eggs per kilogram Female	Eggs per Female
1	2.8 million	460,000	86.8%	131,500	174,200
2	0.9 million	368,000	71.3%	109,600	230,300
3	0.8 million	152,000	87.3%	88,000	123,000

Table 1. Results from pompano spawning trials. Fertilization rates represent the mean values of all spawning events within each trial.

Although considerable variation existed among individual spawns with respect to the fertilization rate of floating eggs, overall fertilization during each trial was consistent and relatively high compared to that observed in previous work with pompano by Hoff and associates. The fecundity of the female pompano was also rather consistent between trials and comparable to that of other subtropical marine finfish species.

Larviculture

After collection and sampling, floating eggs from each spawning event were placed into 100-liter circular tanks or 450-liter rectangular troughs at a density of 50 to 100 eggs per milliliter. Temperature was maintained at 24 to 25 degrees-C and salinity at 34 to 35 grams per liter. Fourteen hours of light and 10 hours of darkness were provided daily with fluorescent lighting.

Hatching occurred in 30-34 hours. The hatching rate of the incubated eggs was not determined. However, in concurrent small-scale studies conducted to determine the effects of copper, ammonia, nitrite, and salinity on eggs and larvae, hatching rates ranged from 20-90 percent between spawns. Unhatched eggs and egg debris were siphoned from the culture units and paper towels were used to remove surface oils. Aeration during incubation and initial larval rearing was maintained at a minimal rate to minimize injury.

Larvae began feeding approximately 72 hours posthatch, which corresponded to the time of complete yolk absorption. As shown in Fig. 1, one day after hatching, *Nannochloropsis* algal paste was added to culture tanks at approximately 30 million cells per milligram to achieve green water conditions, which were maintained for 11 days. Two days after hatching, enriched rotifers were added three times daily at a rate of 1 to 3 individuals per milligram.




Fig. 1: Feeding regime for pompano larviculture.

At seven days, instar artemia were added in addition to rotifers at 1 to 2 individuals per milligram. At 11 days, rotifer feeding was discontinued, and at 12 days, enriched artemia metanauplii were fed at a rate of 1 to 2 individuals per milligram. Micro-particulate diets were first offered at 18 days after hatching. Active consumption began at approximately 23 days, near the onset of metamorphosis, and artemia feeding was discontinued.

Larval survival

Larval survival was highly variable between spawns, with mean survival below 5 percent. Significant mortalities were generally confined to the periods of seven to nine days and 22 to 24 days post-hatch, with the former resulting in the bulk of larval losses.

The early-stage mortalities may have been due to inadequate nutrition, suboptimal light levels, physiological anomalies, bacterial pathogens, or a combination thereof. Latter-stage mortalities were likely due to physiological changes associated with metamorphosis. While no direct cannibalism was observed, postmetamorphic larvae occasionally chased premetamorphic larvae.

At 18 to 22 days post-hatch, pre-metamorphic larvae were noticeably darker in coloration than the post-metamorphic larvae. Postmetamorphic larvae immediately exhibited schooling and swimming behaviors identical to those of older juveniles and adult pompano.

Further research

The 2004 trials were cut short due to the passage of two hurricanes. Since then, considerable renovation of the USDA-ARS and HBOI facilities has occurred. In addition, a five-year USDA-ARS project plan calls for trials to be conducted this year to refine pompano spawning procedures with the goal of attaining year-round natural spawning. Further efforts will focus on improving larviculture methods for increased survival and production.

(Editor's Note: This article was originally published in the June 2005 print edition of the Global Aquaculture Advocate.)

Now that you've finished reading the article ...


... we hope you'll consider supporting our mission to document the evolution of the global aquaculture industry and share our vast network of contributors' expansive knowledge every week.

By becoming a Global Seafood Alliance member, you're ensuring that all of the pre-competitive work we do through member benefits, resources and events can continue. Individual membership costs just \$50 a year.

Not a GSA member? Join us.

[Support GSA and Become a Member](#)

Authors


-  Charles R. Weirich, Ph.D.

Charles R. Weirich, Ph.D.

United States Department of Agriculture
Agricultural Research Service

Sustainable Marine Aquaculture Systems Program
Fort Pierce, Florida 34946 USA

[118,111,103,46,97,100,115,117,46,115,114,97,46,97,112,115,64,104,99,105,114,105,101,119,99]

-  Kenneth Riley, M.S.

Kenneth Riley, M.S.





Harbor Branch Oceanographic Institute
Aquaculture Division
Fort Pierce, Florida, USA

-  Megan Davis, Ph.D.

Megan Davis, Ph.D.

Harbor Branch Oceanographic Institute
Aquaculture Division
Fort Pierce, Florida, USA

Share

-  [Share via Email](#)
-  [Share on Twitter](#)
-  [Share on Facebook](#)
-  [Share on LinkedIn](#)

Tagged With

[larviculture](#) [Florida pompano](#) [Megan Davis](#) [Charles R. Weirich](#) [Kenneth Riley](#)

Related Posts

Responsibility

[Haiti's hatchery of hope](#)

Built on a dream of feeding some of the world's poorest and most vulnerable people, a charity-built tilapia hatchery in Haiti now learns to stand on its own.

Innovation & Investment

[Artemia, the 'magic powder' fueling a multi-billion-dollar industry](#)

Artemia, microscopic brine shrimp used as feed in hatcheries, are the unsung heroes of aquaculture. Experts say artemia is still inspiring innovation more than 50 years after initial commercialization. These creatures are much more than Sea-Monkeys.

Aquafeeds

[Aquaculture Exchange: Rick Barrows](#)

After 14 years with the USDA's Agricultural Research Service, Rick Barrows talks about the importance of finding 'complete' and commercially viable alternative sources of omega-3 fatty acids and continuing innovation in the aquafeed sector.

Health & Welfare

[Pompano reared to market size in RAS](#)

In a trial, rearing density affected the mean weight and weight gain of pompano, although differences between treatments became apparent toward the end of the study.

About The Advocate

The Responsible Seafood Advocate supports the Global Seafood Alliance's (GSA) mission to advance responsible seafood practices through education, advocacy and third-party assurances.

[Learn More](#)

Search Responsible Seafood Advocate

Search 

Advertising Opportunities

[2022 Media & Events Kit](#)

Categories

[Aquafeeds](#) > [Health & Welfare](#) [Health & Welfare](#) > [From Our Sponsors](#) > [Innovation & Investment](#) > [Intelligence](#) > [Responsibility](#) > [Fisheries](#) > [Artículos en Español](#) >

Don't Miss an Article

Featured

- [Health & Welfare](#) [An update on vibriosis, the major bacterial disease shrimp farmers face](#)
- [Intelligence](#) [A seat at the table: Fed By Blue team says aquaculture needs a stronger voice](#)
- [Responsibility](#) [Quantifying habitat provisioning at macroalgae cultivation locations](#)

Popular Tags



Recent

- [Fisheries](#) [Second Test: Another filler for the fisheries category](#)
- [Fisheries](#) [Test: This is filler for the fisheries Category](#)
- [Aquafeeds](#) [Test Article](#)
- [Responsibility](#) [Study: Climate change will shuffle marine ecosystems in unexpected ways as ocean temperature warms](#)
- [Health & Welfare](#) [Indian shrimp researchers earn a patent for WSSV diagnostic tool](#)



- [About](#)
- [Membership](#)
- [Resources](#)
- [Best Aquaculture Practices \(BAP\)](#)
- [Best Seafood Practices \(BSP\)](#)
- [GOAL Events](#)
- [Advocate Magazine](#)
- [Aquademia Podcast](#)
- [Blog](#)
- [Contact](#)

Stay up to date with GSA

- 
- 
- 
- 
- 

Copyright © 2024 Global Seafood Alliance
All rights reserved.

[Privacy](#)

[Terms of Use](#)

[Glossary](#)