Global Seafood Alliance Logo

- GOAL Events
- <u>Advocate Magazine</u>
- <u>Aquademia Podcast</u>
- <u>Blog</u>
- <u>Contact</u>
- 0
- f
- 🗙
- in
- 🕨
- <u>Log In</u>
- \square
 - <u>About</u>
 - Who We Are
 - <u>Our History</u>
 - <u>Our Team</u>
 - Sustainable Development Goals
 - <u>Careers</u>
 - <u>Membership</u>
 - <u>Overview</u>
 - <u>Our Members</u>
 - Corporate Membership
 - <u>Resources</u>
 - <u>Certification</u>
 - Best Aquaculture Practices
 - Best Seafood Practices

Search...

Q

- <u>Log In</u>
 - <u>About</u>
 - <u>Who We Are</u>
 - <u>Our History</u>
 - Our Team
 - Sustainable Development Goals
 - <u>Careers</u>
 - <u>Membership</u>
 - <u>Overview</u>
 - Our Members
 - Corporate Membership
 - <u>Resources</u>
 - <u>Certification</u>
 - <u>Best Aquaculture Practices</u>
 - Best Seafood Practices
 - GOAL Events
 - <u>Advocate Magazine</u>

- <u>Aquademia Podcast</u>
- <u>Blog</u>
- Contact



Health & Welfare Health & Welfare

CART regulates food intake in channel catfish

Responsible Seafood Advocate logo 1 July 2008 Yasuhiro Kobayashi, Ph.D. Brian C. Peterson, Ph.D. Geoffrey C. Waldbieser, Ph.D.

Understanding gene functions to improve production efficiency



Fast-growing lines of channel catfish could pass on their superior performance through breeding programs.

Recent high feed costs have caused some channel catfish farmers to reduce feeding frequency or use less-expensive protein sources in feed. Although they save money, these changes can result in slower growth using standard practices.

Another method to help farmers reduce their feed costs is the use of catfish that are more efficient converters of feed to muscle. Fish with superior feed conversion must be identified and lines developed from these superior individuals.

The traditional method of selecting broodstock that are efficient consumers of feed involves lengthy, expensive and labor-intensive studies involving several lines of catfish families over several generations. However, identifying feed-efficient individuals through DNA analysis could be cheaper and require fewer lines of fish.

The first step in the analysis is to understand what gene products are involved in the regulation of feed intake and how these gene products function. For example, what genes are involved in signaling hunger, and what genes are involved in telling the fish to stop eating?

Feed intake regulation

Feed consumption in both mammals and fish is regulated by many hormones and chemicals called neurotransmitters, which are produced by nerve cells within the brain. One group of nerve cells produces neurotransmitters that increase feed intake, whereas another group of nerve cells produces neurotransmitters that inhibit feed intake.

One neurotransmitter that inhibits feed intake is cocaine-and amphetamine-regulated transcript (CART). Food deprivation reduces CART production in both fish and mice. Injection of CART into the brain decreases feed intake by inhibiting the action of neuropeptide Y (NPY), which stimulates feed intake. Both CART and NPY are neurotransmitter proteins that are encoded by their respective genes.

Previous research

The authors' studies at the Thad Cochran Warmwater Aquaculture Center in Stoneville, Mississippi, USA, have showed that lines of channel catfish selected for increased growth generally consumed more feed than catfish with slower growth.

The authors examined differences in NPY production between faster- and slower-growing catfish because NPY is known to be involved in feed intake, but there was no difference in NPY gene activity between faster-growing and slower-growing catfish. Based on these results, they speculated that differences in feed intake were caused by differences in the amount of neurotransmitters that decreased, rather than increased, feed intake.

If so, differences in the levels of a feeding inhibitor could lead to differences in growth between genetic lines of fish. More recent research has therefore focused on CART, the inhibitory neurotransmitter.

Current research



Neurotransmitters in the brains of fish are coded to either increase or inhibit feed intake.

The authors first used molecular biotechnology to identify the catfish CART gene sequence, which allowed them to deduce the amino acid sequence of the CART protein. The CART sequence was very similar to that of mammals

and other fish species. The authors then developed a molecular tool to measure changes in the levels of CART gene activity in the catfish brain after prolonged food deprivation.

A group of juvenile channel catfish was fasted for 30 days, while another group received feed daily during the same period. After the fasting period, brain tissue was collected from five fish in each group. The remaining fish in each group were fed for 15 additional days before their tissue was collected.

CART gene activity was compared among the four groups of fish. The study showed that 30 days of fasting reduced CART gene activity, and 15 days of refeeding restored CART gene activity to a level observed in the fish fed daily.

CART expression

Another study tested the hypothesis that differences in feed intake between lines of catfish with different growth rates were caused by differences in the amount of CART in their brains. The authors measured the feed intake and deficiency, weight gain and CART expression in the brains of fast- and slow-growing channel catfish.

Fish with faster growth consumed, on average, 97.1 grams more feed and weighed 80.7 grams more at the end of a seven-week period than fish with slower growth, while feed-conversion efficiency was similar for the two fish groups. The faster-growing fish also showed less CART gene activity in their brains than the slower-growing fish.

(Editor's Note: This article was originally published in the July/August 2008 print edition of the Global Aquaculture Advocate.*)*

Now that you've finished reading the article ...

... we hope you'll consider supporting our mission to document the evolution of the global aquaculture industry and share our vast network of contributors' expansive knowledge every week.

By becoming a Global Seafood Alliance member, you're ensuring that all of the pre-competitive work we do through member benefits, resources and events can continue. Individual membership costs just \$50 a year.

Not a GSA member? Join us.

Support GSA and Become a Member

Authors

Pasuhiro Kobayashi, Ph.D.

Yasuhiro Kobayashi, Ph.D.

Division of Biological and Physical Sciences Delta State University Cleveland, Mississippi 38733 USA

[117,100,101,46,101,116,97,116,115,97,116,108,101,100,64,105,104,115,97,121,97,98,111,107]

• 戻 Brian C. Peterson, Ph.D.

Brian C. Peterson, Ph.D.

Thad Cochran National Warmwater Aquaculture Center USDA -ARS Stoneville, Mississippi, USA

Geoffrey C. Waldbieser, Ph.D.

Thad Cochran National Warmwater Aquaculture Center USDA -ARS Stoneville, Mississippi, USA

Share

- Share via Email
- **Share on Twitter**
- **f** Share on Facebook
- in <u>Share on LinkedIn</u>

Tagged With

Geoffrey C. Waldbieser catfish CART Brian C. Peterson Yasuhiro Kobayashi

Related Posts

Innovation & Investment

Assessing coloration in channel catfish fillets

Because consumers look at color to gauge quality of catfish fillets, the authors developed a digital photography measurement method to assess yellowness.

Health & Welfare

Biofloc technology reduces common off-flavors in channel catfish

In studies that used biofloc systems to culture channel catfish, culture tanks were susceptible to episodes of geosmin and 2-methylisoborneol and subsequent bioaccumulation of off-flavors in catfish flesh.

Health & Welfare

<u>A study of Zoea-2 Syndrome in hatcheries in India, part 2</u>

Indian shrimp hatcheries have experienced larval mortality in the zoea-2 stage, with molt deterioration and resulting in heavy mortality. Authors considered biotic and abiotic factors. Part 2 describes results of their study.

Health & Welfare

<u>A study of Zoea-2 Syndrome in hatcheries in India, part 3</u>

6/20/2024

CART regulates food intake in channel catfish - Responsible Seafood Advocate

In this third and final part, authors present recommendations to help reduce the incidence of Zoea-2 Syndrome, which is not caused by any known infectious agents in P. vannamei hatcheries in India.

About The Advocate

The Responsible Seafood Advocate supports the Global Seafood Alliance's (GSA) mission to advance responsible seafood practices through education, advocacy and third-party assurances.

Learn More



Listen to the seafood industry's top podcast

Advertising Opportunities

2022 Media & Events Kit

Categories

Aquafeeds > Health & Welfare Health & Welfare > From Our Sponsors > Innovation & Investment > Intelligence > Responsibility > Fisheries > Artículos en Español >

Don't Miss an Article

Featured

- <u>Health & Welfare An update on vibriosis, the major bacterial disease shrimp farmers face</u>
- Uncategorized <u>A seat at the table: Fed By Blue team says aquaculture needs a stronger voice</u>
- <u>Responsibility Quantifying habitat provisioning at macroalgae cultivation locations</u>

Popular Tags

All Tags 🔹

Recent

- Fisheries Second Test: Another filler for the fisheries category
- Fisheries Test: This is filler for the fisheries Category
- Aquafeeds Test Article
- <u>Responsibility Study: Climate change will shuffle marine ecosystems in unexpected ways as ocean</u> <u>temperature warms</u>
- Health & Welfare Indian shrimp researchers earn a patent for WSSV diagnostic tool



Listen to the seafood industry's top podcast

• <u>About</u>

- <u>Membership</u>
- <u>Resources</u>
- Best Aquaculture Practices (BAP)
- Best Seafood Practices (BSP)
- GOAL Events
- Advocate Magazine
- <u>Aquademia Podcast</u>
- <u>Blog</u>
- <u>Contact</u>

Stay up to date with GSA

- Ø
- **f**
- 🗙
- in
- 🕨

Copyright © 2024 Global Seafood Alliance All rights reserved. <u>Privacy</u> <u>Terms of Use</u> <u>Glossary</u>