



- [GOAL Events](#)
- [Advocate Magazine](#)
- [Aquademia Podcast](#)
- [Blog](#)
- [Contact](#)
- 
- 
- 
- 
- 
- [Log In](#)



- [About](#)
  - [Who We Are](#)
  - [Our History](#)
  - [Our Team](#)
  - [Sustainable Development Goals](#)
  - [Careers](#)
- [Membership](#)
  - [Overview](#)
  - [Our Members](#)
  - [Corporate Membership](#)
- [Resources](#)
- [Certification](#)
  - [Best Aquaculture Practices](#)
  - [Best Seafood Practices](#)

Search...



[Log In](#)

- [About](#)
  - [Who We Are](#)
  - [Our History](#)
  - [Our Team](#)
  - [Sustainable Development Goals](#)
  - [Careers](#)
- [Membership](#)
  - [Overview](#)
  - [Our Members](#)
  - [Corporate Membership](#)
- [Resources](#)
- [Certification](#)
  - [Best Aquaculture Practices](#)
  - [Best Seafood Practices](#)
- [GOAL Events](#)
- [Advocate Magazine](#)
- [Aquademia Podcast](#)
- [Blog](#)
- [Contact](#)



Health & Welfare  
Health & Welfare


## Children’s hospital Boston makes zebrafish feeding regime breakthrough



2 January 2013 Jason Holland Christian Lawrence Nick King



## Feed formula with artemia replacement a more cost-effective solution



Children's Hospital Boston researchers examined whether the husbandry of zebrafish could be simplified to reduce the number of inputs without compromising embryo production. Photo courtesy of Isaac Adatto, Harvard University.

Despite being a diminutive species whose members weigh no more than 10 grams when fully grown, zebrafish (*Danio rerio*) are hugely important to a wide variety of scientific fields, including human disease research, developmental biology and genetics, environmental toxicology, drug screening, evolution and also aquaculture. There are an estimated 2,000 zebrafish laboratories in the United States alone, and a priority for every facility is to produce embryos of high quality that are suitable for the aforementioned research.

Care for these animals has traditionally been a very labor-intensive process, but a new study published by the Aquatic Resources Program at Children's Hospital Boston in Massachusetts, USA, identified a valid alternative requiring less labor. Results indicated that feeding juvenile zebrafish just once a day with a specially formulated feed that replaces the conventional *Artemia salina* nauplii can improve growth and offer a more cost-effective feeding solution with no adverse impact on reproductive performance.

### Mainstream model

The zebrafish model system has been around for a long time, but in the last two or three decades in particular, its usage has accelerated in a wide variety of scientific disciplines. In recent years, therefore, a number of attempts have been made to improve the understanding of the animal's key biological characteristics, particularly in the areas of nutrition, reproductive biology, larviculture, behavior and natural history.

The hospital report, "The Effects Of Feeding Frequency On Growth And Reproduction In Zebrafish," found these and other emerging data hold promise for use as a framework for developing more sophisticated, performance-driven management practices befitting the animals' status as a mainstream research model.

## Laborious inputs

While humans and fish are separated by more than 450 million years of evolution, there is still a great degree of genetic similarity between the two – as much as 75 percent, which is why biomedical researchers increasingly favor the zebrafish model. Other important features of zebrafish include their ease of culture and very short generation time with rapid development and maturity.

Zebrafish laboratories typically have thousands of 3-L aquariums, each housing a dozen fish. Each laboratory has to dedicate a staff member to repetitively service each aquarium by taking the water temperature or feeding the fish. Each visit to a tank is called an input.

The standard practice is to feed postlarval zebrafish two to five times daily, 365 days per year, usually with a combination of *Artemia salina* nauplii and a formulated rotifer diet. While many laboratories have employed this extremely labor-intensive feeding strategy for decades, its effects on performance had not been evaluated.

The Children's Hospital Boston researchers wanted to see if the husbandry could be simplified to reduce the number of inputs without compromising embryo production. These trials were the first formal evaluation of feeding practices in zebrafish beyond the juvenile stage.

## Feeding trials

Starting at 30 days post-fertilization (dpf), the researchers fed eight replicate groups of age- and strain-matched fish a commercial pelleted diet – Gemma Micro 300, manufactured by aquaculture feeds specialist Skretting. The groups received feed either once every other day or one time, three times or five times daily to achieve a total feed input of 5 percent body weight/day compared against a standard control of *Artemia salina* nauplii/metanauplii given three times daily to apparent satiation.

Once every other week, beginning at 30 days and ending at 150 dpf, each replicate tank was taken off the system and photographed from above with a digital camera. At least 10 randomly selected fish in each replicate were measured for fork length from the tip of the snout to the end of the middle of the caudal fin rays. Immediately after they were photographed, the fish in each replicate group were weighed.

Specific growth rates were calculated for each replicate tank for each sampling date. Aside from the first week, efforts were made to conduct growth measurements on a biweekly basis. At 150 dpf, the sex ratio of each treatment group was determined by counting the number of males and females in each tank.

## Positive results

Researchers found that in general, the fish fed the commercial diet at 5 percent body weight one, three or five times daily showed superior growth performance when compared with both the control and every-other-day feeding regimes, and did not compromise embryo quality.

They further learned that while increasing the feeding frequency beyond once daily did result in heavier females, on average, it did not significantly impact the rate of growth. This has potentially important implications for how zebrafish aquaculture facilities are managed, particularly with respect to the time and labor spent on feeding.

From start to finish, the study took eight months, which resulted in fish grown to sexual maturity. The researchers found the difference in the time it takes staff to feed a given facility containing hundreds or thousands of tanks once versus two or three times daily is significant.

The time savings gained could translate to reductions in operating costs and/or an increased dedication of personnel to performing higher-order management tasks, such as genetic management and pathogen control, which are often neglected because of staffing shortages.

## Artemia replacement

Another important aspect of the study was to establish if *Artemia* could be eliminated completely from the zebrafish diet because *Artemia* is a live animal and therefore quite variable – from its nutritional value and where it is harvested, to how it is enriched and packaged. *Artemia* also require a labor-heavy laboratory process of hatching and treating before they can be fed to study fish.

The superior growth of the fish in the one, three and five daily feeding groups may have also been attributed to the differences between the nutrient profiles of the commercial diet and the *Artemia* used in the study's controls.

In particular, newly hatched *Artemia* are known to be deficient in essential polyunsaturated fatty acids, particularly docosahexaenoic acid (DHA) and arachidonic acid (ARA), both of which are important for zebrafish growth and development. The commercial diet, meanwhile, had defined levels of DHA, EPA and other essential fatty acids.

## Robust embryos

Much of the research on zebrafish in the biomedical arena is on the production of embryos that are used as research subjects. The main aim of all zebrafish laboratories is to produce very robust embryos for use in scientific studies.

Yet no study prior to that undertaken by Children's Hospital Boston had extended experiments out to repeatedly measure reproductive performance over such a long period of 190 days. This was a unique aspect of the Boston study.

The hospital's spawning trials commenced at 76 dpf. Once every other week, two males and three females were randomly selected from each replicate tank and set up in a standard mating cage, yielding a total of 8 crosses/treatment and 40 total crosses/event. With nine events, there were a total of 360 crosses, 72 per treatment.

If crosses were successful, the embryos were collected, rinsed with system water, transferred to petri dishes and incubated overnight at 280 degrees-C. This was repeated once every other week for nine weeks until the end of the experiment at 191 dpf.

The fact that the commercial diet was able to give better growth and economics while eliminating artemia would be immaterial if reproductive performance was negatively impacted, and embryo production was compromised. However, the study found that in fish fed the formulated diet, there was no negative impact on reproductive performance as measured by sex ratio, crossing success, fecundity and fertility. Furthermore, the embryos produced during a spawning event were of superior quality.

*(Editor's Note: This article was originally published in the January/February 2013 print edition of the Global Aquaculture Advocate.)*

## Authors

-  Jason Holland

Jason Holland

Fish and Seafood Consultant  
London, United Kingdom

-  Christian Lawrence

Christian Lawrence

Aquatic Resources Program  
Children's Hospital Boston  
One Blackfan Circle  
Boston, Massachusetts 02115 USA

[117,100,101,46,100,114,97,118,114,97,104,46,104,99,116,46,115,114,101,100,110,101,64,101,99,110,101,114,119,97,108,99]

-  Nick King

Nick King

Technical Manager  
Marine Hatchery Feeds  
Skretting

## Share

-  [Share via Email](#)
-  [Share on Twitter](#)
-  [Share on Facebook](#)
-  [Share on LinkedIn](#)

## Tagged With

[Artemia](#) [EPA](#) [DHA](#) [feeding](#) [zebrafish](#) [nauplii](#) [Jason Holland](#) [Christian Lawrence](#) [embryos](#)

## Related Posts

### Intelligence

### [On GM foods, part 4: Let's examine our needs](#)

Scott Nichols concludes his four-part series about GM foods with a simple question about transgenic animals: Are they truly meeting a need? We need more food and we need better nutrition. Both desires should be achieved without increasing environmental impacts.

### Health & Welfare

### [A holistic management approach to EMS](#)

Early Mortality Syndrome has devastated farmed shrimp in Asia and Latin America. With better understanding of the pathogen and the development and improvement of novel strategies, shrimp farmers are now able to better manage the disease.

### Health & Welfare

## [Algae shows promise as alternative DHA source in rainbow trout diets](#)

A growth trial in Canada evaluated the use of algae biomass to increase the concentration of long-chain polyunsaturated fatty acids in the tissues of rainbow trout.

[Aquafeeds](#)

## [A look at phospholipids in aquafeeds](#)

Phospholipids are the major constituents of cell membranes and are vital to the normal function of every cell and organ. The inclusion of phospholipids in aquafeeds ensures increased growth, better survival and stress resistance, and prevention of skeletal deformities of larval and juvenile stages of fish and shellfish species.

### About The Advocate

The Responsible Seafood Advocate supports the Global Seafood Alliance's (GSA) mission to advance responsible seafood practices through education, advocacy and third-party assurances.

[Learn More](#)

Search Responsible Seafood Advocate



### Advertising Opportunities

[2022 Media & Events Kit](#)

### Categories

[Aquafeeds](#) > [Health & Welfare](#) [Health & Welfare](#) > [From Our Sponsors](#) > [Innovation & Investment](#) > [Intelligence](#) > [Responsibility](#) > [Fisheries](#) > [Artículos en Español](#) >

### Don't Miss an Article

#### Featured

- [Health & Welfare](#) [An update on vibriosis, the major bacterial disease shrimp farmers face](#)
- [Uncategorized](#) [A seat at the table: Fed By Blue team says aquaculture needs a stronger voice](#)
- [Responsibility](#) [Quantifying habitat provisioning at macroalgae cultivation locations](#)

Popular Tags

All Tags

Recent

- Fisheries [Second Test: Another filler for the fisheries category.](#)
- Fisheries [Test: This is filler for the fisheries Category.](#)
- [Aquafeeds Test Article](#)
- [Responsibility Study: Climate change will shuffle marine ecosystems in unexpected ways as ocean temperature warms](#)
- Health & Welfare [Indian shrimp researchers earn a patent for WSSV diagnostic tool](#)



- [About](#)
- [Membership](#)
- [Resources](#)
- [Best Aquaculture Practices \(BAP\)](#)
- [Best Seafood Practices \(BSP\)](#)
- [GOAL Events](#)
- [Advocate Magazine](#)
- [Aquademia Podcast](#)
- [Blog](#)
- [Contact](#)

Stay up to date with GSA

- 
- 
- 
- 
- 

Copyright © 2024 Global Seafood Alliance  
All rights reserved.  
[Privacy](#)  
[Terms of Use](#)  
[Glossary](#)