Global Seafood Alliance Logo

- GOAL Events
- Advocate Magazine
- Aquademia Podcast
- Blog
- Contact
- (O)

- Log In

- About
 - Who We Are
 - Our History
 - o Our Team
 - Sustainable Development Goals
 - Careers
- <u>Membership</u>
 - Overview
 - Our Members
 - Corporate Membership
- Resources
- Certification
 - Best Aquaculture Practices
 - Best Seafood Practices

Search...



Log In

- About
 - Who We Are
 - Our History
 - o Our Team
 - Sustainable Development Goals
 - Careers
- <u>Membership</u>
 - o <u>Overview</u>
 - Our Members
 - o Corporate Membership
- Resources
- **Certification**
 - Best Aquaculture Practices
 - Best Seafood Practices
- GOAL Events
- Advocate Magazine

- <u>Aquademia Podcast</u>
- Blog
- Contact




Compound microdiets for marine fish larvae: Key parameters

Responsible Seafood Advocate logo

1 August 2004 Carlos E. Medina-Reyna, Ph.D. Juan P. Lazo, Ph.D. R. Pedroza-Islas, Ph.D.



Authors recommend diet stability and palatability assays



Scanning electron microscopy photograph of a microdiet.

Production of live food for the rearing of larval marine fish constitutes a major operational cost and bottleneck in commercial marine fish culture. The variability and poor nutritional quality of live food are major concerns during production.

Research has been directed toward developing compound microdiets to replace live food to ameliorate this problem. Although extensive research has been performed during the last 25 years, studies to determine specific nutritional requirements such as amino acid and vitamin requirements are scarce and the formulation of adequate diets is still elusive.

Formulating complete diets

Larval performance on compound microdiets alone is, in general, inferior to that of live food. One question that arises from previous studies is whether the technological processes utilized to manufacture microdiets is adequate to emulate live food acceptability and digestibility. The authors recommend that diet research incorporate comprehensive evaluations that include diet stability and palatability assays, which are not usually performed for marine larvae diets.

Much work has been devoted to biological evaluation, such as bioassays that last longer than six weeks, but the only parameters evaluated in most feeding trials are growth and survival. The authors propose that several key parameters be investigated during the manufacturing of practical compound microdiets for first-feeding marine fish larvae.

Research process

The ideal compositions of compound microdiets are species-dependent, but common guidelines can be followed based on previous research (Table 1). The authors have modified the approach suggested by Yufera and coauthors in 2000, which included a three-step process: diet design, biological evaluation and digestibility evaluation. The following research strategy is proposed.

- Selection of ingredients. Use fresh by-products from marine organism processing plants (although they represent potential vectors for disease), fishmeal, and a protein hydrolyzate to provide intact proteins, peptides, and free amino acids, as well as highly unsaturated fatty acids (Table 2).
- Selection of binder/technology. Use a mixture of polysaccharides and proteins that are highly digestible. The application of combing fluidized bed dryer and microbound flake technology is suggested.
- Stability evaluation. Test the leaching rate of soluble protein, peptides, and/or amino acids. The authors evaluated soluble protein using a modified Bradford technique.
- Acceptability evaluation. Study ingestion rates with and without gentle aeration during five hours using larvae not fed for 24 hours. The authors evaluated animals at 16 days posthatch.
- *In vitro* digestibility. Compute using the following equation. Relative protein digestibility = (Δ Abs.280_{CMD} / Δ Abs.280_{Caseine}) x 100

Medina-Reyna, Recommended nutrient levels for compound microdiets, Table 1

Nutrient	-	g/100 g diet Maximum
Proteins	50	65
Free amino acids		10
Peptides		20
Intact protein	30	
Lipids	10	20
Triglycerides		5
Phospholipids	10	
22:6n-3	2	
20:5n-3	1	
20:4n-6	0.1	
DHA:EPA	2	
EPE:ARA	5	10
HUFAs n-3	3	5
Carbohydrates		10

Nutrient	0	g/100 g diet Maximum
Fiber		2
Ash		10
Vitamins	5	
Ascorbic acid	0.5	
Minerals	4	
Attractants	2	4

Table 1. Recommended nutrient levels for compound microdiets for marine fish larvae.

Medina-Reyna, Experimental diet formulation, Table 2

Ingredient	g/100 g diet	
Menhaden fishmeal	49.4	
CPSP 90	14.8	
Skipjack muscle	9.8	
Skipjack orbital oil	14.8	
Cod liver oil	4.9	
Carrageenan	0.62	
Mesquite gum	3.1	
Sodium alginate	0.6	
Whey protein concentrate	0.6	
Vitamin premix	0.1	
Mineral premix	0.2	
Proximate composition (%)		
Protein	67.4	
Lipid	21.3	
Dry matter	93	
m11 o m 1 11	1	

Table 2. Experimental diet formulation.

Experimental diet tested

In a study that applied the above stategy, the authors selected several ingredients available locally and combined them with fishmeal to provide a nutritionally balanced diet for marine fish larvae. Although the 67 percent protein content seems high, previous experience has shown lower ingestion rates with microdiets. The level was chosen to satisfy requirements using a nutrient-dense approach.

A binder mixture enhanced the mechanical properties of the dough during the flaking process, but pieces of fishmeal were not hydrated satisfactorily or homogenized as fresh ingredients. The combination of these factors resulted in a diet that was easy to prepare.

Although the ingredients were finely ground to 10-20 μ to produce a microdiet of 250 μ , the soluble proteinleaching rate was a satisfactory 0.5 percent per hour. Leaching of other nutrients should be evaluated to provide a more comprehensive evaluation of diet stability. The relative protein digestibility of 80.4 percent seemed adequate, but should be confirmed using larval digestive enzymes and ultimately correlated with *in vivo* assays.

Acceptability tests with California halibut larvae were initially performed in tanks without aeration. The compound microdiet tended to sink quickly in less than 30 minutes. When gentle aeration was provided, ingestion rates significantly increased, with even higher rates in the presence of microalgae. The behavior of the diet in the water column must be evaluated and optimized prior to biological evaluation trials.

(Editor's Note: This article was originally published in the August 2004 print edition of the Global Aquaculture Advocate.)

Now that you've finished reading the article ...

... we hope you'll consider supporting our mission to document the evolution of the global aquaculture industry and share our vast network of contributors' expansive knowledge every week.

By becoming a Global Seafood Alliance member, you're ensuring that all of the pre-competitive work we do through member benefits, resources and events can continue. Individual membership costs just \$50 a year.

Not a GSA member? Join us.

Support GSA and Become a Member

Authors

• Carlos E. Medina-Reyna, Ph.D.

Carlos E. Medina-Reyna, Ph.D.

Department of Aquaculture Centro de Investigación Cientifíca y de Educación Superior de Ensenada Km. 107 Carretera Tijuana-Ensenada Código Postal 22860 Apdo. Postal 2732 Ensenada, B.C., México

[120,109,46,101,115,101,99,105,99,64,97,110,105,100,101,109,99]

Juan P. Lazo, Ph.D.

Juan P. Lazo, Ph.D.

Department of Aquaculture Centro de Investigación Cientifíca y de Educación Superior de Ensenada Km. 107 Carretera Tijuana-Ensenada Código Postal 22860 Apdo. Postal 2732 Ensenada, B.C., México

R. Pedroza-Islas, Ph.D.

R. Pedroza-Islas, Ph.D.

Universidad Iberoamericana Santa Fe. México D.F. México

Share

- Share via Email
- **Y** Share on Twitter
- **f** Share on Facebook
- **in** Share on LinkedIn

Tagged With

marine fish R. Pedroza-Islas Juan P. Lazo Carlos E. Medina-Reyna key parameters

Related Posts

Health & Welfare

Protein hydrolysates in larval fish nutrition

Protein hydrolysates are currently used as partial substitutes for fishmeal in high-quality artificial diets since their inclusion in feeds promotes growth and survival.

Health & Welfare

Atlantic red porgy aquaculture

With high value in U.S. markets and its overfished status, red porgy appears to be a viable candidate for aquaculture in North America.

Health & Welfare

UNC-Wilmington conducts research on black sea bass

Studies at UNC-Wilmington found that growth and survival of black sea bass larvae were improved through supplementation of DHA and ARA in live feed.

Aquafeeds

Artemia replacement diets for penaeid shrimp

At shrimp hatcheries, artemia replacement diets are used to replace live artemia, but complete replacement in commercial conditions remains a challenge.

About The Advocate

The Responsible Seafood Advocate supports the Global Seafood Alliance's (GSA) mission to advance responsible seafood practices through education, advocacy and third-party assurances.

Learn More

Search Responsible Seafood Advocate | Search

Search **Q**



Advertising Opportunities

2022 Media & Events Kit

Categories

Aquafeeds > Health & Welfare Health & Welfare > From Our Sponsors > Innovation & Investment > Intelligence > Responsibility > Fisheries > Artículos en Español >

Don't Miss an Article

Featured

- Health & Welfare An update on vibriosis, the major bacterial disease shrimp farmers face
- <u>Intelligence A seat at the table: Fed By Blue team says aquaculture needs a stronger voice</u>
- Responsibility Quantifying habitat provisioning at macroalgae cultivation locations

Popular Tags

All Tags

Recent

- Fisheries Second Test: Another filler for the fisheries category
- Fisheries Test: This is filler for the fisheries Category
- Aguafeeds Test Article
- Responsibility Study: Climate change will shuffle marine ecosystems in unexpected ways as ocean temperature warms
- Health & Welfare Indian shrimp researchers earn a patent for WSSV diagnostic tool



- **About**
- **Membership**
- **Resources**
- Best Aquaculture Practices (BAP)
- Best Seafood Practices (BSP)
- **GOAL Events**
- **Advocate Magazine**
- Aquademia Podcast
- **Blog**
- **Contact**

Stay up to date with GSA

Copyright © 2024 Global Seafood Alliance All rights reserved.

Privacy Terms of Use Glossary