### Global Seafood Alliance Logo

- GOAL Events
- Advocate Magazine
- Aquademia Podcast
- Blog
- Contact
- (O)

- Log In

- About
  - Who We Are
  - Our History
  - o Our Team
  - Sustainable Development Goals
  - Careers
- <u>Membership</u>
  - Overview
  - Our Members
  - Corporate Membership
- Resources
- **Certification** 
  - Best Aquaculture Practices
  - Best Seafood Practices

Search...



# Log In

- About
  - Who We Are
  - Our History
  - o Our Team
  - Sustainable Development Goals
  - Careers
- <u>Membership</u>
  - o <u>Overview</u>
  - Our Members
  - o Corporate Membership
- Resources
- **Certification** 
  - Best Aquaculture Practices
  - Best Seafood Practices
- GOAL Events
- Advocate Magazine

- Aquademia Podcast
- Blog
- Contact



## Ecuador study shows wild postlarvae use down

Responsible Seafood Advocate logo

1 October 2002 Stanislaus Sonnenholzner, Ph.D. Laurence Massaut, Ph.D. Claude E. Boyd, Ph.D.



# Between 1995 and 2000, the use of wild PLs declined from 61 percent to 6 percent



Wild postlarvae are collected in scissor nets in shallow waters off the coast of Ecuador.

Over the past 30 years, Ecuador has developed a large shrimp-farming industry characterized by extensive and semi-intensive production systems. During the 1970s and 1980s, the industry relied almost entirely on wild postlarvae (PL). However, because of the natural variability in the supply of wild PL, a large shrimp hatchery industry also developed in Ecuador.

## Wild, hatchery postlarvae

Most farmers traditionally used wild PL when they were available and hatchery PL when the supply of wild PL was short. There is an obvious advantage to having a constant supply of PL so that ponds can be stocked in a timely manner following harvests.

Many observers of shrimp farming in Ecuador believe the shrimp farming industry moved toward greater reliance on hatchery PL during the past 10 to 12 years. Nevertheless, wild PL are still used in large numbers and environmentalists often criticize this practice as being ecologically degrading.

## Survey conducted

To assess the opinions of farm managers on PL performance in growout ponds and their preferred sources for larvae in the last five years of production, the National Center for Aquaculture and Marine Research (Centro Nacional de Acui-cultura e Investigaciones Marinas, CENAIM) conducted a survey by questionnaire in 2000. The work was conducted with the financial support of the Mac Arthur Fund, World Wildlife Fund, World Bank, and Network of Aquaculture Centers in Asia Pacific. Of more than 100 questionnaires distributed, only 11 were returned.

# Wild PL use declining

Pond records indicated a declining trend of wild PL stocking from 61 percent in 1995 to 6 percent in 2000, with 1998 the exception to this trend (Fig. 1). The percentage of ponds stocked with wild PL during this year, which was characterized by a strong El Niño southern oscillation (ENSO) event, increased to 69 percent. An increase of wild PL abundance during ENSO events has been documented for Ecuador. Higher supply and probably lower prices of wild PL likely contributed to the observed stocking preference in 1998.

### La Niña and WSSV




Fig. 1: Proportion of shrimp ponds stocked with wild and hatchery postlarvae.

The significant decline of wild PL stocked into growout ponds in 1999 and 2000 can be attributed to two factors: one climatological (La Niña), and the other epidemiological in nature (the appearance of White Spot Syndrome Virus).

Contrary to the above-normal seawater temperatures characteristic of ENSO events, La Niña brought cooler water and a shortage of wild seed to the Ecuadorian coast. The first cases of disease problems in Ecuador associated with WSSV were reported in April 1999. Since then, production in Ecuador has declined significantly.

Statistical reports of the Cámara Nacional de Acuicultura of Ecuador showed a 66 percent reduction in exports for the year 2000. In an attempt to exclude the virus and reduce the risk of infection, farmers adopted the practice of stocking hatchery PL in ponds. Only 3 and 6 percent of all ponds surveyed in this study were stocked with wild PL during 1999 and 2000, respectively.

## **Performance**

There is a general belief among farm managers that wild PL outperform hatchery PL during pond growout, but the survey showed similar yields (p > 0.05) and absolute growth rates between wild and hatchery PL. However, survival rates of wild PL were significantly greater (p > 0.05) than those of hatchery PL (Table 1).

# Sonnenholzner, Comparison of pond production stocked with hatchery, Table 1

Numbe Ponds	r Number Ponds	Growth (g/week)	Growth (g/week)	Survival (50 %)	Survival (50%)	Yield (lb/ha)	Yield (lb/ha)	_`	g Stocking (PL/m <sup>2</sup> )
Н	W	Н	W	Н	W	H	W	H	W
1995 37	57	0.78	0.77	37	57	1,188	1,526	12.1	9.2
1996 118	73	0.74	0.68	38	64	1,374	1,342	13.9	7.4
1997 202	70	0.68	0.75	39	70	1,467	1,704	15.9	9.3
1998 90	199	0.63	0.65	50	61	1,536	1,661	14.3	10.6
1999 290	10	0.64	0.62	36	56	1,208	2,168	14.5	14.5
2000 162	11	0.83	0.83	20	23	449	460	10.0	8.3

Table 1. Comparison of pond production stocked with hatchery (H) and wild (W) Litopenaeus vannamei postlarvae.

The survival rates reported for wild PL could be attributed to an underestimation of stocking densities, possibly due to both technical and social issues. PL counting methods are not standard and have a wide margin of error, usually favoring the farmer. And many managers and biologists take advantage of the need fishermen and middlemen have to sell their wild PL, setting purchase prices and practically assuring an overestimation of PL numbers in any particular batch. As a result, ponds are typically stocked with 10 to 30 percent more PL than are registered in the record books of the farms.

In contrast, hatcheries have better methods for estimating the number of PL in their tanks, and are in a better position to control the prices they charge farmers. As a result, counts of hatchery PL are probably more accurate.

## Motives for PL selection

In 2000, about 73 percent of managers preferred hatchery seed, 18 percent preferred wild seed, and one respondent did not have a preference for one source of PL over the other. To gain insight into their motives for PL selection, farmers were asked about the issues that contributed to their preference for PL.

Reported changes were related to economic factors such as price and availability of wild seed, and a desire for PL that were stronger and more resistant to diseases – achievable through genetic selection at integrated farm-hatchery programs. Environmental concerns such as wild larvae overfishing or by-catch mortality were not a major factor in selecting hatchery PL.

Disease resistance of PL was of low priority in the decision-making process when selecting hatchery seed. Surprisingly, wild PL performance was not considered an important factor. The timing of the survey may have affected this response, because wild PL were achieving survival similar to hatchery PL in WSSV-infected ponds. In the past, the common opinion was that wild PL were stronger and produced higher yields than hatchery PL.

### **Conclusion**

A small-scale survey of shrimp farms in Ecuador to assess the use and performance of wild and hatchery postlarvae showed that between 1995 and 2000, the use of wild PL declined from 61 percent to 6 percent, except for the El Niño year of 1998, which was characterized by an overabundance of wild seed.

Results also showed similar yields and growth rates between wild and hatchery PL, but the survival rates of wild PL were significantly greater than those of hatchery PL. The latter can probably be attributed to PL counting methods. Hatchery PL were preferred by 73 percent of the managers surveyed.

(Editor's Note: This article was originally published in the October 2002 print edition of the Global Aquaculture Advocate.)

# Now that you've finished reading the article ...

... we hope you'll consider supporting our mission to document the evolution of the global aquaculture industry and share our vast network of contributors' expansive knowledge every week.

By becoming a Global Seafood Alliance member, you're ensuring that all of the pre-competitive work we do through member benefits, resources and events can continue. Individual membership costs just \$50 a year.

Not a GSA member? Join us.

Support GSA and Become a Member

#### Authors

Stanislaus Sonnenholzner, Ph.D.

Stanislaus Sonnenholzner, Ph.D.

Centro Nacional de Acuicultura e Investigaciones Marinas Campus Politécnico Casilla 09-01-4519 Guayaquil, Ecuador

[99,101,46,117,100,101,46,108,111,112,115,101,46,109,105,97,110,101,99,64,110,101,110,110,111,115,115]

Laurence Massaut, Ph.D.

Laurence Massaut, Ph.D.

Centro Nacional de Acuicultura e Investigaciones Marinas Campus Politécnico Casilla 09-01-4519 Guayaguil, Ecuador

Claude E. Boyd, Ph.D.

Claude E. Boyd, Ph.D.

Department of Fisheries and Allied Aquacultures Auburn University, Alabama, USA

#### Share

- Share via Email
- **Share on Twitter**
- **f** Share on Facebook
- in Share on LinkedIn

#### **Tagged With**

Claude E. Boyd Stanislaus Sonnenholzner Laurence Massaut wild postlarvae

#### **Related Posts**

Responsibility

## 'Cash-SIS' culture yields money crop, family food

In a study of pond polyculture, manipulation of species composition improved fish yield and corresponding income. Selling the whole production increased income 27 percent.

Health & Welfare

## **Dissolved oxygen dynamics**

Dissolved oxygen management is the most important requirement of aquaculture pond water quality. DO concentration below 3 mg/L is stressful to shrimp.

Intelligence

## Freshwater prawn farming in Brazil

Freshwater prawns are farmed in almost all states in Brazil, although the industry is concentrated in the southeast region, mainly the state of Espirito Santo.

Health & Welfare

## New nursery diet for shrimp performs well at super-high stocking densities

Search **Q** 

A shrimp diet developed in trials is the first dry nursery diet shown to ensure good performance at stocking densities up to 45,000 per square meter.

#### **About The Advocate**

The Responsible Seafood Advocate supports the Global Seafood Alliance's (GSA) mission to advance responsible seafood practices through education, advocacy and third-party assurances.

### Learn More



### **Advertising Opportunities**

2022 Media & Events Kit

#### Categories

Aquafeeds > Health & Welfare Health & Welfare > From Our Sponsors > Innovation & Investment > Intelligence > Responsibility > Fisheries > Artículos en Español >

#### Don't Miss an Article

#### Featured

- Health & Welfare An update on vibriosis, the major bacterial disease shrimp farmers face
- Intelligence A seat at the table: Fed By Blue team says aquaculture needs a stronger voice
- Responsibility Quantifying habitat provisioning at macroalgae cultivation locations

#### **Popular Tags**

All Tags	~
, iii lage	

#### Recent

- Fisheries Second Test: Another filler for the fisheries category
- Fisheries Test: This is filler for the fisheries Category
- Aquafeeds Test Article
- Responsibility Study: Climate change will shuffle marine ecosystems in unexpected ways as ocean temperature warms
- Health & Welfare Indian shrimp researchers earn a patent for WSSV diagnostic tool



**About** 

- <u>Membership</u>
- Resources
- Best Aquaculture Practices (BAP)
- Best Seafood Practices (BSP)
- GOAL Events
- Advocate Magazine
- Aquademia Podcast
- Blog
- Contact

### Stay up to date with GSA

- 0

Copyright © 2024 Global Seafood Alliance All rights reserved.

**Privacy** 

Terms of Use

Glossary