



ALLIANCE™

(<https://gsa.rakadev.com>).



**Responsible
Seafood**
ADVOCATE



Health &
Welfare

Effective management avoids gas bubble trauma

2 July 2012

By Claude E. Boyd, Ph.D.

Still water can remain undersaturated or supersaturated for several hours or days



Aerators that splash water into the air can lessen afternoon supersaturation with dissolved oxygen caused by a high rate of photosynthesis.

Atmospheric pressure results from the weight of atmospheric gases pressing down on the earth's surface. The atmospheric gases and their proportions are: nitrogen, 78.084 percent; oxygen, 20.946 percent; argon, 0.934 percent; carbon dioxide, 0.038 percent; water vapor and other gases, 0.036 percent.

According to Dalton's Law of Partial Pressures, each gas in the atmosphere exerts a pressure (partial pressure) in direct proportion to its percentage composition. For example, standard atmospheric pressure at sea level is 760 mm of mercury (Hg), and the partial pressure of nitrogen is 593.4 mm (760 mm x 0.78084)

Henry's Law states that each gas has a characteristic solubility, and its molecules will diffuse from the atmosphere into water until the pressure in water equals the partial pressure of the gas in the atmosphere. When this state is reached, the pressure of the gas in the atmosphere is at equilibrium with its pressure in water, and no further net exchange of its molecules occurs between the atmosphere and water.

At this stage, the water is said to be saturated with the gas. When the pressures of gases in the atmosphere and in the water are equal, the water is saturated with air.

Under certain conditions, water can contain either a lower or higher concentration of one or more gases than it should at equilibrium. When water is undersaturated with a gas, that gas enters the water from the atmosphere, and an equilibrium state is attained. Likewise, if water has more of a gas than it should – a state called supersaturation – the gas will diffuse from water to the atmosphere until equilibrium is reached.

Such diffusion does not occur rapidly in still water. Still water can remain undersaturated or supersaturated for several hours or days under certain conditions.

Saturation concentration

The saturation concentrations of gases and air vary with water temperature (Table 1). A sudden rise in temperature would result in temporary gas supersaturation, while a drop in temperature would cause temporary gas undersaturation.

Concentrations of air and individual gases in water sometimes are given as percentage saturation. For example, saturation with dissolved air at 20 degrees C is 25.06 mg/L (Table 1), but if water at this temperature contains 30.17 mg/L air, its percentage saturation with air is 120.4 percent $[(30.17/25.06) \times 100]$. Obviously, at equilibrium, the percentage saturation of a gas (or air) in water is 100 percent.

Boyd, Concentrations of individual gases, Table 1

Water Temp. (° C)	Concentration (mg/L) Nitrogen	Concentration (mg/L) Oxygen	Concentration (mg/L) Argon	Concentration (mg/L) Carbon Dioxide	Concentration (mg/L) Air
5	20.33	12.56	0.78	0.89	34.56
10	18.14	11.28	0.69	0.75	30.86
15	16.36	10.07	0.62	0.63	27.68
20	14.88	9.08	0.56	0.54	25.06
25	13.64	8.24	0.50	0.46	22.84
30	12.58	7.54	0.46	0.40	20.98
35	11.68	6.93	0.42	0.35	19.38
40	10.89	6.41	0.39	0.31	18.00

Table 1. Concentrations of individual gases and atmospheric air in freshwater at different temperatures and 760 mm Hg.

Supersaturation

There are several reasons why water becomes supersaturated with air. Natural warming of water in a pond or heating of water in hatcheries are common causes. Air leaks on the suction side of pumps or an improper submergence depth of pump intakes can cause gas supersaturation. Highly efficient submerged aerators have also been reported to cause gas supersaturation.

One of the best-known causes of gas supersaturation is entrainment of air bubbles when water falls over spillways of high dams. In cooler climates, water that infiltrates downward into aquifers in winter can be quite cold and contain a high concentration of air. In warm weather, water from wells in such aquifers tend to be supersaturated with air in respect to ambient air temperature.

Individual gases can be below or above saturation, with the most common example being dissolved oxygen. It is not unusual for surface waters in ponds to have dissolved-oxygen supersaturation of 200 to 300 percent during the afternoon because of photosynthesis. At night, photosynthesis stops, and respiration can cause dissolved-oxygen concentrations to fall to 50 percent or less of saturation.

Gas bubble trauma

Gases dissolve in the blood of fish, shrimp and other aquatic animals. Suppose animals are held in water at a certain temperature, and their blood equilibrates with gases in the water. Then suppose the water is suddenly warmed, resulting in gas supersaturation. The fish blood also will be supersaturated with gases, and gas bubbles can form in the blood. In fact, anytime the blood of animals becomes supersaturated with gas, bubbles can form.

This condition is known as gas bubble trauma, and it can lead to stress or mortality. Eggs may float to the surface, and larvae and fry may exhibit hyperinflation of the swim bladder, cranial swelling, swollen gill lamellae and other abnormalities.

A common symptom of acute gas bubble trauma in juvenile and adult fish is gas bubbles in the blood that can be seen in the surface tissues on the head, in the mouth and in fin rays. The eyes of affected fish also tend to protrude.

Saturation assessment

A variable known as ΔP is used to assess gas supersaturation in water relative to gas bubble trauma. The ΔP is defined as the difference between the total gas pressure in water and the barometric pressure at a given location.

The ΔP can be calculated by measuring the partial pressure of each gas in the water [(percentage saturation/100 x partial pressure in the atmosphere), summing the partial pressures and subtracting from the sum the barometric pressure. Fortunately, a relatively inexpensive instrument called a satumeter can be used to measure ΔP directly.

Aquatic animals exposed to ΔP values of 25 to 75 mm Hg on a continuous basis may exhibit some symptoms of gas bubble trauma, and low-level mortality may occur over an extended period of time. Acute gas bubble trauma occurs at greater levels of ΔP . Symptoms will be more pronounced, and mortalities typically are 50 to 100 percent.

Supersaturation of pond waters with dissolved oxygen during afternoons is a common occurrence. This condition usually does not harm culture animals, for supersaturation does not persist for long and is often limited to surface waters. Animals can move to greater depths, where the combination of lower dissolved-oxygen concentration and greater hydrostatic pressure result in a lower ΔP .

Nevertheless, carp reportedly had a greater frequency of disease when percentage saturation with dissolved oxygen exceeded 150 percent (ΔP above 225 mm Hg). Mortality of fish and shrimp has been reported in culture systems where dissolved-oxygen supersaturation exceeded 300 percent (ΔP above 450 mm Hg).

Management

Supersaturating gases can be removed from water in degassing towers in which water is passed through screens or other media to increase exposure to the atmosphere. In ponds, aerators that splash water into the air can lessen afternoon supersaturation with dissolved oxygen caused by a high rate of photosynthesis. Of course, managers always should guard against gas supersaturation caused by heating water, pumps, submerged aerators and air entrainment.

(Editor's Note: This article was originally published in the July/August 2012 print edition of the Global Aquaculture Advocate.)

Author



CLAUDE E. BOYD, PH.D.

Department of Fisheries
and Allied Aquacultures
Auburn University
Auburn, Alabama 36849 USA

boydce1@auburn.edu (<mailto:boydce1@auburn.edu>).

Copyright © 2023 Global Seafood Alliance

All rights reserved.