



ALLIANCE™

(<https://gsa.rakadev.com>).



**Responsible
Seafood**
ADVOCATE




Health &
Welfare

Effects of suspended solids in shrimp biofloc systems

1 November 2013

By Carlos Gaona , Plínio Furtado , Fabiane Serra , Luis Poersch and Wilson Wasielesky, Ph.D.

Considerations for water quality management



The use of lined tanks increases the confinement of biofloc in the culture environment and restricts the organic particles in the water column from being recycled completely.

Marine shrimp aquaculture systems that incorporate biofloc technology often experience a high density of suspended solids within the systems. The injection of air at the bottoms of culture tanks promotes both the diffusion of oxygen in the water column and mixing of suspended material. The use of lined tanks increases the confinement of biofloc in the culture environment and restricts the organic particles in the water column from being recycled completely.

Suspended solids

Suspended solids consist mainly of organic matter, which is comprised of microbial forms that when decomposing exert a high demand for oxygen. This demand can decrease the dissolved-oxygen concentrations in the culture system to reach levels below the recommended concentration for the cultivated species. An increase in suspended solids can also reduce the water quality within the system (Table 1). Overall, these less-than-optimal conditions reduce system performance.

Gaona, Changes in water quality parameters, Table 1

Parameter	Effect
Ammonia	Increase
Nitrite	Increase
Nitrate	Increase
Phosphorus	Increase
pH	Increase
Alkalinity	Decrease

Dissolved oxygen	Decrease
Carbon dioxide	Increase

Table 1. Changes in water quality parameters with excessive suspended solids.

These conditions can occur throughout the different stages of the marine shrimp production cycle, depending on whether the system starts a cycle with mature biofloc inoculum. However, the use of biofloc inoculum relates to the suspended solids concentration in the water column.

At the beginning of a cycle, interactions among the nitrogenous compounds, ammonia and nitrite are most notable when biofloc is forming in the culture. The reduction of ammonia occurs with the establishment of ammonium-oxidizing bacteria, which does not require the use of organic carbon, and with the absorption of heterotrophic bacteria. Therefore, the nitrite accumulation occurs due to the slow growth of nitrite-oxidizing bacteria.

Total suspended solids levels constantly increase, and consequently nitrite concentrations increase. In contrast, when a cycle begins with a well-stocked biofloc inoculum, the concentrations of ammonia and nitrite are relatively low. However, the concentrations of suspended solids are higher in this phase. Both situations require management of suspended particulate matter within the culture system.

Experimental work

In experiments conducted with *Litopenaeus vannamei* shrimp in biofloc systems at the Marine Aquaculture Station of the Federal University of Rio Grande in southern Brazil, the best growth performance occurred in a study where suspended solids were removed for the maintenance and control of the total suspended solids concentrations in culture inoculated with biofloc.

In another experiment, concentrations of nitrogen compounds differed when compared to different total suspended solids (TSS) concentrations during the formation of biofloc. Higher TSS concentrations resulted in higher concentrations of nitrite.



Greenhouse shrimp production with linear tanks.

In another experiment that used the biofloc inoculum, it was observed that when the concentration of dissolved oxygen was maintained above 5 mg/L, the suspended solids excess was not a problem for the respiration of the reared shrimp. Considering the interactions of water quality parameters listed above, maintenance of the suspended solids levels is required for better water quality. Values for common water quality parameters measured under two average TSS concentrations are shown in Table 2.

Gaona, Water quality parameter values, Table 2

Parameter	200 mg/L TSS	800 mg/L TSS
-----------	--------------	--------------

Dissolved oxygen (mg/L)	5.90	5.10
pH	7.95	7.40
Alkalinity (mg calcium carbonate/L)	160	100
Ammonia (mg/L)	3.50	2.50
Nitrite (mg/L)	1.00	15.50
Nitrate (mg/L)	5.00	80.00
Phosphate (mg/L)	0.50	6.00

Table 2. Water quality parameter values with average total suspended solids concentrations.

Monitoring, intervention

Measurements of suspended solids were made using the gravimetric method, which measures total suspended solids and settleable solids within an Imhoff cone. Various techniques can be applied to reduce and maintain suspended solids concentrations, such as the use of a settling chamber or clarifier to remove solids. Set up in straightforward settling chambers, clarifiers rely on gravity to move particles to the bottom (Fig. 1).




Fig. 1: Settling chambers use gravity to settle particles. The arrows indicate water flow. Water from the culture tank enters via a central pipe, which reduces velocity and turbulence, and returns after settling.

The radial water flow in the settling chamber can be adjusted based on prior analysis of biofloc sedimentation in an Imhoff cone, increasing the efficiency of the method. This method allows the control of TSS and keeps the concentrations near the recommended values (Fig. 2). Another advantage of this method is maintaining a constant flow during the application of settling, so a small amount of water is sufficient for removing suspended solids.

Fig. 2: Total suspended solids concentrations in study with and without clarification. The arrow indicates the seventh week since the start of the solids removal process.

(Editor's Note: This article was originally published in the November/December 2013 print edition of the Global Aquaculture Advocate.)

Authors



CARLOS GAONA

Laboratório de Carcinocultura
Universidade Federal do Rio Grande
Rio Grande (R.S.) 96201-900 Brazil



PLÍNIO FURTADO

Laboratório de Carcinocultura
Universidade Federal do Rio Grande
Rio Grande (R.S.) 96201-900 Brazil



FABIANE SERRA

Laboratório de Carcinocultura
Universidade Federal do Rio Grande
Rio Grande (R.S.) 96201-900 Brazil



LUIS POERSCH

Laboratório de Carcinocultura
Universidade Federal do Rio Grande
Rio Grande (R.S.) 96201-900 Brazil



WILSON WASIELESKY, PH.D.

Laboratório de Carcinocultura
Universidade Federal do Rio Grande
Rio Grande (R.S.) 96201-900 Brazil

manow@mikrus.com.br (<mailto:manow@mikrus.com.br>).

Copyright © 2022 Global Seafood Alliance

All rights reserved.