Global Seafood Alliance Logo

- GOAL Events
- Advocate Magazine
- Aquademia Podcast
- <u>Blog</u>
- Contact
- 0
- X
- in
- Log In

- About
 - Who We Are
 - Our History
 - o Our Team
 - Sustainable Development Goals
 - Careers
- <u>Membership</u>
 - o <u>Overview</u>
 - Our Members
 - Corporate Membership
- Resources
- Certification
 - Best Aquaculture Practices
 - Best Seafood Practices

Search...

Q

Log In

- About
 - Who We Are
 - o Our History
 - o Our Team
 - Sustainable Development Goals
 - o <u>Careers</u>
- Membership
 - o <u>Overview</u>
 - o Our Members
 - Corporate Membership
- Resources
- Certification
 - Best Aquaculture Practices
 - Best Seafood Practices
- GOAL Events
- Advocate Magazine
- Aquademia Podcast
- **Blog**
- Contact



Essential oils increase weight gain in channel catfish

Responsible Seafood Advocate logo

1 July 2011 Brian C. Peterson, Ph.D. Brian G. Bosworth, Ph.D. Monica L. Wood Menghe H. Li, Ph.D. Ruben Beltran, M.S.



Fillets tended to have more protein, less fat



Research with domestic livestock that suggested essential oils can improve growth, feed efficiency and the ability to ward off diseases is carrying over to aquaculture species.

Essential oils, also known as volatile oils, are highly scented compounds of fragrant grasses, trees and plants. The word "essential" suggests that essential oils are vital to the life of plants, but this is not the case. The word "essential" is derived from the word "quintessence," which can be defined as the "pure and concentrated essence of a substance." Essential oils contain most of the plants' active substances.

Many essential oil mixtures have been used for centuries in traditional medicine in the treatment of bacterial and viral diseases, inflammation, pain and even some forms of cancer. Today, they are commonly found as ingredients in cosmetics, perfumes, cleaning products and flavorings for food and drinks.

As the popularity of essential oils increases, products specific to the domestic livestock and aquaculture markets have emerged. These products – often sold as feed additives – appear to be associated with realized performance gains. Several studies have focused on unraveling the mode of action of the oils. It seems the modulation of gut microflora plays an important role, but many questions remain on how the additives provide their apparent medicinal

At the animal level, research with domestic livestock suggests that essential oils may improve growth and feed efficiency, and increase the ability to ward off diseases. Although fewer studies using the oils in fish have been conducted, the data also suggest that essential oils have benefits in improving growth and controlling diseases.

Current research

At the Thad Cochran National Warmwater Aquaculture Center in Stoneville, Mississippi, USA, the authors conducted a trial to examine the effects of a commercial matrix-encapsulated essential oil (Biomin P.E.P. MGE) on weight gain, specific growth rate, feed-conversion ratio (FCR) and survival of

channel catfish.

Groups of 50, 32.4 ± 1.7-gram fish per tank were randomly assigned to two treatments with five replicate tanks per treatment. Treatment 1 was a control with a 32 percent-crude protein floating commercial diet. Treatment 2 used the same diet with the addition of the essential oil at 200 g/mt.

The 1.15-m³ tanks were supplied with recirculated pond water and aeration. The fish were fed once a day to apparent satiation for 12 weeks and weighed every four weeks. Water quality was monitored throughout the study. At the end of the study, fillets from two fish per tank were analyzed for protein and

Results

Water temperature and dissolved-oxygen concentrations averaged 32.9 degrees-C and 9.3 mg/L, respectively. Total ammonia nitrogen and nitrite averaged 0.34 and 0.01 mg/L, respectively. By week 8, fish in treatment 2 gained more weight (51.4 ± 1.9 versus 37.3 ± 5.1 grams/fish) and had a higher specific growth rate (1.8 \pm 0.1 versus 1.5 \pm 0.1) than the controls (P < 0.03).

At the end of the study, fish that received the essential oil product gained more weight (76.9 ± 2.0 versus 53.4 ± 3.2 g/fish) and had a higher specific growth rate $(1.5 \pm 0.1 \text{ versus } 1.3 \pm 0.1)$ than the controls (P < 0.001) (Table 1). In addition, fish fed the essential oil consumed more feed (104.3 ± 3.6) versus 79.6 ± 3.0 grams/fish), suggesting an increase in feed palatability.

Peterson, Growth performance of channel catfish fed diets, Table 1

Treatment	Initial Weight (g)	Final Weight (g)	Weight Gain (g)	Specific Growth Weight	Feed Conversion Ratio	Survival (%)
Control diet	33.40	88.40 ^a	53.4 ^a	1.30 ^a	1.51	89.5
Diet with essential oil	31.40	112.80 ^b	76.90 ^b	1.50 ^b	1.36	90.0
Standard error	1.94	2.48	2.17	0.03	0.06	4.8

Table 1. Growth performance of channel catfish fed diets with and without an essential oil additive for 12 weeks. Values with different letters within columns are significantly different (P < 0.001).

There was an improvement in FCR (1.36 versus 1.51) in fish that received the essential oil, although it was not statistically different (P > 0.05). The survival rates of about 90 percent were similar in both treatments, as no natural outbreaks of disease were recorded. Fillet composition analysis showed that the amount of fat in the fillets of fish fed the essential oil was lower (16.1 versus 18.7 percent), and the amount of protein was higher (79.6 versus 76.5 percent) compared to controls (P < 0.09).

Perspectives

The results showed that fish fed an essential oil consumed more feed and gained more weight. In addition, fillets from fish fed the oil tended to have higher amounts of protein and lower amounts of fat.

The mechanisms through which the essential oil increased weight gain were not determined in this study, but may be related to an increase in appetite. The addition of essential oils to catfish diets may prove beneficial in improving the palatability of feed as well as the growth efficiency of channel catfish.

(Editor's Note: This article was originally published in the July/August 2011 print edition of the Global Aquaculture Advocate.)

Authors

• Brian C. Peterson, Ph.D.

Brian C. Peterson, Ph.D.

Thad Cochran National Warmwater Aquaculture Center U.S. Department of Agriculture Agricultural Research Service 127 Experiment Station Road Stoneville, Mississippi 38776 USA

[118,111,103,46,97,100,115,117,46,115,114,97,64,110,111,115,114,101,116,101,112,46,110,97,105,114,98]

Brian G. Bosworth, Ph.D.

Brian G. Bosworth, Ph.D.

Thad Cochran National Warmwater Aquaculture Center U.S. Department of Agriculture Agricultural Research Service 127 Experiment Station Road Stoneville, Mississippi 38776 USA

Monica L. Wood

Monica L. Wood

Thad Cochran National Warmwater Aquaculture Center U.S. Department of Agriculture Agricultural Research Service 127 Experiment Station Road Stoneville, Mississippi 38776 USA

• Menghe H. Li, Ph.D.

Menghe H. Li, Ph.D.

Thad Cochran National Warmwater Aquaculture Center U.S. Department of Agriculture Agricultural Research Service 127 Experiment Station Road Stoneville, Mississippi 38776 USA

• Ruben Beltran, M.S.

Ruben Beltran, M.S.

Biomin America, Inc. San Antonio, Texas, USA

Share

- Share via Email
- **Share on Twitter**
- Share on Facebook
- in Share on LinkedIn

Tagged With

feed efficiency channel catfish Essential oils Brian C. Peterson Brian G. Bosworth Menghe H. Li Ruben Beltran Monica L. Wood weight gain

Related Posts

Aquafeeds

Alternative feed ingredients support continued aquaculture expansion

Identifying sources for essential macro- and micronutrients is important, as well as understanding how best to manufacture feed to required physical specifications when using these new raw materials.

Innovation & Investment

Assessing coloration in channel catfish fillets

Because consumers look at color to gauge quality of catfish fillets, the authors developed a digital photography measurement method to assess yellowness.

Health & Welfare

Biofloc technology reduces common off-flavors in channel catfish

In studies that used biofloc systems to culture channel catfish, culture tanks were susceptible to episodes of geosmin and 2-methylisoborneol and subsequent bioaccumulation of off-flavors in catfish flesh.

Health & Welfare

Blue catfish outproduce channel catfish under low-D.O. conditions

Although there is increasing interest in blue catfish, a potential disadvantage of the fish when compared to channel catfish is their reported poorer tolerance of low dissolved-oxygen concentrations.

About The Advocate

The Responsible Seafood Advocate supports the Global Seafood Alliance's (GSA) mission to advance responsible seafood practices through education, advocacy and third-party assurances.

Search **Q**

Learn More

Search Responsible Seafood Advocate Search Aquademia Listen to the seafood industry's top podcast

Advertising Opportunities

2022 Media & Events Kit

Categories

Aquafeeds > Health & Welfare Health & Welfare > From Our Sponsors > Innovation & Investment > Intelligence > Responsibility > Fisheries > Artículos en Español >

Don't Miss an Article

Featured

- Health & Welfare An update on vibriosis, the major bacterial disease shrimp farmers face
- Intelligence A seat at the table: Fed By Blue team says aquaculture needs a stronger voice
- Responsibility Quantifying habitat provisioning at macroalgae cultivation locations

Popular Tags



Recent

- Fisheries Second Test: Another filler for the fisheries category
- <u>Fisheries Test: This is filler for the fisheries Category</u>
- **Aquafeeds Test Article**
- Responsibility Study: Climate change will shuffle marine ecosystems in unexpected ways as ocean temperature warms
- Health & Welfare Indian shrimp researchers earn a patent for WSSV diagnostic tool



- **About**
- <u>Membership</u>
- Resources
- Best Aquaculture Practices (BAP)
- Best Seafood Practices (BSP)
- GOAL Events
- Advocate Magazine
- Aquademia Podcast
- Blog
- Contact

Stay up to date with GSA

- 0

Copyright © 2024 Global Seafood Alliance All rights reserved.

Privacy Terms of Use <u>Glossary</u>