



- [GOAL Events](#)
- [Advocate Magazine](#)
- [Aquademia Podcast](#)
- [Blog](#)
- [Contact](#)
-
-
-
-
-
- [Log In](#)



- [About](#)
 - [Who We Are](#)
 - [Our History](#)
 - [Our Team](#)
 - [Sustainable Development Goals](#)
 - [Careers](#)
- [Membership](#)
 - [Overview](#)
 - [Our Members](#)
 - [Corporate Membership](#)
- [Resources](#)
- [Certification](#)
 - [Best Aquaculture Practices](#)
 - [Best Seafood Practices](#)

Search...



[Log In](#)

- [About](#)
 - [Who We Are](#)
 - [Our History](#)
 - [Our Team](#)
 - [Sustainable Development Goals](#)
 - [Careers](#)
- [Membership](#)
 - [Overview](#)
 - [Our Members](#)
 - [Corporate Membership](#)
- [Resources](#)
- [Certification](#)
 - [Best Aquaculture Practices](#)
 - [Best Seafood Practices](#)
- [GOAL Events](#)
- [Advocate Magazine](#)
- [Aquademia Podcast](#)
- [Blog](#)
- [Contact](#)



Evaluating the aquaculture potential of the neotropical fish trahira



24 February 2020 Marcos Daniel Renó Faria, M.S. Dr. Aderbal Cavalcante-Neto Dr. Ivan Bezerra Allaman Dr. Aline Dal'Olio Gomes Dr. Renata Guimarães Moreira Eric M. Hallerman, Ph.D. Dr. Alexandre Wagner Silva Hilsdorf



Results of growth rates, fatty acid content, hardness and market support its production



Study results showed that trahira could be a viable aquaculture species in Neotropical regions. Photo by Vassil, Wikimedia Creative Commons.

Aquaculture development has driven the introduction of many alien species worldwide. The environmental risks posed by such introductions may threaten native fish diversity, so the proper evaluation of native species as viable aquaculture candidates is important for further development of sustainable fish farming.

The neotropical region [the tropical terrestrial eco-regions of the Americas and the entire South American temperate zone] is rich in native fish genetic resources potentially useful for aquaculture. Among these, there are eight species of the genus *Hoplias* or trahiras that occur from Costa Rica to Argentina. The different trahiras are economically important for local artisanal fisheries, but have not been farmed yet at any significant scale; for example, the aquaculture production of all trahiras in Brazil in 2016 was just over 806 metric tons (MT).

The wolf fish, tiger fish or trahira (*Hoplias malabaricus*) is a candidate species for aquaculture production. It is a carnivorous fish that preferentially inhabits lotic [moving] water environments but also occurs in lentic [still] waters such as lakes and reservoirs. It is a sturdy species well adapted to, and can reproduce in, hypoxic [low oxygen] waters. The nutritional quality of its meat and new processing techniques producing boneless, quality fillets and high fillet yield (~44 percent) have attracted many consumers to trahira as a delicacy fish that fetches high prices in markets.

There are some constraints to the development of its aquaculture production, including commercial availability of feed-trained fingerlings; cannibalism in early life stages; limited information on nutritional requirements and growth production data; undesirable intramuscular bones; and the continuing availability of market-sized fish from capture fisheries.

This article – adapted and summarized from the [original](#) (Faria, M.D.R. et al. 2019. The potential of *Hoplias malabaricus*, a Neotropical carnivore, for aquaculture) evaluated its aquaculture potential by determining the best growth model through one year of semi-intensive production; quantified its growth rate from fry through grow-out under two production densities; and assessed fillet quality in terms of total protein, lipid levels and fatty acid profile.

Study setup

The growth trial was carried out at the Ponte Nova Fish Farm located in Salesópolis County, State of São Paulo, Brazil, a region with subtropical climate with mean annual temperature of 19 degrees-C. A total of 270 *H. malabaricus* juveniles were obtained from the Brumado Fish Hatchery, Ltd. in the State of São Paulo, which had produced them from wild broodstock.

The juvenile fish were stocked in four 12.6 cubic meters concrete ponds using a completely randomized design and at two stocking densities, 5 fish per square meter (90 juveniles) and 10 fish per square meter (180 juveniles). Fish were fed exclusively using an extruded commercial diet (NutriPeixe TC 40, Purina, Brazil), and the feeding regime was based on that commercially proposed for carnivorous fishes. The growth trial lasted 11 months from September to August. Every 40 days through the experiment, all individuals were anesthetized, measured and weighed.

For information on the experimental setup; fish and rearing conditions; grow-out performance and growth model data analyses; and lipids and protein content profiles, refer to original publication.

Results and discussion

The mean weight after 11 months of grow-out did not differ significantly between the two densities, 263.0 grams and 246.7 grams for 5 and 10 fish per square meter, respectively. It should be noted that the juveniles we used originated from wild broodstock which had not undergone genetic improvement. The high coefficient of variation at the last weight measurement may reflect the full range of phenotypic variation for the trait.

Our results showed that stocking densities had little influence on growth rate, protein and lipid content, or fatty acid profile of fillets of *H. malabaricus*. The survival rates were 69 percent at a density of 5 fish per square meter and 76 percent at 10 fish per square meter.

After the eleventh month of rearing, including the winter months, fish weighing more than 450 grams were found among the trial fish. The market demand for an individual portion (plate size) and the growth rate achieved in this study indicate the commercially interesting weight for trahira is between 500 and 600 grams. Observation of fish with higher growth rates under semi-intensive culture management suggests that *H. malabaricus* may have potential for improved growth by selective breeding. Domestication and selective breeding to improve feed conversion rate would address a factor critical for adoption of *H. malabaricus* for commercial aquaculture.

Regarding grow-out performance, initial total length, final total length and body weight, as well as the majority of the growth parameters, were statistically equal at both stocking densities. Conversely, feed conversion rate and survival rate showed statistically significant differences between stocking densities by the end of the experiment. The coefficients of variation for growth traits were high and within the range found in fish and shellfish (CV = 17 to 29 percent).

Successful cultivation of an aquaculture candidate species depends on genetic make-up and physiological plasticity so that the species can adjust to culture conditions and show a viable growth rate for commercialization. Therefore, modeling fish growth is a first step for assessing an aquaculture candidate species because it supports predictions of how well the fish will grow under various production systems. Our results showed that *H. malabaricus* under semi-intensive farming conditions presents positive allometric growth, i.e., individuals increase in relative body thickness, becoming heavier at a given length as the fish grows (Fig. 1).




Fig. 1: Growth of weight based on the values observed and predicted by the logistic model, with a 95 percent confidence interval.

Our assessment of growth models for cultured *H. malabaricus* showed that the mixed-model approach provided a better fit to the data than traditional modeling with only fixed effects.

The typical lifetime growth curve of fish has a sigmoid shape, i.e., growth during early life is slow and is followed by a period of acceleration, and then growth reaches its maximum around puberty, when deceleration occurs. Our results for trahira weight growth show that at the lower stocking density (5 fish per square meter), the convex phase of the growth curve started earlier than at the higher stocking density (10 fish per square meter). However, the growth rate in the fish at a higher stocking density increased until the sixth month of grow-out.

Growth models for farmed fishes may be affected by environmental factors, particularly oxygen availability and temperature, as well as culture-dependent factors such as stocking density and feeding. To assess the growth curve for a candidate fish for aquaculture, such as *H. malabaricus*, it is important to understand and manage culture variables in order to improve growth rate, and to determine the optimal cost-benefit weight for harvest size. The grow-out performance of *H. malabaricus* during our one year-production cycle had similar outcomes for the two stocking densities, as can be seen in Table 1.

Faria, trahira, Table 1

Growth parameters	Stocking density 5 fish per square meter	Stocking density 10 fish per square meter
Initial total length (cm)	12.0±0.06a	12.2±0.04a
Final total length (cm)	27.5±0.38a	27.0±0.24a
Initial body weight (g)	18.6±0.26a	19.4±0.18a

Growth parameters	Stocking density 5 fish per square meter	Stocking density 10 fish per square meter
Final body weight (g)	263.0±10.59a	246.7±6.89a
Relative growth (%)	43.03±12.51a	39.58±9.85a
Specific growth rate (%/day)	0.36±0.08a	0.35±0.07a
FCR	3.90:1a	3.34:1b
Condition factor (K)	1.23±0.03a	1.23±0.30a
Feed efficiency (%)	34.24±8.22a	37.05±6.53a
Coefficient of variation (%)	24.8±1.29a	23.67±0.82a
Survival rate (%)	69a	76b

Table 1. Growth rate parameters from *H. malabaricus* juveniles reared at two stocking densities (5 fish per square meter and 10 fish per square meter) under on-farm grow-out conditions. Data are shown as mean ± standard error. Means in a row followed by the same superscript are not significantly different ($P \leq 0.05$).

Regarding the body composition, total protein concentrations in fillets decreased through the experiment at both stocking densities; however, the values still remained higher than those for other species analyzed, such as Murray cod, gilthead sea bream, Tiete tetra, goldfish and Nile tilapia. The decrease in fillet protein may be due to utilization of protein as an energy source for growth, suggesting that use of a higher protein diet (> 40 percent) may be appropriate for grow-out of *H. malabaricus*.

Lipid content is also an important aspect of fillet quality. Fillets can be classified as “lean” when the lipid content is below 2.5 percent and as “semi-skimmed” between 2.5 and 10 percent. In our study, high concentrations of lipids in the muscle of the *H. malabaricus* juveniles were such that the fillet would be classified as high-quality in the semi-skimmed range. The fillet fatty acid (FA) profile showed few changes over time, but we noted an interesting content of saturated fatty acids (SFAs) and polyunsaturated fatty acids (PUFAs). The major FAs in *H. malabaricus* juvenile fillets were linoleic acid (LA), arachidonic acid (ARA), and docosahexaenoic acid (DHA).

H. malabaricus juveniles in our study had a high percentage of linoleic acid (almost 90 percent of total PUFAs), suggesting that they tend to accumulate desirable omega-3 PUFAs, mainly DHA, in their fillets, even at high stocking densities and with a relatively low percentage of this FA in the diet.

Perspectives

Based on our evaluation of the aquaculture potential of trahira under semi-intensive culture conditions – including growth rates and fillet quality – and considering the species hardiness, tolerance for low oxygen conditions, acceptance and consumption of manufactured aquafeeds, considerable growth potential, and high nutritional quality and superior meat flavor, *H. malabaricus* appears to be a viable aquaculture species for Central and South American countries, where the development of native species can contribute to sustainable aquaculture production.

References available from the original publication.

Now that you've finished reading the article ...

... we hope you'll consider supporting our mission to document the evolution of the global aquaculture industry and share our vast network of contributors' expansive knowledge every week.

By becoming a Global Seafood Alliance member, you're ensuring that all of the pre-competitive work we do through member benefits, resources and events can continue. Individual membership costs just \$50 a year.

Not a GSA member? Join us.

[Support GSA and Become a Member](#)

Authors

-  Marcos Daniel Renó Faria, M.S.

Marcos Daniel Renó Faria, M.S.

Unit of Biotechnology
University of Mogi das Cruzes
Mogi das Cruzes, SP, 08780-970, Brazil

-  Dr. Aderbal Cavalcante-Neto

Dr. Aderbal Cavalcante-Neto

Unit of Biotechnology
University of Mogi das Cruzes
Mogi das Cruzes, SP, 08780-970, Brazil

-  Dr. Ivan Bezerra Allaman

Dr. Ivan Bezerra Allaman

Department of Exact Sciences and Technology
State University of Santa Cruz
Ilhéus, BA, 45662-900, Brazil

-  Dr. Aline Dal'Olio Gomes

Dr. Aline Dal'Olio Gomes

Department of Physiology, Bioscience Institute
University of São Paulo
São Paulo, SP, 05508-020, Brazil

-  Dr. Renata Guimarães Moreira

Dr. Renata Guimarães Moreira

Department of Physiology, Bioscience Institute
University of São Paulo
São Paulo, SP, 05508-020, Brazil

-  Eric M. Hallerman, Ph.D.

Eric M. Hallerman, Ph.D.

Department of Fish and Wildlife Conservation
Virginia Polytechnic Institute and State University
Blacksburg, VA 23061-0321 USA





-  Dr. Alexandre Wagner Silva Hilsdorf

Dr. Alexandre Wagner Silva Hilsdorf

Corresponding author
Unit of Biotechnology
University of Mogi das Cruzes
Mogi das Cruzes, SP, 08780-970, Brazil

[114,98,46,99,109,117,64,114,101,110,103,97,119]

Share

-  [Share via Email](#)
-  [Share on Twitter](#)
-  [Share on Facebook](#)
-  [Share on LinkedIn](#)

Tagged With

[Hoplias malabaricus](#) [Brazil](#) [Eric M. Hallerman](#) [Marcos Daniel Renó Faria](#) [Aderbal Cavalcante-Neto](#) [Ivan Bezerra Allaman](#) [Aline Dal'Olio Gomes](#) [Renata Guimarães Moreira](#) [Alexandre Wagner Silva Hilsdorf](#) [Trahira](#)

Related Posts

[Responsibility](#)

[Brazil's inland aquaculture](#)

Freshwater fish culture is practiced in every state in Brazil, primarily at small-scale earthen pond facilities, but tilapia culture in reservoir cages is increasing.

[Innovation & Investment](#)

[Pirarucu culture in the Brazilian Amazon](#)

With their fast growth and ample light-flavored flesh, pirarucu present great potential. Industry growth is hampered by limited technology and management of domestication practices. Also, current commercial diets are typically not fully suited to the nutritional needs of pirarucu.

[Health & Welfare](#)

[Arapaima a candidate for intensive freshwater culture](#)

Arapaima, also known as pirarucu or paiche, offers appealing flesh color and texture, low fat content and high levels of protein and fatty acids.

[Aquafeeds](#)

[Soy-based feeds evaluated for production of Amazonian paiche](#)

A cooperative feeding demonstration conducted by USSEC and Amazon Fish Products S.A. evaluated the use of soy to partially replace fishmeal in diets for paiche, or pirarucu, the largest freshwater fish species.

About The Advocate

The Responsible Seafood Advocate supports the Global Seafood Alliance's (GSA) mission to advance responsible seafood practices through education, advocacy and third-party assurances.

[Learn More](#)

Search Responsible Seafood Advocate



Advertising Opportunities

[2022 Media & Events Kit](#)

Categories

[Aquafeeds](#) > [Health & Welfare](#) > [From Our Sponsors](#) > [Innovation & Investment](#) > [Intelligence](#) > [Responsibility](#) > [Fisheries](#) > [Artículos en Español](#) >

Don't Miss an Article

Featured

- [Health & Welfare](#) [An update on vibriosis, the major bacterial disease shrimp farmers face](#)
- [Intelligence](#) [A seat at the table: Fed By Blue team says aquaculture needs a stronger voice](#)
- [Responsibility](#) [Quantifying habitat provisioning at macroalgae cultivation locations](#)

Popular Tags

All Tags ▼

Recent

- Fisheries [Second Test: Another filler for the fisheries category.](#)
- Fisheries [Test: This is filler for the fisheries Category.](#)
- [Aquafeeds Test Article](#)
- [Responsibility Study: Climate change will shuffle marine ecosystems in unexpected ways as ocean temperature warms](#)
- Health & Welfare [Indian shrimp researchers earn a patent for WSSV diagnostic tool](#)



- [About](#)
- [Membership](#)
- [Resources](#)
- [Best Aquaculture Practices \(BAP\)](#)
- [Best Seafood Practices \(BSP\)](#)
- [GOAL Events](#)
- [Advocate Magazine](#)
- [Aquademia Podcast](#)
- [Blog](#)
- [Contact](#)

Stay up to date with GSA

- 
- 
- 
- 
- 

Copyright © 2024 Global Seafood Alliance
All rights reserved.

[Privacy](#)
[Terms of Use](#)
[Glossary](#)