



ALLIANCE™

<https://gsa.rakadev.com>**Responsible
Seafood**
ADVOCATE

Aquafeeds

Fatty acid nutrition of juvenile *Litopenaeus vannamei*

1 October 2002


By Mayra Lizett González-Félix, Ph.D. , Martin Perez-Velazquez, Ph.D. , Addison L. Lawrence, Ph.D. and Delbert M. Gatlin III, Ph.D.

Recirculating-system research conducted at Texas A&M

Among lipids, some omega-3 and omega-6 polyunsaturated fatty acids (PUFA) such as linolenic (18:3 omega-3) and linoleic (18:2 omega-6) acid, as well as highly unsaturated fatty acids (HUFA) such as arachidonic (20:4 omega-6), eicosapentaenoic (20:5 omega-3), and docosahexaenoic (22:6 omega-3) acid are considered essential nutrients for penaeid shrimp. In spite of this, knowledge of the dietary requirements for these nutrients in Pacific white shrimp (*Litopenaeus vannamei*), an economically important penaeid for shrimp farming in the Americas, is limited.

The authors conducted a six-week experiment at the Shrimp Mariculture Project of Texas A&M University in Port Aransas, Texas, USA to test the dietary inclusion of 18:3 omega-3, 18:2 omega-6, 20:4 omega-6, 20:5 omega-3, and 22:6 omega-3 fatty acids at 0.5 percent of diet. An additional diet contained 0.5 percent of an omega-3 HUFA mix, which included 416 milligrams per gram of 20:522:6 omega-3-3 and 237 milligrams per gram of 22:6 omega-3. A basal diet contained only palmitic (16:0) and stearic (18:0) acid, also used as fillers for a 5 percent total dietary lipid.

L. vannamei juveniles with a mean initial weight of 0.4 grams were stocked at 44.4 animals per square meter into 10, 32-liter replicate tanks per dietary treatment. The indoor aerated recirculating system was kept at a temperature of 30 degrees-C and a salinity of 25 ppt.



Nutrition lab at Texas A&M Shrimp Mariculture Project.

Shrimp growth and survival

Differences in weight gain of shrimp were observed at the end of the feeding trial (Fig. 1). HUFA like 20:4 omega-6, 20:5 omega-3, and 22:6 omega-3, as well as the omega-3 HUFA mix, showed higher nutritional value than PUFA like 18:3 omega-3 and 18:2 omega-6 for juvenile *L. vannamei*.

Neither 18:3 omega-3 or 18:2 omega-6 improved shrimp growth significantly when compared to animals fed the basal diet with 16:0 and 18:0 only. Survival of shrimp averaged almost 94 percent and was not significantly affected by the dietary treatments.




Fig. 1: Initial weight and weight gain of juvenile *Litopenaeus vannamei* fed different PUFA and HUFA (mean values). Bars with the same superscript are not significantly different.

Conclusion

As demonstrated for juvenile *L. vannamei* in the study, HUFA generally show higher nutritive value than PUFA. However, no preferential activity as essential fatty acids was observed for the omega-3 over the n-6 family of fatty acids under our experimental conditions, as different reports have indicated.

Results suggested that for *L. vannamei*, the essential fatty acid value may be determined by chain length and degree of unsaturation, with long-chain unsaturated fatty acids having greater nutritional value than shorter-chain fatty acids, regardless of the family to which they belong.

(Editor's Note: This article was originally published in the October 2002 print edition of the Global Aquaculture Advocate.)

Authors

**MAYRA LIZETT GONZÁLEZ-FÉLIX, PH.D.**

Departamento de Investigaciones Científicas y Tecnológicas
Universidad de Sonora
Rosales y Niños Héroes
A.P. 1819 C.P. 83000
Hermosillo, Sonora, México

mgonzale@correom.uson.mx (<mailto:mgonzale@correom.uson.mx>)

**MARTIN PEREZ-VELAZQUEZ, PH.D.**

Departamento de Investigaciones Científicas y Tecnológicas
Universidad de Sonora
Rosales y Niños Héroes
A.P. 1819 C.P. 83000
Hermosillo, Sonora, México

**ADDISON L. LAWRENCE, PH.D.**

Texas A&M University
Shrimp Mariculture Project
Port Aransas, Texas, USA

**DELBERT M. GATLIN III, PH.D.**

Department of Wildlife and Fisheries Sciences
Texas A&M University
College Station, Texas, USA

Copyright © 2023 Global Seafood Alliance

All rights reserved.