

 [Global Seafood Alliance Logo](#)

- [GOAL Events](#)
- [Advocate Magazine](#)
- [Aquademia Podcast](#)
- [Blog](#)
- [Contact](#)

- 
- 
- 
- 
- 

- [Log In](#)

—

- [About](#)
 - [Who We Are](#)
 - [Our History](#)
 - [Our Team](#)
 - [Sustainable Development Goals](#)
 - [Careers](#)
- [Membership](#)
 - [Overview](#)
 - [Our Members](#)
 - [Corporate Membership](#)
- [Resources](#)
- [Certification](#)
 - [Best Aquaculture Practices](#)
 - [Best Seafood Practices](#)

Search...



[Log In](#)

- [About](#)
 - [Who We Are](#)
 - [Our History](#)
 - [Our Team](#)
 - [Sustainable Development Goals](#)
 - [Careers](#)
- [Membership](#)
 - [Overview](#)
 - [Our Members](#)
 - [Corporate Membership](#)
- [Resources](#)
- [Certification](#)
 - [Best Aquaculture Practices](#)
 - [Best Seafood Practices](#)
- [GOAL Events](#)
- [Advocate Magazine](#)
- [Aquademia Podcast](#)
- [Blog](#)
- [Contact](#)



 Aquafeeds
Aquafeeds

Feed management for RAS, part 2

 [Responsible Seafood Advocate logo](#)

1 April 2001 Phillip G. Lee, Ph.D. Addison L. Lawrence, Ph.D.



Improving feed utilization, disease resistance and immune response



Improved feed management in RAS can offset the cost of the quality feeds required.

The potentially higher cost of the complete feeds needed for recirculating aquaculture systems (RAS) can be offset by improved feed management, since it is easier to feed frequently, place feed in the proximity of aquatic animals, and monitor uneaten or wasted feed in RAS. If such improved management is realized, the overall costs of feeding can actually decrease.

Overfeeding

Overfeeding is one of the major causes of environmental pollution resulting from aquaculture. This is a more significant problem for aquatic animals that require sinking feeds, such as shrimp, versus most fish, which need floating feeds. With shrimp, pond managers must estimate the correct amount of feed to deliver, since they cannot directly observe feed being consumed.

An increase in feeding frequency and reduction in volume per feeding is usually associated with a decrease in feed conversion ratios (FCR) or improvement in feed utilization. This is one of the major advantages associated with intensive RAS, since automatic feeders can be programmed to optimize feed utilization and reduce waste, improving on manual techniques. A parallel example from pond aquaculture is the use of feeding trays to quantify daily feeding behavior, adjust feeding rate and frequency, and minimize wasted feed.

Automated feeding systems

Many automated feeding systems for RAS and intensive cage systems that result in cost savings to producers have been developed in recent years. Improvements in feed use are associated with a subsequent reduction in feed waste and nutrient leaching.

Feed waste has been estimated at 25-75 percent, depending on the farming method. This waste should be drastically reduced in an RAS, since feeding behavior can be more easily assessed, the solids created by wasted food can be visually or automatically monitored, and the addition of food can be controlled by automated feeding systems.

A similar savings will be associated with leaching, since shorter but more frequent feeding periods result in shorter exposure of the feed to water before consumption, improving the nutrient availability. Finally, there are some reports that adding saponin compounds to the feed will actually result in reduced levels of ammonia in the culture water.

Attractants, incitants, and stimulants

Effective use of feeding attractants, incitants, or stimulants in RAS feeds may help reduce the use of expensive animal meals currently used, allow the reduction in protein level proposed in [Part 1](#) of this article, by standardizing palatability and reduce leaching and uneaten feed. These factors would subsequently further increase feed utilization and reduce feed waste.

Palatability

Differences in feed palatability have been demonstrated using different feed ingredients. For example, the protein requirement for *L. vannamei* was determined to be 15 percent in the presence of 2 percent artemia meal, but over 30 percent in the absence of artemia meal, emphasizing the need for adequate feed palatability.

Presently, many commercial feeds depend upon expensive squid meal, krill meal, fish solubles, and/or fishmeal to serve as “attractants.” Applied research is needed to assay aquatic chemoreception and develop cost-effective attractants for commercial feeds. These attractants should allow feed manufacturers to standardize palatability from one feed batch to another, without the market price of a single ingredient adversely affecting the selling price.

Improving disease resistance and immune response



Improved feed management in RAS can offset the cost of the quality feeds required.

Crowding has been shown to have a major correlation to disease outbreaks in aquatic animals, and RAS lead to much higher densities than traditional pond culture. Research suggests the addition of probiotics (e.g., carotenoids, nucleotides, vitamins, and β -glucans) can decrease such losses.

Probiotics

Vitamins E and C, selenium, and carotenoids increase disease resistance and/or immune response in several aquatic species. The use of docosahexaenoic acids and phospholipids for increased stress resistance, and dietary vitamin C for increased resistance to *Vibrio sp.* infections have been reported for aquatic animals.

Recently, the feed industry has begun to investigate the addition of β -glucans and various pigments to feed for the purpose of protecting aquatic animals from disease. The potential use of β -glucans has been established, but questions remain. When should they be used? How much should be used per

treatment? How many treatments are required per crop? What is the best treatment mode? Answers are needed before β -glucans can be added to feed cost-effectively.

Antibiotics

The addition of antibiotics to feed is a common practice that requires further research. The extended need for antibiotic treatment may be reduced in RAS, since acute treatment can more easily be accomplished due to the greater control and monitoring capabilities available in tanks versus ponds. Some antibiotics have also been found to act as growth enhancers, and are added as routine components of feed.

However, a significant consideration must be the effect of the antibiotics leaching from the feed on the biological filter bacteria. The nitrifying bacteria in these biological filters are susceptible to many common antibiotics. Misapplication of antibiotics can result in the loss of complete ammonia oxidation, leading to an acute rapid increase in total ammonia nitrogen and nitrite concentrations. The reestablishment of nitrifying bacteria can take several weeks; the added stress of increasing dissolved nitrogen concentrations on already diseased aquatic animals can be fatal.

This problem can be avoided by taking the filtration off-line during exposure to the medicated feed, and subsequently pumping the culture tank water back into the filtration system. The water can pass through a bed of activated carbon to absorb the antibiotic, and/or through an oxidizing environment like an ozone column to destroy the antibiotic. It is relatively easy to bioassay the effects of feed antibiotics on biological filter bacteria before adding the feed to large production systems.

Conclusion

The elements of feed management presented here include the most economically significant factors influencing the development and use of RAS feeds. However, research must be continued in many areas for truly cost-effective feeds and feed management strategies to be developed.

Future goals for feeds used in RAS should include:

- Lower feed ingredient costs (as a percentage of operating costs) due to improved species-specific formulations (i.e., improved assimilation and palatability).
- FCR as close to 1.0 as possible.
- Reduced environmental impact through better feed management and significantly reduced effluent discharge.
- Increased survival and disease resistance from improved feeds.
- Improved biosecurity of all production.

(Editor's Note: This article was originally published in the April 2001 print edition of the Global Aquaculture Advocate.)

Now that you've finished reading the article ...

... we hope you'll consider supporting our mission to document the evolution of the global aquaculture industry and share our vast network of contributors' expansive knowledge every week.

By becoming a Global Seafood Alliance member, you're ensuring that all of the pre-competitive work we do through member benefits, resources and events can continue. Individual membership costs just \$50 a year.

Not a GSA member? Join us.

[Support GSA and Become a Member](#)

Authors

-  Phillip G. Lee, Ph.D.

Phillip G. Lee, Ph.D.

Director, National Resource Center for Cephalopods
University of Texas Medical Branch
Galveston, Texas, USA

[117,100,101,46,98,109,116,117,64,101,101,108,103,112]

-  Addison L. Lawrence, Ph.D.

Addison L. Lawrence, Ph.D.

Reagent Professor and Project Leader
Texas A&M Shrimp Mariculture Project
Port Aransas, Texas, USA

[109,111,99,46,111,111,104,97,121,64,108,108,97,112,109,115]

Share

- [✉ Share via Email](#)
- [🐦 Share on Twitter](#)
- [f Share on Facebook](#)
- [in Share on LinkedIn](#)

Tagged With

[recirculating aquaculture systems](#) [Addison L. Lawrence](#) [Phillip G. Lee](#)

Related Posts

Innovation & Investment

[AquaMaof cultivating a ‘new generation’ of fish farmers](#)

Israel-based AquaMaof has seen its RAS technology rolled out to 10 facilities around the world, with more still in the pipeline.

Health & Welfare

[Bacterial pathogens in biofilms pose health risks in recirculating systems](#)

Recirculating aquaculture presents an increased potential for pathogenic bacteria to become established in the system through the formation of biofilms.

Responsibility

[Biofloc reduces feed, filtration costs in recirculating shrimp nursery system](#)

Trials with recirculating shrimp nursery systems in India supported the use of biofloc. To control ammonia is a way to reduce filtration costs.

Intelligence

[Full speed ahead for Florida pompano RAS farm](#)

While many attempts to farm Florida pompano have fallen short, Joe Cardenas is undaunted, confident that recirculating aquaculture systems are the key.

About The Advocate

The Responsible Seafood Advocate supports the Global Seafood Alliance's (GSA) mission to advance responsible seafood practices through education, advocacy and third-party assurances.

[Learn More](#)

Search Responsible Seafood Advocate



Advertising Opportunities

[2022 Media & Events Kit](#)

Categories

[Aquafeeds](#) [Aquafeeds](#) > [Health & Welfare](#) > [From Our Sponsors](#) > [Innovation & Investment](#) > [Intelligence](#) > [Responsibility](#) > [Fisheries](#) > [Artículos en Español](#) >

Don't Miss an Article

Featured

- [Health & Welfare](#) [An update on vibriosis, the major bacterial disease shrimp farmers face](#)
- [Intelligence](#) [A seat at the table: Fed By Blue team says aquaculture needs a stronger voice](#)
- [Responsibility](#) [Quantifying habitat provisioning at macroalgae cultivation locations](#)

Popular Tags

All Tags ▼




Recent

- [Fisheries](#) [Second Test: Another filler for the fisheries category](#)
- [Fisheries](#) [Test: This is filler for the fisheries Category](#)
- [Aquafeeds](#) [Test Article](#)
- [Responsibility](#) [Study: Climate change will shuffle marine ecosystems in unexpected ways as ocean temperature warms](#)
- [Health & Welfare](#) [Indian shrimp researchers earn a patent for WSSV diagnostic tool](#)



- [About](#)
- [Membership](#)
- [Resources](#)
- [Best Aquaculture Practices \(BAP\)](#)
- [Best Seafood Practices \(BSP\)](#)
- [GOAL Events](#)
- [Advocate Magazine](#)
- [Aquademia Podcast](#)
- [Blog](#)
- [Contact](#)

Stay up to date with GSA

- 
- 
- 
- 
- 

Copyright © 2024 Global Seafood Alliance
All rights reserved.

[Privacy](#)

[Terms of Use](#)

[Glossary](#)