



- [GOAL Events](#)
- [Advocate Magazine](#)
- [Aquademia Podcast](#)
- [Blog](#)
- [Contact](#)

-
-
-
-
-

- [Log In](#)



- [About](#)
 - [Who We Are](#)
 - [Our History](#)
 - [Our Team](#)
 - [Sustainable Development Goals](#)
 - [Careers](#)
- [Membership](#)
 - [Overview](#)
 - [Our Members](#)
 - [Corporate Membership](#)
- [Resources](#)
- [Certification](#)
 - [Best Aquaculture Practices](#)
 - [Best Seafood Practices](#)

[Log In](#)

- [About](#)
 - [Who We Are](#)
 - [Our History](#)
 - [Our Team](#)
 - [Sustainable Development Goals](#)
 - [Careers](#)
- [Membership](#)
 - [Overview](#)
 - [Our Members](#)
 - [Corporate Membership](#)
- [Resources](#)
- [Certification](#)
 - [Best Aquaculture Practices](#)
 - [Best Seafood Practices](#)
- [GOAL Events](#)
- [Advocate Magazine](#)

- [Aquademia Podcast](#)
- [Blog](#)
- [Contact](#)



 Aquafeeds
Aquafeeds

Feed study tests vegetable replacements for marine ingredients

 [Responsible Seafood Advocate logo](#)

1 April 2002 David Z. McMahon Bart Baca, Ph.D. Andrew Rogerson, Ph.D. Tzachi M. Samocha, Ph.D. John Reguzzoni



Experimental shrimp diet with low levels of vegetable protein and lipids produced acceptable results



Feed was manually broadcast to ponds during the study.

Most aquafeed manufacturers use expensive squid meal, fishmeal, and marine fish oils to meet protein requirements in their products. However, recent research has shown that vegetable and terrestrial animal proteins and oils could be used to successfully replace marine proteins and oils in feeds for many aquatic species, including penaeid shrimp.

Feed study

In late 2001, the authors conducted a 69-day grow-out trial in outdoor ponds at the OceanBoy Farms facility in Labelle, Florida, USA to compare the performance of various aquafeeds. OceanBoy is engaged in applied research to increase protein levels from vegetable sources in its shrimp feeds.

The diets were tested at varying stocking densities, ranging 33 to 138 shrimp per square meter. Animals used in the study were first reared for 21 days in a greenhouse nursery system and then moved to 12 outdoor, 1-ha, plastic-lined ponds with strong aeration and zero water exchange.

Water quality

Dissolved oxygen, pH, salinity, and temperature were monitored twice daily. Ammonia, nitrite, and nitrate levels were monitored weekly. All water-quality parameters measured remained within acceptable levels.

Feeds tested

Four diets were used in the study. Table 1 shows proximate analyses of the feeds described below.

- Diet 1: Experimental OceanBoy formulation with low levels of vegetable protein and lipids, and standard vitamin and enzyme input.
- Diet 2: Feed with lipids of vegetable origin and protein from vegetable and marine sources (OceanBoy formulation).
- Diet 3: Formulation provided by Dr. Allen Davis of Auburn University in Auburn, Alabama, USA with poultry meal instead of fishmeal and fish oil for palatability.
- Diet 4: Formulated by Robins McIntosh (formerly with Belize Aquaculture Ltd.), this diet contained medium levels of marine proteins and oils.

McMahon, Proximate analysis of selected feeds, Table 1

Crude Feed	Moisture	Protein	Crude Fiber	Ash	Crude Fat	Metabolize Energy
Diet 1	9.35	22.32	2.9	6.1	6.1	1,484
Diet 2	8.72	27.6	1.8	6.05	8.2	1,539
Diet 3	8.44	33.45	2.6	7.94	10.9	1,535
Diet 4	9.15	33.84	1.8	8.04	10.1	1,517

Table 1. Proximate analysis of selected feeds.

Results

Table 2 summarizes survival, growth, feed conversion and stocking density after 69 days of culture. As shown in the table, diet 4 produced the highest mean final weight (10.40 grams), followed by 6.97 grams for diet 3, 6.23 grams for diet 2, and 5.40 grams for diet 1. Diets 3 and 4 produced survival results of about 87 percent, closely followed by diet 2 and then diet 1 at 73 percent. The lowest feed conversion ratio (1.17) was obtained with diet 2.

McMahon, Summary of survival, growth, feed conversion, Table 2

Feed	Survival (%)	Average Weight (g)	Feed-Conversion Ratio	Stocking Density (animals/m²)
Diet 1	60.0	5.8	5.5	33.2
	71.0	6.0	2.1	70.6
	89.0	4.4	2.4	67.5
Diet 2	85.0	5.9	1.2	106
	83.0	6.3	1.5	84
	88.0	6.5	0.8	116.5
Diet 3	7.0	7.0	1.9	111.7
	7.6	7.6	1.9	137.8
	6.3	6.3	2.1	84
Diet 4	12.5	12.5	3.8	99.9
	8.7	8.7	4.2	45.8
	10.0	10.0	8.3	131.6

Table 2. Summary of survival, growth, feed conversion, and stocking density after 69 days of culture.

Conclusion

The study of feed performance showed that an experimental diet with low levels of vegetable protein and lipids, although ranking last in growth and survival among the diets tested, still produced acceptable results. Feed protein requirements can effectively be reduced and vegetable protein sources increased in shrimp feeds.

(Editor's Note: This article was originally published in the April 2002 print edition of the Global Aquaculture Advocate.)

Now that you've finished reading the article ...


... we hope you'll consider supporting our mission to document the evolution of the global aquaculture industry and share our vast network of contributors' expansive knowledge every week.

By becoming a Global Seafood Alliance member, you're ensuring that all of the pre-competitive work we do through member benefits, resources and events can continue. Individual membership costs just \$50 a year.

Not a GSA member? Join us.

[Support GSA and Become a Member](#)

Authors

-  David Z. McMahon

David Z. McMahon

Nova Southeastern University
Oceanographic Center
8000 North Ocean Drive
Dania, Florida 33004 USA

-  Bart Baca, Ph.D.


Bart Baca, Ph.D.

Nova Southeastern University
Oceanographic Center
8000 North Ocean Drive
Dania, Florida 33004 USA

-  Andrew Rogerson, Ph.D.

Andrew Rogerson, Ph.D.

Nova Southeastern University
Oceanographic Center
8000 North Ocean Drive
Dania, Florida 33004 USA

-  Tzachi M. Samocha, Ph.D.

Tzachi M. Samocha, Ph.D.





Texas Agricultural Experiment Station
Shrimp Mariculture Research Facility
Corpus Christi, Texas, USA

-  John Reguzzoni

John Reguzzoni

OceanBoy Farms, Inc.
Clewiston, Florida, USA

Share

-  [Share via Email](#)
-  [Share on Twitter](#)
-  [Share on Facebook](#)
-  [Share on LinkedIn](#)

Tagged With

[Tzachi M. Samocha](#) [vegetable replacements](#) [David Z. McMahon](#) [Bart Baca](#) [Andrew Rogerson](#) [John Reguzzoni](#) [Shrimp](#)

Related Posts

[Aquafeeds](#)

[Fish oil alternatives are waiting in the wings](#)

The Future of Fish Feed (F3) initiative aims to jumpstart innovation in terms of viable fish-free substitutes to fish oil through the global F3 Fish Oil Challenge.

Health & Welfare

[Fishmeal replacement in cobia diets reduces costs, improves sustainability](#)

Scientists have been working to understand cobia's nutritional requirements and advance the economic and environmental sustainability of feeding carnivorous fish using fishmeal alternatives.

Aquafeeds

[Fishmeal-free feeds for hybrid tilapia](#)

A series of experiments showed that fishmeal is not an essential ingredient in hybrid tilapia feeds and that plant protein-based diets can yield results similar to those obtained with diets containing 10 percent fishmeal.

Aquafeeds

[Meta-analysis examines ingredient substitution for fishmeal in salmon, trout feeds](#)

In a meta-analysis of 28 nutrition studies, the authors found limited possibilities for ingredient substitution for fishmeal in diets for salmon and trout.

About The Advocate

The Responsible Seafood Advocate supports the Global Seafood Alliance's (GSA) mission to advance responsible seafood practices through education, advocacy and third-party assurances.

[Learn More](#)

Search Responsible Seafood Advocate



Advertising Opportunities

[2022 Media & Events Kit](#)

Categories

[Aquafeeds](#) [Aquafeeds](#) > [Health & Welfare](#) > [From Our Sponsors](#) > [Innovation & Investment](#) > [Intelligence](#) > [Responsibility](#) > [Fisheries](#) > [Artículos en Español](#) >

Don't Miss an Article

Featured

- [Health & Welfare](#) [An update on vibriosis, the major bacterial disease shrimp farmers face](#)
- [Uncategorized](#) [A seat at the table: Fed By Blue team says aquaculture needs a stronger voice](#)
- [Responsibility](#) [Quantifying habitat provisioning at macroalgae cultivation locations](#)

Popular Tags

All Tags





Recent

- [Fisheries Second Test: Another filler for the fisheries category](#)
- [Fisheries Test: This is filler for the fisheries Category](#)
- [Aquafeeds Test Article](#)
- [Responsibility Study: Climate change will shuffle marine ecosystems in unexpected ways as ocean temperature warms](#)
- [Health & Welfare Indian shrimp researchers earn a patent for WSSV diagnostic tool](#)



- [About](#)
- [Membership](#)
- [Resources](#)
- [Best Aquaculture Practices \(BAP\)](#)
- [Best Seafood Practices \(BSP\)](#)
- [GOAL Events](#)
- [Advocate Magazine](#)
- [Aquademia Podcast](#)
- [Blog](#)
- [Contact](#)

Stay up to date with GSA

- 
- 
- 
- 
- 

Copyright © 2024 Global Seafood Alliance

All rights reserved.

[Privacy](#)

[Terms of Use](#)

[Glossary](#)