



- [GOAL Events](#)
- [Advocate Magazine](#)
- [Aquademia Podcast](#)
- [Blog](#)
- [Contact](#)
  
- 
- 
- 
- 
- 
  
- [Log In](#)



- [About](#)
  - [Who We Are](#)
  - [Our History](#)
  - [Our Team](#)
  - [Sustainable Development Goals](#)
  - [Careers](#)
- [Membership](#)
  - [Overview](#)
  - [Our Members](#)
  - [Corporate Membership](#)
- [Resources](#)
- [Certification](#)
  - [Best Aquaculture Practices](#)
  - [Best Seafood Practices](#)

Search...



[Log In](#)

- [About](#)
  - [Who We Are](#)
  - [Our History](#)
  - [Our Team](#)
  - [Sustainable Development Goals](#)
  - [Careers](#)
- [Membership](#)
  - [Overview](#)
  - [Our Members](#)
  - [Corporate Membership](#)
- [Resources](#)
- [Certification](#)
  - [Best Aquaculture Practices](#)
  - [Best Seafood Practices](#)
- [GOAL Events](#)
- [Advocate Magazine](#)
- [Aquademia Podcast](#)
- [Blog](#)
- [Contact](#)



Health & Welfare  
Health & Welfare


Genetic improvement of marine fish



1 February 2002 Wayne Knibb, Ph.D.



## Methods, values for sea bream, other species



One avenue of genetic improvement is mutation. The “ebony” mutation expressed in the sea bream with dark tails above is a form of heterotic growth that is also used for commercial protection of the improved strains.

Many aquaculture industries fail to enhance productivity and competitiveness with genetic improvement. This may be due, in part, to uncertainty over which methods should be used. Indeed, genetics comprises many subdisciplines, each with advocates who promote one field or another as the best approach. For the Mediterranean sea bream, as well as other species, various genetic methods have been employed to advance commercial value.

## Benefits of genetic improvement

Genetic improvement programs can shape cultured animals to suit a given purpose or environment – as is evident, for example, in the variety of the foods we consume. Genetic programs can produce many benefits, including important economic ones.

### *Increased productivity, reliability*

Genetic improvement is often the primary method used to increase productivity in terrestrial agriculture. Nearly all the 300-400 percent growth improvement of broiler chicken strains since the 1950s was due to selection. Faster growth translates into greater production with existing facilities and, often, improved food-conversion efficiency.

Today, chicken is inexpensive, of uniform quality and appearance, and reliable in supply. Unfortunately, the reverse is typical of many commercial species of marine fish.

### *Increased profitability, new industry*

Long-term profit margins for commodity aquaculture companies are usually narrow. For marginal companies that operate with 5 percent profit, even modest genetic gains of 10 percent can double profits. For sea bream, where profitability is constrained by environmental regulations and limits on sea cages, genetically improved strains can increase production with existing infrastructure. In addition, new “seed” industries that produce eggs and fingerlings for local or overseas markets can develop from genetic improvement programs.

### *Resistance to pathogens*

The emergence of new pathogens is almost predictable in farmed Mediterranean sea bream and European sea bass. The development of fish strains that are genetically resistant to important pathogens is one approach to this problem.

## Improved ROI

Genetic gains compound with each generation of selection, and economic value is proportional to the size of the industry. It is possible to calculate with some precision the return on investment for genetic breeding programs (e.g., 1:15 for salmon, 1:50 for pigs).

## Selection among strains

Strains differ in their growth and survival performance. The choice of the best existing strains could equal the genetic gains made by years of within-strain selection using inferior strains. Hence, strain testing should precede within-strain selection.

Most finfish species, including sea bream, carp, and tilapia, show some strain variation for growth, disease resistance, or cold tolerance. To varying degrees, these differences have been exploited in commercial farms (Table 1).

**Table 1. Although each species reflects constraints, most finfish show strain variation with commercial potential.**

Method	Species	Genetic Gain	Cost	\$ Benefit	\$ Risk	Major Constraints*
Between-Strain Selection	Carp	++	++	+++	++	Disease propagation among strains
	Tilapia	++	++	+	++	
	Sea Bream	+	++	+	+++	
Within-Strain Selection	Carp	++	+++	+++	++	Long lag times (up to 10 years) Undesirable correlated selection response
	Tilapia	++	+++	+++	++	
	Sea Bream	+++	+++	+++	+	
Mutations of Large Effect/Qualitative Traits	Carp	++	+	++	+++	Unpredictability
	Tilapia	++	+	++	+++	
	Sea Bream	++	+	++	+++	
Crossbreeding Among Strains	Carp	++	++	++	++	Disease propagation
	Tilapia	+	++	+	+++	
	Sea Bream	+	++	+	+++	
Hybridization Between Species	Carp	++	+	+	+++	Reproduction Acceptance Unpredictability
	Tilapia	++	+	++	+++	
	Sea Bream	+	+	+	+++	
Chromosome Manipulation	Carp	+	+	+	+++	Low GSI values
	Tilapia	+	+	+	+++	
	Sea Bream	+	+	+	+++	
Genetic Engineering	Carp	++	++	+	+++	Integration Acceptance Growout fitness
	Tilapia	++	++	+	+++	
	Sea Bream	+			+++	

+ = low potential, ++ = medium potential, +++ = high potential. \* Potential constraints common to all: ecological, strain testing, and commercial protection.

## Selection within strains

Within-strain selection can have various objectives (e.g., traits that contribute to profit) using different selection criteria. Individuals can be ranked with information from themselves, from their relatives, or using additional information.

### Genetic differences, gains

Selection response cannot occur unless there are genetic differences among individuals that can be inherited by offspring. Luckily, most populations have genetic differences for commercial traits. For example, in Israel, an industrial mass-selection program increased body weight in sea bream 5 to 10 percent per generation.

Elsewhere, selection response has been achieved for disease resistance, body weight, and body proportion in carp, and body weight in tilapia. Most selection experiments for body weight in finfish have yielded gains of 10-20 percent per generation.

### Optimize accuracy

It is generally desirable to optimize the accuracy of selection and increase response. However, optimization can incur high costs for tagging and management to obtain pedigree information. Cost-benefit analyses of genetic options for each species should be conducted to calculate total costs and gains from different types of selection.

### Business evaluation

In addition to genetic constraints, the business evaluation of selection options should consider temporal, biological, and technical constraints. Without feasibility studies, businesses should adopt proven genetic technologies, rather than technologies without precedent of commercial success.

## Traits, genes, and mutations

Unlike growth, thought to be due to many loci with minor effects, some traits are due to one or few loci. Various single-gene mutations, including ebony, yellow, and white body color, have been isolated for sea bream. The ebony mutation is heterotic growth, used for commercial protection of the improved strains. However, overall, there are few qualitative traits of commercial value in foodfish aquaculture. Examples include red body color in tilapia and scale patterns in carp.

## Crossbreeding

In terrestrial agriculture, crossbreeding between different strains of the same species can produce heterosis and uniformity. For sea bream, commercial crossbreeding is used to limit inbreeding in selection lines. Indeed, for most aquaculture applications, crossbreeding should be considered simply as the reverse of inbreeding, rather than as a method to achieve gains over the base outbred population.

## Hybrids

Hybrids between species are created to combine parental traits like fast growth and tolerance to disease or temperature, or produce single-sex offspring. They can also be used to achieve superior performance or sterility.

Hybrids usually resemble the average of their parents. Given the large number of attempts to produce new fish hybrids, remarkably few (apparently less than 1 percent, and none for sea bream) have resulted in sustained commercial production.

## Chromosome manipulation

The potential applications for chromosome set manipulations in fish are diverse. They include growth acceleration (with triploids via suppression of gonadal development), same-sex offspring, rapid inbreeding, and restoration of whole fish from cryopreserved sperm.

However, in practice, triploid sea bream have little growth acceleration. A similar conclusion has been reached for triploid carp. Many fish triploids have been produced, but except for some salmonids, little information exists regarding their commercial performance, and few are actually used by farmers. Even so, ecological issues may sustain future research on sterile and triploid fish.

Managed genetics programs usually minimize inbreeding to avoid mutations like those reflected in these sea bream.

## Genetic engineering

Presently, there is no commercial use of genetically engineered fish. But genetic engineering could be used to minimize inbreeding in wild and captive populations.

Without formal genetics programs, local industries often attempt selection in house, with few broodstock individuals and small effective population sizes. Even without deliberate selection, farm management typically results in severe inbreeding and domestication selection. Managed genetics programs usually attempt to minimize inbreeding.

## Conclusion

In theory, there are many methods to genetically improve fish strains. In practice, few improved strains have reached the industry, and of these, most have been developed with traditional selection that exploits additive genetic variance. Those strains that have reached the aquaculture industry, however, have been of substantial economic value.

*(Editor's Note: This article was originally published in the February 2002 print edition of the Global Aquaculture Advocate.)*

## Now that you've finished reading the article ...

... we hope you'll consider supporting our mission to document the evolution of the global aquaculture industry and share our vast network of contributors' expansive knowledge every week.

By becoming a Global Seafood Alliance member, you're ensuring that all of the pre-competitive work we do through member benefits, resources and events can continue. Individual membership costs just \$50 a year.

**Not a GSA member? Join us.**

[Support GSA and Become a Member](#)

#### Author





-  Wayne Knibb, Ph.D.

Wayne Knibb, Ph.D.

Bribie Island Aquaculture Research Centre  
144 North Street, Woorim  
P.O. Box 2066  
Bribie Island  
Queensland 4507 Australia

[46,68,46,104,80,32,44,98,98,105,110,75,32,101,110,121,97,87]

#### Share

-  [Share via Email](#)
-  [Share on Twitter](#)
-  [Share on Facebook](#)
-  [Share on LinkedIn](#)

#### Tagged With

[genetic improvement](#) [Wayne Knibb](#)

#### Related Posts

##### Intelligence

#### [Byproduct utilization for increased profitability, part 5](#)

Fish protein hydrolysates have been used in the food industry for the development of antioxidants, emulsifiers, flavoring agents, anti-bacterial agents and feed. As natural products, they have a growing acceptance, since natural products are preferred by some consumers over industrial products.

##### Aquafeeds

#### [Feed study examines effects of fishmeal storage](#)

A study investigated the effects of storage temperature and duration on the quality of finished fishmeal fed to juvenile shrimp. Diets with 15 percent fishmeal were stored at 4 or 30 degrees-C for six months or 12 months.

##### Innovation & Investment

#### [Unleashing the genetic potential of black tiger shrimp](#)

The black tiger shrimp is the second most important, farmed shrimp species globally. Because of its importance in Australia, the new ARC Hub for Advanced Prawn Breeding will develop a scalable, advanced breeding program for the species as well as bring its genetic knowledge to a level comparable to that of livestock.

##### Health & Welfare

#### [Screening probiotic bacteria for Atlantic cod larvae](#)

The addition of probiotic bacteria to marine larvae can reduce pathogenic bacteria and subsequent infection by establishing a protective flora on the mucosal surfaces of the larvae.

#### About The Advocate

The Responsible Seafood Advocate supports the Global Seafood Alliance's (GSA) mission to advance responsible seafood practices through education, advocacy and third-party assurances.

[Learn More](#)

Search Responsible Seafood Advocate



## Advertising Opportunities

[2022 Media & Events Kit](#)

## Categories

[Aquafeeds](#) > [Health & Welfare](#) [Health & Welfare](#) > [From Our Sponsors](#) > [Innovation & Investment](#) > [Intelligence](#) > [Responsibility](#) > [Fisheries](#) > [Artículos en Español](#) >

## Don't Miss an Article

## Featured

- [Health & Welfare](#) [An update on vibriosis, the major bacterial disease shrimp farmers face](#)
- [Uncategorized](#) [A seat at the table: Fed By Blue team says aquaculture needs a stronger voice](#)
- [Responsibility](#) [Quantifying habitat provisioning at macroalgae cultivation locations](#)

## Popular Tags

All Tags ▼




## Recent

- [Fisheries](#) [Second Test: Another filler for the fisheries category.](#)
- [Fisheries](#) [Test: This is filler for the fisheries Category.](#)
- [Aquafeeds](#) [Test Article](#)
- [Responsibility](#) [Study: Climate change will shuffle marine ecosystems in unexpected ways as ocean temperature warms](#)
- [Health & Welfare](#) [Indian shrimp researchers earn a patent for WSSV diagnostic tool](#)



- [About](#)
- [Membership](#)
- [Resources](#)
- [Best Aquaculture Practices \(BAP\)](#)
- [Best Seafood Practices \(BSP\)](#)
- [GOAL Events](#)
- [Advocate Magazine](#)
- [Aquademia Podcast](#)
- [Blog](#)
- [Contact](#)

#### Stay up to date with GSA

- 
- 
- 
- 
- 

Copyright © 2024 Global Seafood Alliance  
All rights reserved.

[Privacy](#)

[Terms of Use](#)

[Glossary](#)