



- [GOAL Events](#)
- [Advocate Magazine](#)
- [Aquademia Podcast](#)
- [Blog](#)
- [Contact](#)

- 
- 
- 
- 
- 

- [Log In](#)



- [About](#)
  - [Who We Are](#)
  - [Our History](#)
  - [Our Team](#)
  - [Sustainable Development Goals](#)
  - [Careers](#)
- [Membership](#)
  - [Overview](#)
  - [Our Members](#)
  - [Corporate Membership](#)
- [Resources](#)
- [Certification](#)
  - [Best Aquaculture Practices](#)
  - [Best Seafood Practices](#)

Search...



[Log In](#)

- [About](#)
  - [Who We Are](#)
  - [Our History](#)
  - [Our Team](#)
  - [Sustainable Development Goals](#)
  - [Careers](#)
- [Membership](#)
  - [Overview](#)
  - [Our Members](#)
  - [Corporate Membership](#)
- [Resources](#)
- [Certification](#)
  - [Best Aquaculture Practices](#)
  - [Best Seafood Practices](#)
- [GOAL Events](#)
- [Advocate Magazine](#)
- [Aquademia Podcast](#)
- [Blog](#)
- [Contact](#)



[Health & Welfare](#)  
Health & Welfare

## Genetic selection can increase feed efficiency of salmon



1 December 2001 Bjarne Gjerde Jørn Thodesen Barbara Grisdale-Helland



## Can salmon farmers follow poultry's lead?



Waste feed collection for determining daily feed intake.

Feed is the largest variable cost in intensive production of farmed fish. In Norway, for example, feed constitutes more than 50 percent of the cost of Atlantic salmon production. Increased feed efficiency is therefore of great importance in order to increase economic margins and improve production in situations with limited feed resources. Greater efficiency can be obtained through improved feeds and production management, but also through direct or indirect genetic selection for increased feed utilization.

Direct selection is not currently possible, because accurate feed intake measurements are not feasible in largescale breeding programs with present technology, or are very costly to measure. However, selection for increased growth rate will result in an indirect (correlated) response for increased feed efficiency. This has proven to be the case for poultry, swine, sheep, and cattle; and also in a study with Atlantic salmon in Norway.

### Improved feed efficiency

Salmon selected for increased growth rate over five generations were compared with nonselected wild salmon in a 14-week growth study (Thodesen et al., 1999). The selected salmon had 40 percent higher relative feed intake per unit of body weight per day, 113 percent higher growth rate, and 25 percent higher feed efficiency ratio. Given Norway's current salmon production of 450,000 metric tons (MT), this improvement in feed efficiency could be worth approximately U.S. \$110 million.

### Wild vs. selected salmon

The difference in feed efficiency between the selected and wild salmon indicated a correlated response in feed efficiency of 4.6 percent per generation. Therefore, the higher growth rate of the selected salmon was the result of both greater feed consumption (larger appetite) and more efficient utilization of the feed for growth. The main reason for this was an increased amount of metabolizable energy available for gain relative to the energy needed for maintenance. Feed Efficiency and Growth Rates In the above experiment, feed efficiency ratio showed a high positive correlation with growth rate ( $r = 0.90$ ). This strong association between feed efficiency and growth rate in Atlantic salmon was recently confirmed in a 10-week growth experiment with 14 replicated full-sib family groups (Thodesen et al., 2001), as shown in Fig. 1 ( $r = 0.79$ ).




Fig. 1: Relationship between growth rate and feed efficiency.

This relationship, however, was not linear. Increasing growth rate was associated with a decreasing response in feed efficiency ratio. This suggests that at some stage, it will be more beneficial to select directly for improved feed efficiency in Atlantic salmon, as has earlier been concluded for poultry.

## Energy component of weight gain

Results from several experiments with Atlantic salmon indicate a significant genetic variation in deposition of energy in the carcass and viscera per unit of body weight gain. Feed efficiency can therefore be improved indirectly by reducing the energy content of the weight gain, assuming the energy content is higher than optimum. Thus, increased feed efficiency in Atlantic salmon may be achieved by selection for increased growth rate or less energy in weight gain.

However, this indirect selection response will decline with increasing growth rate and when the energy in the weight gain has reached an optimum level. Further improvement in feed efficiency must be obtained by a direct selection procedure. This will require new and cheaper technology that can be used to measure feed intake of individual fish or full-sib family groups. Development of such technology should be encouraged.

## Conclusion

Genetic selection of salmon for growth rate is indirectly improving feed efficiency. Additional indirect gains can be achieved through reduction in the energy content of weight gain. Ultimately, technology will be required for direct selection based on feed efficiency.

*Note: Cited references are available from the first author.*

*(Editor's Note: This article was originally published in the December 2001 print edition of the Global Aquaculture Advocate.)*

## Now that you've finished reading the article ...




... we hope you'll consider supporting our mission to document the evolution of the global aquaculture industry and share our vast network of contributors' expansive knowledge every week.

By becoming a Global Seafood Alliance member, you're ensuring that all of the pre-competitive work we do through member benefits, resources and events can continue. Individual membership costs just \$50 a year.

**Not a GSA member? Join us.**

[Support GSA and Become a Member](#)

## Authors

-  Bjarne Gjerde  
Bjarne Gjerde  
AKVAFORSK  
P.O. Box 5010, N-1432 Ås  
Norway
-  Jørn Thodesen  
Jørn Thodesen  
ICLARM, Penang  
Malaysia
-  Barbara Grisdale-Helland  
Barbara Grisdale-Helland  
AKVAFORSK  
Sunnalsøra  
Norway

## Share

- [✉ Share via Email](#)
- [🐦 Share on Twitter](#)
- [f Share on Facebook](#)
- [in Share on LinkedIn](#)

## Tagged With

[Bjarne Gjerde](#) [feed efficiency of salmon](#) [Jørn Thodesen](#) [Barbara Grisdale-Helland](#)

## Related Posts

### Intelligence

#### [Better to eat – farmed or wild salmon?](#)

There is a widely promoted misconception that eating wild-caught fish is better for the oceans than eating farmed seafood. On a global basis, however, sustainably farmed fish may represent 60 times more efficient use of anchovies and other baitfish resources than wild fish.

### Aquafeeds

#### [Feed Innovation Network: A streamlined aquafeed ingredient evaluation tool](#)

To position aquaculture for future growth, it needs tools to evaluate a growing spectrum of alternative ingredients and formulated diets designed to optimize fish health and efficient production.

### Health & Welfare

#### [Chile's salmon industry addresses health crises](#)

Environmental and fish health problems have affected Chile's salmon farmers since 2004. As production ramped up, 2006 saw an increase in sea lice that reduced output. In 2007, infectious salmon anemia spread across the farming region and quickly cut production.

### Innovation & Investment

#### [AquaGen CEO: Genomics are transforming aquaculture](#)

The CEO of AquaGen knew that the Norwegian research group's work in genomics was key to the salmon industry's future. And that was before she even worked there.

## About The Advocate

The Responsible Seafood Advocate supports the Global Seafood Alliance’s (GSA) mission to advance responsible seafood practices through education, advocacy and third-party assurances.

[Learn More](#)

Search Responsible Seafood Advocate



**Advertising Opportunities**

[2022 Media & Events Kit](#)

**Categories**

[Aquafeeds](#) > [Health & Welfare](#) [Health & Welfare](#) > [From Our Sponsors](#) > [Innovation & Investment](#) > [Intelligence](#) > [Responsibility](#) > [Fisheries](#) > [Artículos en Español](#) >

**Don't Miss an Article**

**Featured**

- [Health & Welfare](#) [An update on vibriosis, the major bacterial disease shrimp farmers face](#)
- [Uncategorized](#) [A seat at the table: Fed By Blue team says aquaculture needs a stronger voice](#)
- [Responsibility](#) [Quantifying habitat provisioning at macroalgae cultivation locations](#)

**Popular Tags**

All Tags






**Recent**

- [Fisheries](#) [Second Test: Another filler for the fisheries category.](#)
- [Fisheries](#) [Test: This is filler for the fisheries Category.](#)
- [Aquafeeds](#) [Test Article](#)
- [Responsibility](#) [Study: Climate change will shuffle marine ecosystems in unexpected ways as ocean temperature warms](#)
- [Health & Welfare](#) [Indian shrimp researchers earn a patent for WSSV diagnostic tool](#)



- [About](#)
- [Membership](#)
- [Resources](#)
- [Best Aquaculture Practices \(BAP\)](#)
- [Best Seafood Practices \(BSP\)](#)
- [GOAL Events](#)
- [Advocate Magazine](#)
- [Aquademia Podcast](#)
- [Blog](#)
- [Contact](#)

**Stay up to date with GSA**

- 
- 
- 
- 
- 

Copyright © 2024 Global Seafood Alliance  
All rights reserved.

- [Privacy](#)
- [Terms of Use](#)
- [Glossary](#)