Global Seafood Alliance Logo

- GOAL Events
- Advocate Magazine
- <u>Aquademia Podcast</u>
- <u>Blog</u>
- <u>Contact</u>
- 0
- ·f
- 🕅
- in
- 🕨
- <u>Log In</u>

- <u>About</u>
 - <u>Who We Are</u>
 - <u>Our History</u>
 - <u>Our Team</u>
 - Sustainable Development Goals
 - <u>Careers</u>
 - <u>Membership</u>
 - <u>Overview</u>
 - Our Members
 - <u>Corporate Membership</u>
- <u>Resources</u>
- <u>Certification</u>
 - Best Aquaculture Practices
 - Best Seafood Practices

Search...
Q

<u>Log In</u>

- <u>About</u>
 - Who We Are
 - <u>Our History</u>
 - <u>Our Team</u>
 - Sustainable Development Goals
 - <u>Careers</u>
- <u>Membership</u>
 - <u>Overview</u>
 - Our Members
 - Corporate Membership
- <u>Resources</u>
- <u>Certification</u>
 - Best Aquaculture Practices
 - Best Seafood Practices
- GOAL Events
- Advocate Magazine
- <u>Aquademia Podcast</u>
- <u>Blog</u>
- Contact



Health & Welfare

Gracilaria cultivation can provide bioremediation in Chinese mariculture

Responsible Seafood Advocate logo

1 January 2011 Dr. Yufeng Yang Dr. Charles Yarish



Seaweed can decrease nitrogen and phosphorus loadings



Gracilaria seaweed is effective at removing inorganic nutrients from water.

Within China's mariculture sector, culture of the red agarophyte *Gracilaria* has rapidly expanded over the past 10 years. Production of *Gracilaria* reached 99,451 metric tons in 2007 and for seaweeds ranked only behind the kelps *Saccharina* and *Undaria*.

The principal *Gracilaria* species cultured throughout China is *G. lemaneiformis*. Growth rates for the seaweed range up to 13.9 percent/day in Jiaozhou Bay in Shandong Province.

G. lemaneiformis is very effective in decreasing nitrogen and phosphorus loadings. The seaweed is also able to inhibit the growth of some microalgae and may increase dissolved oxygen in the water column. Large-scale *Gracilaria* cultivation can be an effective means of improving water quality and promoting a more sustainable mariculture industry in China.

Bioremediation benefits

Nanao is an island county of Guangdong Province with a population of about 70,000. Of these, about 5,000 people are now engaged in the cultivation of *Gracilaria*. The area of cultivation rose from 0.06 hectares (ha) in 1999 to 800 ha in 2006. The seaweed provides several beneficial functions.

Biofiltration

The rapid development of the mariculture industry has aroused concerns about the effects of these activities on the Chinese coastal environment, which can include deterioration of water quality and an increase in contaminants.

Mesocosm experiments demonstrated that *G. lemaneiformis* can effectively remove inorganic nutrients from water. Concentrations of ammonium nitrogen decreased by 85.53 and 69.45 percent, and concentrations of phosphate decreased by 65.97 and 26.74 percent in the mesocosms with *Gracilaria* in comparison to mesocosms without the seaweed. In 24-hour enclosure experiments, *Gracilaria* removed 68.44 percent of ammonium nitrogen, 23.03 percent of nitrate nitrogen and 13.04 percent of nitrite nitrogen.

Gracilaria cultivation can provide bioremediation in Chinese mariculture - Responsible Seafood Advocate

The maximum uptake rates of nitrate nitrogen, ammonium nitrogen and phosphate by G. lichenoides were 55.88, 35.17 and 3.106 umol/g/h, respectively. The corresponding rates for *G. lemaneiformis* were 53.17, 32.24 and 3.064 umol/g/h, respectively. These studies confirmed that *Gracilaria* species are good candidates for nutrient removal.

Increased D.O. concentrations

Testing during 17 visits to the Shenao culture area in Nanao showed that dissolved-oxygen (D.O.) levels were highest in cages with Gracilaria, second highest in the surrounding sea water outside the cages and lowest in cages with fish.

A 12-day experiment found that concentrations of D.O. were always higher in 1-m³ mesocosms with *Gracilaria* than those without it (Fig. 1). These results demonstrated that cultivated *Gracilaria* is very effective in improving D.O. levels in mariculture areas.



Fig. 1: Dissolved-oxygen concentrations in mesocosms with and without Gracilaria.

Decreased microalgae densities

The mesocosm experiments also demonstrated that *G. lemameiformis* limited microalgae growth. The densities of phytoplankton increased from 3.017×10^4 to 105.500×10^4 cells/L in the mesocosms without *Gracilaria*, whereas the densities increased from 2.387×10^4 to 26.500×10^4 cell/L in those with *Gracilaria*. The densities of phytoplankton were always lower in the mesocosms with *Gracilaria* (Fig. 2).



Fig. 2: Phytoplankton densities in mesocosms with and without Gracilaria.

Other experiments demonstrated that the fresh thalli of *G. lemaneiformis* significantly inhibited the growth of selected microalgae: *Prorocentrum donghaiense*, *Alexandrium tamarense*, *Amphidinium cartera*, *Scrippsiella trochoide* and *Chaetotoceros curvisetus*.

The results showed that *Gracilaria* can suppress growth and decrease densities of these microalgae. Large-scale cultivation of *Gracilaria* may be an effective ecological strategy to control harmful algal blooms in Chinese coastal waters.

Gracilaria cultivation can help control harmful algae blooms.

(Editor's Note: This article was originally published in the January/February 2011 print edition of the Global Aquaculture Advocate.)

Authors

• Dr. Yufeng Yang

Dr. Yufeng Yang

College of Life Science and Technology Jinan University Guangzhou, China

• 💭 Dr. Charles Yarish

Dr. Charles Yarish

Department of Ecology and Evolutionary Biology University of Connecticut Stamford, Connecticut, USA 75 North Eagleville Road Unit 3043 Storrs, Connecticut 06269 USA

[117,100,101,46,110,110,111,99,117,64,104,115,105,114,97,121,46,115,101,108,114,97,104,99]

Share

• <u>Share via Email</u>

- **J** Share on Twitter
- **F** Share on Facebook
- in <u>Share on LinkedIn</u>

Tagged With

Yufeng Yang Charles Yarish bioremediation nitrogen China phosphorous microalgae gracilaria D.O.

Related Posts

Responsibility

A look at integrated multi-trophic aquaculture

In integrated multi-trophic aquaculture, farmers combine the cultivation of fed species such as finfish or shrimp with extractive seaweeds, aquatic plants and shellfish and other invertebrates that recapture organic and inorganic particulate nutrients for their growth.

Health & Welfare

<u>A study of Zoea-2 Syndrome in hatcheries in India, part 3</u>

In this third and final part, authors present recommendations to help reduce the incidence of Zoea-2 Syndrome, which is not caused by any known infectious agents in P. vannamei hatcheries in India.

Innovation & Investment

American Unagi brings eel farming back 'home'

Sara Rademaker launched American Unagi to shift eel farming to American soil, where the eels are from. Why? Because of the novelty, and because she saw an opportunity to do things better.

Responsibility

An African prison where fish will be farmed and lives saved

A nonprofit organization working to improve conditions in African prisons is hoping tilapia ponds, tended to by inmates, will aid in their nutrition. A small donation from the Global Aquaculture Alliance will go a long way.

About The Advocate

The Responsible Seafood Advocate supports the Global Seafood Alliance's (GSA) mission to advance responsible seafood practices through education, advocacy and third-party assurances.

Learn More

Search Responsible Seafood Advocate Search Search Q





Listen to the seafood industry's top podcast

Advertising Opportunities

2022 Media & Events Kit

Categories

Aquafeeds > Health & Welfare Health & Welfare > From Our Sponsors > Innovation & Investment > Intelligence > Responsibility > Fisheries > Artículos en Español >

Don't Miss an Article

Featured

- Health & Welfare An update on vibriosis, the major bacterial disease shrimp farmers face
- Intelligence A seat at the table: Fed By Blue team says aquaculture needs a stronger voice
- <u>Responsibility Quantifying habitat provisioning at macroalgae cultivation locations</u>

Popular Tags

All Tags 🗸

Recent

- Fisheries Second Test: Another filler for the fisheries category
- Fisheries Test: This is filler for the fisheries Category
- <u>Aquafeeds Test Article</u>
- <u>Responsibility Study: Climate change will shuffle marine ecosystems in unexpected ways as ocean temperature warms</u>
- Health & Welfare Indian shrimp researchers earn a patent for WSSV diagnostic tool





Listen to the seafood industry's top podcast

- <u>About</u>
- <u>Membership</u>
- <u>Resources</u>
- Best Aquaculture Practices (BAP)
- Best Seafood Practices (BSP)
- GOAL Events
- <u>Advocate Magazine</u>
- <u>Aquademia Podcast</u>
- <u>Blog</u>
- <u>Contact</u>

Stay up to date with GSA

- 🞯
- f
- X
- in
- •

Copyright © 2024 Global Seafood Alliance All rights reserved. <u>Privacy</u> <u>Terms of Use</u> <u>Glossary</u>