



- [GOAL Events](#)
- [Advocate Magazine](#)
- [Aquademia Podcast](#)
- [Blog](#)
- [Contact](#)
  
- 
- 
- 
- 
- 
  
- [Log In](#)



- [About](#)
  - [Who We Are](#)
  - [Our History](#)
  - [Our Team](#)
  - [Sustainable Development Goals](#)
  - [Careers](#)
- [Membership](#)
  - [Overview](#)
  - [Our Members](#)
  - [Corporate Membership](#)
- [Resources](#)
- [Certification](#)
  - [Best Aquaculture Practices](#)
  - [Best Seafood Practices](#)

Search...



[Log In](#)

- [About](#)
  - [Who We Are](#)
  - [Our History](#)
  - [Our Team](#)
  - [Sustainable Development Goals](#)
  - [Careers](#)
- [Membership](#)
  - [Overview](#)
  - [Our Members](#)
  - [Corporate Membership](#)
- [Resources](#)
- [Certification](#)
  - [Best Aquaculture Practices](#)
  - [Best Seafood Practices](#)
- [GOAL Events](#)
- [Advocate Magazine](#)
- [Aquademia Podcast](#)
- [Blog](#)
- [Contact](#)



Health & Welfare  
Health & Welfare

Green water culture of California halibut larvae



1 April 2005 Jean-Benoit Muguet Douglas E. Bush Douglas E. Conklin, Ph.D. Raul H. Piedrahita, Ph.D. German E. Merino, Ph.D.



## Beneficial effects of microalgae during larviculture include fivefold increase in survival



Left: California halibut larvae 12 days after hatching. Right: Halibut near the end of metamorphosis at 30 days posthatch (7.5 mm). Bottom: California halibut juvenile at 45 days posthatch (12.8 mm).

The California halibut (*Paralichthys californicus*), an endemic species along the California coast of the United States, is an important sport fish with potential for culture in aquaculture systems.

Small-scale production of California halibut juveniles was originally realized in a clearwater flow-through system at the California Halibut Hatchery Program in Redondo Beach. The systems used conical-bottom tanks for the pelagic phase and shallow raceways for the benthic period thereafter.

A larger recirculation system was developed at the University of California in Davis, where larger numbers of juveniles have been produced. However, the larval growth and survival in both systems was too limited to be applied to large-scale production.

## Microalgae improves performance

Numerous studies have shown that the addition of microalgae to tanks during early rearing can improve larval performance for a number of species. The authors recently investigated the potential benefits of green water larviculture for California halibut using both live algae and algae paste. The study was funded by the National Sea Grant College Program of the U.S. Department of Commerce National Oceanic and Atmospheric Administration and the California State Resource Agency.

## California study

A batch of California halibut eggs was collected at the California Halibut Hatchery Program facility. The newly hatched larvae were transported to the University of California in Davis and divided into three groups.

The first group was raised in clear water in two 180-liter cylindrical tanks of 0.8 meters diameter and 1.1 meters maximum depth with conical bottoms connected to a recirculation system. The second group was reared in two static 120-liter cylindrical tanks of 0.5 meters diameter and 0.6 meters maximum depth. Additions of *Isochrysis* algae paste were made twice a day to maintain the algae concentration between 800,000 and 1 million cells per milligram.

The third group was cultured in 100-liter static raceways that measured 2.2 meters long, 0.35 meters wide, and 0.15 meters deep. Live *Isochrysis* were used to maintain the green water environment.

For all groups, water quality parameters were kept as follows: temperature –  $21.0 \pm 0.5$  degrees-C, dissolved oxygen –  $7.7 \pm 1.0$  milligrams per liter, salinity:  $30.5 \pm 1.0$  grams per liter, and photoperiod – 16 hours light and eight hours dark. Starting at mouth opening three days posthatch, the larvae were given rotifers fed baker's yeast and enriched with a rotifer grow-out diet. During the 15 to 40 days after hatching, the fish were fed enriched artemia nauplii and meta-nauplii.

Twenty larvae were sampled at regular intervals in each treatment for total length measurement. Survival was evaluated by counting at 28 and 38 days posthatch for the clear- and green water tanks, respectively.

## Positive results

Fish survival was significantly affected by the presence of the microalgae. Both treatments with microalgae had a better survival rate than the clear water treatment (Table 1). Larvae growth was also positively affected by the algae.

Muguet, Larval halibut growth and survival, Table 1

	Live Algae	Algae Paste	Clearwater
Total length at 17 days posthatch (mm)	5.90	5.80	5.60
Total length at 35 days posthatch (mm)	8.20	7.90	6.80
Survival at 38 days posthatch (%)	54.55	52.65	7.70

Table 1. Larval halibut growth and survival.

Increase in size was similar for all treatments during the premetamorphic period. However, 35 days posthatch, the larvae raised in green water were larger. Growth was similar for fish raised on algae paste or live algae.

Other parameters were also influenced by the use of algae. The number of unpigmented juveniles was reduced dramatically, and subsequent weaning onto artificial diets was more successful in the green water fish, probably due to their larger size.

Algae use recommended

The beneficial effects of microalgae during larviculture have been reported for numerous fish. The fivefold increase in survival reported in this research, for example, was consistent with similar experiments carried out with different marine species. Multiple factors are thought to improve larval quality: better localization of the live prey, stabilization or improvement of the water quality, direct or indirect nutritional effects, and increased pancreatic and intestinal enzymatic production.

The use of dead but intact microalgae in the algal paste did not affect the results relative to the use of live algae, even though no beneficial effects on water quality could be attributed to the algal paste. This contrasted with the live algae, which should contribute to improved water quality. Given the improved survival and weaning, green water culture of California halibut larvae using live algae or algae paste is strongly recommended.

(Editor’s Note: This article was originally published in the April 2005 print edition of the Global Aquaculture Advocate.)

Now that you've finished reading the article ...




... we hope you’ll consider supporting our mission to document the evolution of the global aquaculture industry and share our vast network of contributors’ expansive knowledge every week.

By becoming a Global Seafood Alliance member, you’re ensuring that all of the pre-competitive work we do through member benefits, resources and events can continue. Individual membership costs just \$50 a year.

Not a GSA member? Join us.

[Support GSA and Become a Member](#)

Authors


-  Jean-Benoit Muguet  
  
Jean-Benoit Muguet  
  
Department of Animal Science  
University of California  
One Shields Avenue  
Davis, California 95616 USA  
  
[117,100,101,46,115,105,118,97,100,99,117,64,116,101,117,103,117,109,97,106]
-  Douglas E. Bush  
  
Douglas E. Bush  
  
Department of Animal Science  
University of California  
One Shields Avenue  
Davis, California 95616 USA
-  Douglas E. Conklin, Ph.D.  
  
Douglas E. Conklin, Ph.D.  
  
Department of Animal Science  
University of California

One Shields Avenue  
Davis, California 95616 USA

-  Raul H. Piedrahita, Ph.D.

Raul H. Piedrahita, Ph.D.





Department of Biological and Agricultural Engineering  
University of California  
Davis, California, USA

-  German E. Merino, Ph.D.

German E. Merino, Ph.D.

Departamento de Acuicultura  
Universidad Catolica del Norte  
Coquimbo, Chile

## Share

-  [Share via Email](#)
-  [Share on Twitter](#)
-  [Share on Facebook](#)
-  [Share on LinkedIn](#)

## Tagged With

[halibut](#) [German E. Merino](#) [Raul H. Piedrahita](#) [Douglas E. Conklin](#) [Douglas E. Bush](#) [Jean-Benoit Muguet](#)

## Related Posts

Health & Welfare

### [Integrated larval-nursery system for California halibut](#)

Hubbs-SeaWorld Research Institute developed an integrated larval-nursery recirculating aquaculture system for California halibut utilizing shallow raceways for juvenile culture.

Aquafeeds

### [Amino acid requirements in developing marine fish](#)

Most marine fish have pelagic eggs with large pools of free amino acids of fairly constant proportion regardless of species. The amino acids provide energy for metabolism and are an important nutritional asset for growing embryos.

Intelligence

### [Halibut farming in Norway](#)

While stable production of juveniles remains an issue for halibut farming in Norway, advances such as diets to replace live artemia are improving the industry.

Innovation & Investment

### [Norway showcases halibut farm as sustainability star](#)

Recognizing its potential as a sustainable fish producer, Norway highlights land-based halibut farmer Sogn Aqua in the debut of The Explorer digital showroom. The Advocate recently paid a visit to the farm in Ortnevik.

## About The Advocate

The Responsible Seafood Advocate supports the Global Seafood Alliance's (GSA) mission to advance responsible seafood practices through education, advocacy and third-party assurances.

[Learn More](#)

Search Responsible Seafood Advocate



## Advertising Opportunities

[2022 Media & Events Kit](#)

## Categories

[Aquafeeds](#) > [Health & Welfare](#) [Health & Welfare](#) > [From Our Sponsors](#) > [Innovation & Investment](#) > [Intelligence](#) > [Responsibility](#) > [Fisheries](#) > [Artículos en Español](#) >

## Don't Miss an Article

## Featured

- [Health & Welfare](#) [An update on vibriosis, the major bacterial disease shrimp farmers face](#)
- [Uncategorized](#) [A seat at the table: Fed By Blue team says aquaculture needs a stronger voice](#)
- [Responsibility](#) [Quantifying habitat provisioning at macroalgae cultivation locations](#)

## Popular Tags

All Tags ▼

## Recent

- [Fisheries](#) [Second Test: Another filler for the fisheries category](#)
- [Fisheries](#) [Test: This is filler for the fisheries Category](#)
- [Aquafeeds](#) [Test Article](#)
- [Responsibility](#) [Study: Climate change will shuffle marine ecosystems in unexpected ways as ocean temperature warms](#)
- [Health & Welfare](#) [Indian shrimp researchers earn a patent for WSSV diagnostic tool](#)



- [About](#)
- [Membership](#)
- [Resources](#)
- [Best Aquaculture Practices \(BAP\)](#)
- [Best Seafood Practices \(BSP\)](#)
- [GOAL Events](#)
- [Advocate Magazine](#)
- [Aquademia Podcast](#)
- [Blog](#)
- [Contact](#)

#### Stay up to date with GSA

- 
- 
- 
- 
- 

Copyright © 2024 Global Seafood Alliance  
All rights reserved.

[Privacy](#)

[Terms of Use](#)

[Glossary](#)