



- [GOAL Events](#)
- [Advocate Magazine](#)
- [Aquademia Podcast](#)
- [Blog](#)
- [Contact](#)

-
-
-
-
-

- [Log In](#)



- [About](#)
 - [Who We Are](#)
 - [Our History](#)
 - [Our Team](#)
 - [Sustainable Development Goals](#)
 - [Careers](#)
- [Membership](#)
 - [Overview](#)
 - [Our Members](#)
 - [Corporate Membership](#)
- [Resources](#)
- [Certification](#)
 - [Best Aquaculture Practices](#)
 - [Best Seafood Practices](#)

Search...



[Log In](#)

- [About](#)
 - [Who We Are](#)
 - [Our History](#)
 - [Our Team](#)
 - [Sustainable Development Goals](#)
 - [Careers](#)
- [Membership](#)
 - [Overview](#)
 - [Our Members](#)
 - [Corporate Membership](#)
- [Resources](#)
- [Certification](#)
 - [Best Aquaculture Practices](#)
 - [Best Seafood Practices](#)
- [GOAL Events](#)
- [Advocate Magazine](#)
- [Aquademia Podcast](#)
- [Blog](#)
- [Contact](#)



Health & Welfare
Health & Welfare

Low-salinity culture water controls Vibrios in shrimp postlarvae



1 November 2014 Dr. Carlos A. Ching Ing. Juan Portal Ing. Alfredo Salinas



Halophytic bacteria growth inhibited when exposed to low-salinity water



Inland farms in southern Ecuador can produce up to 10 mt/ha of shrimp using underground water with 2- to 3-ppt salinity.

Low-salinity shrimp culture in southern Ecuador is done at inland farms using underground water that is pumped into 0.5- to 1.0-hectare (ha) ponds with liners and plastic greenhouse covers. Paddlewheel aeration is continuous during the whole production cycle, which can yield 7-10 metric tons (MT) per ha in 90 to 120 days.

The hatcheries that supply postlarvae to these farms acclimate the shrimp in water from 30 to 5 ppt salinity before transportation to the farms (Table 1). Once at the farms, postlarvae are further acclimated to water with 2 ppt salinity in pondside tanks for direct stocking or in nursery ponds before they are finally stocked in grow-out ponds.

Ching, Acclimation protocol, Table 1

Salinity Range	Acclimation Time
20-30 ppt	2 ppt reduction every 20 minutes
15-20 ppt	2 ppt reduction every 30 minutes
10-15 ppt	1 ppt reduction every 30 minutes
5-10 ppt	1 ppt reduction every hour

Table 1. Acclimation protocol used by hatcheries that provide shrimp postlarvae to low-salinity inland farms in Ecuador.

Monitoring, results

In research by the authors, the first set of bacteriological analyses consisted of three samples taken from two hatchery tanks containing water at 30 ppt salinity that provided postlarvae to each farm. Macerates of P.L.₆ were cultured in agar, and *Vibrio* counts in colony-forming units per gram (CFU/g) were recorded either as yellow (sucrose-positive) or green (sucrose-negative) colonies.

A second set of these analyses were performed when P.L.₁₂ postlarvae arrived at each farm in 5-ppt salinity water before they were stocked. The last set of analyses was done either in the nursery pond or in the grow-out pond during the first days of culture. Green colonies of *Vibrio* were later identified as *V. parahaemolyticus*.

Postlarvae from the hatchery tank with direct stocking averaged 442,400 yellow CFU/g and 29,933 green CFU/g (Fig. 1), while the postlarvae with a nursery phase had 390,000 yellow CFU/g and 20,933 green CFU/g (Fig. 2).




Fig. 1: Average concentrations of *Vibrio* species in shrimp postlarvae from the hatchery until the first days of culture in the farm with direct postlarvae stocking.




Fig. 2: Average concentrations of *Vibrio* species in shrimp postlarvae from the hatchery until the first days in a nursery pond.

On arrival at the farms, an average of three samples of P.L.12 taken randomly from three transportation bags with 5-ppt salinity water indicated that in the farm with direct stocking, yellow *Vibrio* counts went down to 1,236 CFU/g, and green *Vibrio* counts went down to less than 100 CFU/g. At the other farm, average counts of yellow colonies went down to 3,000 CFU/g, and green colonies fell to 102 CFU/g.

Early culture

Finally, the last monitoring of *Vibrio* species in postlarvae was carried out during the first days of culture, either in the nursery pond for the two-phase farm or in a growout pond for the direct-stocking farm. In both cases, green colonies of *V. parahaemolyticus* were eradicated from the postlarvae in five days in the nursery pond and in 10 days in the growout pond. The difference between results for these two farms may be attributed to the higher minimum oxygen levels in the nursery pond (5.0 mg/L) than in the growout pond (4.0 mg/L).

Monitoring of the farms continued during the growout period to a final salinity of 3 ppt at harvest. It was noticed that *V. parahaemolyticus* never appeared again in the samples taken from the hemolymph and hepatopancreas tissues of shrimp, or even from water samples.

Shrimp postlarvae become free from *Vibrio parahaemolyticus* in this low-salinity nursery pond in a few days after stocking.


Perspectives

Vibrios are known as halophytic bacteria, meaning they grow well in high-salinity aquatic environments, and their growth is inhibited when they are exposed to low-salinity water. However, when shrimp postlarvae are infected with pathogenic *Vibrio* bacteria at very high concentrations in the hatchery, disease can become uncontrollable.

There is a good chance that the infection levels determine the fate of these postlarvae at farms, as is the case with early mortality syndrome (EMS) caused by a pathogenic strain of *Vibrio parahaemolyticus*. Even if infected postlarvae are cultured in freshwater, mortalities may occur during the first days of culture.

(Editor's Note: This article was originally published in the November/December 2014 print edition of the Global Aquaculture Advocate.)


Authors

-  Dr. Carlos A. Ching

Dr. Carlos A. Ching

Aquaculture Manager
Nicovita – Vitapro S.A.
Av. Argentina 4793
Callao, Lima, Peru

[101,112,46,109,111,99,46,111,114,112,97,116,105,118,64,109,103,110,105,104,99,99]

-  Ing. Juan Portal

Ing. Juan Portal

Technical Assistance Manager
Nicovita – Vitapro S.A.

-  Ing. Alfredo Salinas

Ing. Alfredo Salinas

Production Manager
Orocam
Huaquillas, Ecuador

Share

- [✉ Share via Email](#)
- [🐦 Share on Twitter](#)
- [f Share on Facebook](#)
- [in Share on LinkedIn](#)

Tagged With

[salinity](#) [Vibrio parahaemolyticus postlarvae](#) [shrimp postlarvae](#) [Carlos A. Ching](#) [Juan Portal](#) [vibrios](#)

Related Posts

Health & Welfare

[Acclimating shrimp postlarvae before pond stocking](#)

Shrimp postlarvae acclimation before stocking into the various growout systems (ponds, raceways, tanks) is a critical – and often overlooked, sometimes taken for granted – step in the shrimp culture process. Various water quality parameters should be changed slowly so that the young shrimp have the time to gradually adapt to the new conditions.

Aquafeeds

[Biofloc consumption by Pacific white shrimp postlarvae](#)

The stable isotopes technique with $\delta^{13}\text{C}$ and $\delta^{15}\text{N}$ can be used to determine the relevance of different food sources to shrimp feeding during the pre-nursery phase of *Litopenaeus vannamei* culture. During this trial, different types of commercial feed, microalgae, *Artemia* sp. nauplii and bioflocs were used as food sources.

Intelligence

[As ocean temperatures rise, so too will vibrio outbreaks](#)

A study using a half-century of data has linked climate change and warming sea temperatures with an increase in illnesses from the common vibrio bacteria. Shellfish growers, fighting a particularly virulent strain of *Vibrio parahaemolyticus*, are changing their harvest protocols.

Health & Welfare

[A holistic management approach to EMS](#)

Early Mortality Syndrome has devastated farmed shrimp in Asia and Latin America. With better understanding of the pathogen and the development and improvement of novel strategies, shrimp farmers are now able to better manage the disease.

About The Advocate

The Responsible Seafood Advocate supports the Global Seafood Alliance's (GSA) mission to advance responsible seafood practices through education, advocacy and third-party assurances.

[Learn More](#)

Search Responsible Seafood Advocate



Advertising Opportunities

[2022 Media & Events Kit](#)

Categories

[Aquafeeds](#) > [Health & Welfare](#) [Health & Welfare](#) > [From Our Sponsors](#) > [Innovation & Investment](#) > [Intelligence](#) > [Responsibility](#) > [Fisheries](#) > [Artículos en Español](#) >

Don't Miss an Article

Featured

- [Health & Welfare](#) [An update on vibriosis, the major bacterial disease shrimp farmers face](#)
- [Intelligence](#) [A seat at the table: Fed By Blue team says aquaculture needs a stronger voice](#)
- [Responsibility](#) [Quantifying habitat provisioning at macroalgae cultivation locations](#)

Popular Tags

All Tags ▼






Recent

- [Fisheries](#) [Second Test: Another filler for the fisheries category](#)
- [Fisheries](#) [Test: This is filler for the fisheries Category](#)
- [Aquafeeds](#) [Test Article](#)
- [Responsibility](#) [Study: Climate change will shuffle marine ecosystems in unexpected ways as ocean temperature warms](#)
- [Health & Welfare](#) [Indian shrimp researchers earn a patent for WSSV diagnostic tool](#)



- [About](#)
- [Membership](#)
- [Resources](#)
- [Best Aquaculture Practices \(BAP\)](#)
- [Best Seafood Practices \(BSP\)](#)
- [GOAL Events](#)
- [Advocate Magazine](#)
- [Aquademia Podcast](#)
- [Blog](#)
- [Contact](#)

Stay up to date with GSA

- 
- 
- 
- 
- 

Copyright © 2024 Global Seafood Alliance
All rights reserved.

[Privacy](#)

[Terms of Use](#)

[Glossary](#)