



(<https://gsa.rakadev.com>).




Health &
Welfare

Metabolomics approaches to improve mussel larval production

30 November 2015

By Andrea C. Alfaro, Ph.D. and Tim Young, Ph.D.

Analyses of metabolites and their ratios can be integrated with gene and protein expression data



Poor-quality mussel larvae (left) grow more slowly than higher-quality – and faster-growing – larvae (right). Scale bars = 20 μ m. Photos by A. Rusk.

Hatchery production of mussel larvae presents a number of biological and technological challenges that continue to hamper growth within the industry. Unexpected batch crashes are routinely observed in some hatcheries, and the causes are often not identified. When solutions are found and improvements made, these tend to be incremental and directed by technical managers.

Unfortunately, technical developments are poorly captured in the primary scientific literature and often not shared due to concerns over losing commercial advantage. In general, low production yields are hampered by high larval mortality rates that stem from the poor quality of broodstock, inadequate feeding regimes or unhealthy culture systems.

With the complex physical and biological factors within the larval-rearing process, it is not surprising larval cultivation still remains a hit-and-miss approach with variable chances of success. In this regard, metabolomics can provide a powerful means of identifying problems, and potentially could be used as an early warning system in subsequent cultures.

Metabolomics

Metabolomics is the study of chemical processes involving metabolites. As Wikipedia said, metabolomics is the “systematic study of the unique chemical fingerprints that specific cellular processes leave behind.” Metabolic profiling can give an instantaneous snapshot of the physiology of cells.

Because this is a new application in larval biology, there are only a few studies to illustrate its advantages for aquaculture. The authors recently performed a study using metabolomics to investigate intraspecific growth variations in New Zealand Greenshell™ *Perna canaliculus* mussel larvae.

Mussel aquaculture in New Zealand

In New Zealand, exports of *P. canaliculus* mussels represent the largest aquaculture sector by value and volume. Small-scale hatchery production of spat currently contributes only marginally toward the industry's seed requirements. However, substantial research over the past decade has led to the ongoing development of selective breeding lines, establishment of a cryopreservation program, optimization of microalgae culture and larval-rearing procedures, and future strategies for extensive growth and upscaling of hatchery facilities.

The ability to provide consistency in larval quality and quantity is a critical step to achieve the commercial goal of successful large-scale production. Identification of biomarkers that reflect the immediate physiological condition of larvae has the potential to provide valuable tools that not only determine larval batch viability early on, but also identify the causes of poor production levels. This information can lead to better management decisions and improve overall hatchery operations.




Figure 1. PLS-DA scores plot (B) of poor (circles) and high (squares) quality larvae. Hierarchical clustering and heat map (A) of poor (■) and high (■) quality larvae based on the top 50 features ranked by their T-test statistic (samples on the y-axis, features on the x-axis). PLS-DA scores plot (B) of poor-

(●) and high- (●) quality larvae. SAM plot (C) of significantly different features (○) between quality classes (upper right = ratios that were higher in poor-quality group, lower left = ratios that were lower). EBAM plot (D) of significantly different features (○) between quality classes (all lower in the poor-quality group). Volcano plot (E) of features (●) with a between-class fold-change greater than 5 and a T-test statistic below 0.01 (upper left = ratios that were higher in the poor-quality group; upper right = ratios that were lower). Venn diagram (E) displays the counts of commonly identified ratios by the four independent feature selection methods.

Study setup

In this study, the physiological condition of mussel larvae was used to assess larval quality during hatchery production. Mussel larvae were grown to umbo stage, separated based on relative sizes to divide poor-quality, slow-growing larvae from high-quality, fast-growing animals, and sampled over time. Six samples including about 80,000 pooled individuals from each of the two size fractions and three replicate rearing tanks were used for metabolite extractions via gas chromatography/mass spectrometry.

Metabolite peaks were identified, and their relative abundances were calculated. The dataset matrix was subjected to a variety of feature selection methods to identify potential biomarkers for discrimination of larval quality classes. Comprehensive Web-based analytical pipeline tools were used to statistically analyze the data sets.

To visualize feature differences between the larval quality classes, metabolite ratios were used to perform agglomerative hierarchical cluster analysis, displayed in dendrograms and heat maps. Four methods of feature reduction were used independently to minimize selection bias and provide robust criteria for assisting candidate biomarker identification.

Results

From the initial 253 metabolite ratios analyzed, candidate biomarkers were identified for assessing mussel larval quality using four independent feature selection methods: Volcano plot, partial least squares discriminant analysis (PLS-DA), significant analysis of microarrays (SAM) and empirical Bayesian analysis of microarrays (EBAM). These methods resulted in the selection of 19 common features.

Based on their performance, these were reduced to a final four metabolite ratios that were good candidates for assessing larval quality: alanine/succinate, glycine/succinate, myristic acid/succinate and pyroglutamate/succinate. Each of the identified ratios was substantially lower in the poor-quality larvae (Figure 1).

Comparing this group with the faster-growing cohort, the trend was characterized by elevated levels of succinate and simultaneous reductions in relative metabolite abundances of alanine, glycine, pyroglutamate and myristic acid. Based on the known functions of these metabolites, it was possible to identify potential biochemical pathways involved in larval performance. These pathways include energy metabolism, osmotic regulation, immune function and cell-to-cell communication.

Perspectives

While this study identified broad areas of larval physiology responsible for larval quality, the authors' ongoing studies are providing more specific biomarkers intended to generate predictive models of larval performance. Furthermore, the aim is to eventually develop easy-to-use tool kits to evaluate the physiological state of larvae throughout the rearing process.

Further analysis of the single metabolites and their ratios will likely reveal additional information, and when integrated with gene and protein expression data, could provide new avenues for selective-breeding programs to consistently yield high-quality larvae. Supplementary experiments incorporating metabolomics-based approaches to investigate other measures of larval quality – such as health after immunological challenge and nutritional condition in response to diet manipulation – have the potential to offer a suite of biomarkers for a range of applications.

The use of metabolomics could also be applied to other areas of hatchery production to help close the loop on full life-cycle culture for many species. For example, routine production of high-quality gametes for successful fertilization and on-growing could be achieved through better broodstock management and understanding of maternal provisioning, paternal effects and factors associated with high fecundity.

Authors



ANDREA C. ALFARO, PH.D.

Institute for Applied Ecology New Zealand
School of Applied Sciences
Auckland University of Technology
Auckland, New Zealand

aalfaro@aut.ac.nz (<mailto:aalfaro@aut.ac.nz>)



TIM YOUNG, PH.D.

Institute for Applied Ecology New Zealand
School of Applied Sciences
Auckland University of Technology

All rights reserved.