### Global Seafood Alliance Logo

- GOAL Events
- Advocate Magazine
- Aquademia Podcast
- Blog
- Contact
- (O)

- <u>Log In</u>

- About
  - Who We Are
  - Our History
  - o Our Team
  - Sustainable Development Goals
  - Careers
- <u>Membership</u>
  - Overview
  - Our Members
  - Corporate Membership
- Resources
- **Certification** 
  - Best Aquaculture Practices
  - Best Seafood Practices

Search...



- About
  - Who We Are
  - Our History
  - o Our Team
  - Sustainable Development Goals
  - Careers
- <u>Membership</u>
  - o <u>Overview</u>
  - Our Members
  - o Corporate Membership
- Resources
- **Certification** 
  - Best Aquaculture Practices
  - Best Seafood Practices
- GOAL Events
- Advocate Magazine

- <u>Aquademia Podcast</u>
- Blog
- Contact



# Microencapsulated organic acids aid shrimp culture

Responsible Seafood Advocate logo

25 January 2016 Nicholas Romano, Ph.D. Chik-Boon Koh, Ph.D., Wing-Keong Ng, Ph.D.



# Novel blends enhance productivity and Vibrio resistance



The farming of Pacific white shrimp is a major industry in Asia that has frequently been challenged with disease outbreaks, including the recent Vibrio infections.

Disease outbreaks in the shrimp farming industry have caused substantial economic losses around the world and, in particular, the recent "early mortality syndrome" (EMS) due to plasmids in Vibrio bacteria is a growing threat. In addition to decreasing immunity, EMS causes substantial hepatopancreatic damage to shrimp, which gradually induces mass mortalities. Moreover, with a greater restriction or altogether ban on antibiotic use, this can leave shrimp farmers with fewer options to protect their animals against bacterial diseases.

# Organic acids in shrimp feeds

In response, researchers have investigated environmentally-friendly dietary alternatives as potential prophylactics. One potential alternative are organic acids, which are "Generally Regarded as Safe," or GRAS, and that have been used for decades with success as growth promoters and antimicrobials in the terrestrial livestock feed industry. However, not much is known about their efficacy to shrimp farming productivity.

Recently, our laboratory research group conducted a controlled study where Pacific white shrimp (*Litopenaeus vannamei*) were fed with different dietary levels of a prototype microencapsulated organic acids blend (OAB) at 0 (control), 1, 2 or 4 percent and evaluated their growth performance, phenoloxidase (PO) activity and hepatopancreatic histopathology after 50 days. Dry matter and phosphorus utilization efficiency were also determined. At the end of the feeding trial, shrimp were challenged with pathogenic *Vibrio harveyi* and their survival monitored along with the associated changes to immunity and hepatopancreatic histopathology.

# **Novel microencapsulated organic acids**

Various organic acids were pre-screened, alone and in combination, for their antagonistic and synergistic activities to *Vibrio* sp. in order to formulate an optimal blend. This was followed by a two-month pilot study on white shrimp to narrow down the most appropriate dietary OAB levels. The OAB was then microencapsulated to minimize leaching, which is particularly important for crustaceans that are relatively slow feeders and tear apart their diets while feeding.

The propriety OAB (Orgacids<sup>TM</sup>-AQUA) was developed and produced in collaboration with Sunzen Feedtech Pte. Ltd. (Malaysia) in which four organic acids (formic, lactic, malic and citric acids) were coated in a specialized lipid matrix using high-speed, centrifugal spray cooling technology that produced micro-capsules of less than 250 microns. The final product was a free- flowing fine powder, non-corrosive and resistant to water leaching of the soluble organic acids. More importantly, the encapsulation process should allow a slower release of the organic acids throughout the entire length of the shrimp gut for optimal efficacy.

The OAB was added to the shrimp practical diets at 0 (control), 1, 2 or 4 percent. Soybean meal and squid liver oil were the main protein and lipid sources, respectively.



A scanning electron micrograph of the microencapsulated organic acids blend manufactured using centrifugal cool-spraying technology by Sunzen Feedtech Pte. Ltd. (Malaysia).

# Shrimp feeding trial

Triplicate groups of shrimp were hand-fed their respective experimental diet three times a day to apparent satiation in a closed recirculating system for 50 days and then we measured their growth, PO activity and hepatopancreatic histopathology. In a separate experiment, shrimp were measured for dry matter and phosphorus (P) digestibility and then challenged with *V. harveyi* for 10 days, followed by measuring their subsequent survival, PO activity and hepatopancreatic histopathology.

Growth, survival and P utilization were improved in all OAB treatments, with 2 percent OAB giving the best results (growth and P utilization were significantly higher than the control), while the PO activity was unaffected. The resistance of shrimp to pathogenic *V. harveyi* when fed the OAB diets was significantly enhanced (Figure 1) as well as the PO activity, while the hepatopancreas showed much less histopathological damage.




Figure 1: The cumulative mortality (%) of white shrimp after being fed different levels of OAB and challenged with *Vibrio harveyi*.

Figure 1: The cumulative mortality (%) of white shrimp after being fed different levels of OAB and challenged with Vibrio harveyi.

# Dietary organic acids efficacy in shrimp

The effectiveness of the tested dietary OAB as a growth promoter was likely due to improved nutrient utilization, which was also supported by observing substantially more lipid storage cells in the hepatopancreas. In particular, P utilization was significantly enhanced, which may have implications to reducing excessive and harmful P discharge to the surrounding aquatic environment. Meanwhile, although feed intake was not directly measured, it was routinely observed that shrimp fed diets supplemented with organic acids were more active upon feed introduction. Some organic acids, such as propionate and butyrate, have recently been reported to act as feed attractants to white shrimp, although these were not in the tested OAB formulation.

The present study demonstrated the immune boosting and hepatopancreatic protective properties of organic acids. Shrimp fed the OAB treated diets had significantly higher PO activity and less hepatopancreatic damage, which likely explains their higher survival during *V. harveyi* challenge. Vibriosis induces shrimp mortalities by decreasing immunity and causing hepatopancretic damage.

# **Perspectives**

The use of the prototype OAB evaluated in this study or other organic acids or their salts in combination or singly, may be a particularly beneficial functional feed additive for the marine shrimp farming industry. Further feeding trials including farm-based trials are currently being conducted to further confirm the beneficial impacts of organic acids in shrimp feeds.

Authors' note: For the full report of this study, please refer to, "Romano, N., Koh, C.B., Ng, W.K. 2015. Dietary microencapsulated organic acid blends enhances growth, phosphorus utilization, immune response, hepatopancreatic integrity and resistance against Vibrio harveyi in white shrimp, Litopenaeus vannamei. Aguaculture 435, 228-236."

#### Authors

• Nicholas Romano, Ph.D.

Nicholas Romano, Ph.D.

Fish Nutrition Laboratory School of Biological Sciences Universiti Sains Malaysia Penang 11800, Malaysia

[121,109,46,109,115,117,64,103,110,107,119]

Chik-Boon Koh, Ph.D., Wing-Keong Ng, Ph.D.

Chik-Boon Koh, Ph.D., Wing-Keong Ng, Ph.D.

Fish Nutrition Laboratory School of Biological Sciences Universiti Sains Malaysia Penang 11800, Malaysia

#### Share

- Share via Email
- **Share on Twitter**
- **f** Share on Facebook
- in Share on LinkedIn

#### **Tagged With**

Shrimp Vibrio microencapsulated organic acids shrimp feeding Romano organic Nicholas Romano

#### **Related Posts**

Health & Welfare

### **Vibrio control in shrimp farming: Part 1**

Control of Vibrio bacteria should focus on minimizing overall bacterial loads and the potential for horizontal transmission. The challenge for hatchery managers is identifying gaps in biosecurity and plugging them without creating niches for other potential pathogens.

Health & Welfare

## Vibrio control in shrimp farming: Part 2

Strategies to control Vibrio bacteria at aquaculture farms must revolve around limiting inputs, including those from postlarvae suppliers, those that come in with the water, those that remain in the soil between crops and those in the feed and accumulated organic matter.

Health & Welfare

### Genetic variation for resistance to WSS, AHPND in Pacific white shrimp

Selection for disease resistance has been used in breeding farm animals and can be a viable option to deal with white spot syndrome and acute hepatopancreatic necrosis disease in commercial shrimp culture. In trials, heritability for AHPND resistance was low, while that for WSS was moderate.

Health & Welfare

# Limited decomposition enhances PCR detection of AHPND Vibrio in shrimp

A study confirmed the utility for improved polymerase chain reaction (PCR) detection of the Vibrio bacteria that cause acute hepatopancreatic necrosis disease (AHPND) in asymptomatic shrimp by permitting the shrimp to expire and decompose for several hours prior to preservation and PCR processing.

Health & Welfare

# Emerging epitheliocystis disease in Mediterranean sparids caused by novel bacteria


Epitheliocystis is an emerging infection among farmed gilthead seabream and is also lethal in mesocosm cultures of sharpsnout seabream larvae. Two studies of this disease at sites in Greece and Crete have characterized the gill and skin cysts in more detail.

### **About The Advocate**

The Responsible Seafood Advocate supports the Global Seafood Alliance's (GSA) mission to advance responsible seafood practices through education, advocacy and third-party assurances.

#### **Learn More**

Search Responsible Seafood Advocate Search Search



### **Advertising Opportunities**

2022 Media & Events Kit

### Categories

Aquafeeds Aquafeeds > Health & Welfare > From Our Sponsors > Innovation & Investment > Intelligence > Responsibility > Fisheries > Artículos en Español >

### Don't Miss an Article

#### **Featured**

- Health & Welfare An update on vibriosis, the major bacterial disease shrimp farmers face
- Uncategorized A seat at the table: Fed By Blue team says aquaculture needs a stronger voice
- Responsibility Quantifying habitat provisioning at macroalgae cultivation locations

### **Popular Tags**

All Tags 

✓

#### Recent

- Fisheries Second Test: Another filler for the fisheries category
- Fisheries Test: This is filler for the fisheries Category
- Aguafeeds Test Article
- Responsibility Study: Climate change will shuffle marine ecosystems in unexpected ways as ocean temperature warms
- Health & Welfare Indian shrimp researchers earn a patent for WSSV diagnostic tool



- About
- Membership
- Resources
- Best Aquaculture Practices (BAP)
- Best Seafood Practices (BSP)
- GOAL Events
- Advocate Magazine
- Aquademia Podcast
- Blog
- Contact

#### Stay up to date with GSA

Copyright © 2024 Global Seafood Alliance All rights reserved.

<u>Privacy</u> Terms of Use Glossary