



- [GOAL Events](#)
- [Advocate Magazine](#)
- [Aquademia Podcast](#)
- [Blog](#)
- [Contact](#)
-
-
-
-
-
- [Log In](#)



- [About](#)
 - [Who We Are](#)
 - [Our History](#)
 - [Our Team](#)
 - [Sustainable Development Goals](#)
 - [Careers](#)
- [Membership](#)
 - [Overview](#)
 - [Our Members](#)
 - [Corporate Membership](#)
- [Resources](#)
- [Certification](#)
 - [Best Aquaculture Practices](#)
 - [Best Seafood Practices](#)

Search...



[Log In](#)

- [About](#)
 - [Who We Are](#)
 - [Our History](#)
 - [Our Team](#)
 - [Sustainable Development Goals](#)
 - [Careers](#)
- [Membership](#)
 - [Overview](#)
 - [Our Members](#)
 - [Corporate Membership](#)
- [Resources](#)
- [Certification](#)
 - [Best Aquaculture Practices](#)
 - [Best Seafood Practices](#)
- [GOAL Events](#)
- [Advocate Magazine](#)
- [Aquademia Podcast](#)
- [Blog](#)
- [Contact](#)



Mixed maturation diets improve shrimp broodstock performance



1 June 2001 P. Coutteau E. Dencece D. Pham E. Pinon Y. Balcazar



Mixture of fresh feeds, dry broodstock pellet results in lower broodstock mortality for male and female *P. vannamei*



P. stylirostris broodstock (without striped coloration) fed a 50/50 mix of fresh and pelleted feed provided quality offspring through repeated spawnings.

Interest in captive reproduction of penaeid shrimp is increasing worldwide due to the urgent need to establish selective breeding programs and produce certified, diseasefree postlarvae. An optimal diet is a crucial factor in the successful sexual maturation and reproduction of shrimp in breeding operations. High-quality food represents the highest operational cost in most broodstock management facilities. Maturation facilities in marine shrimp hatcheries depend mostly on fresh feeds, which can be inconvenient for operators. Dry artificial feeds, on the other hand, have several advantages, including reliable supply, consistent and controlled quality, and ease of handling. Artificial diets also reduce fouling of larval tanks, cut risk for disease transfer, and offer effective delivery of chemotherapeutics, immunostimulants and hormones. Trials in Ecuador and New Caledonia indicated that the incorporation of artificial feeds in broodstock diets can improve broodstock performance in several areas.

A dry, off-the-shelf broodstock feed was formulated to provide optimal palatability, stability, and nutrition for broodstock shrimp (Table 1). The diet contains no marine protein sources derived from aquaculture, and provides a constant supply of all known essential nutrients for broodstock maturation. These include highly unsaturated fatty acids of the omega-3 and -6 families, phospholipids, cholesterol, carotenoids, vitamins, minerals, artemia and squid factors. The combination of appropriate extrusion and binder technology resulted in high water stability.

Coutteau, Nutritional profile of INVE Breed-Shrimp, Table 1

Component	Unit	Specifications
Moisture	%	< 10
Crude fiber	%	< 4
Crude protein	%	> 50
Crude fat (hydrolysis)	%	> 10
Essential Fatty Acids		
ARA (20:4n-6)	mg/g DW	> 1.5
EPA (20:5n-3)	mg/g DW	> 5
DHA (22:6n-3)	mg/g DW	> 10
n-3 HUFA (≥ 20:3n-3)	mg/g DW	> 20
n-6 PUFA (≥ 18:2n-6)	mg/g DW	> 10
Vitamin C	mg/kg	> 3,000
Vitamin A	IU/kg	> 20,000
Vitamin D3	IU/kg	> 8,000
Vitamin E	mg/kg	> 800

Table 1. Nutritional profile of INVE Breed-Shrimp broodstock feed.




Fig. 1: Average daily mortality during a 39-day maturation trial with *Penaeus vannamei* broodstock fed fresh feeds or a mixture of fresh and dry feeds. During the last two weeks (Period II), both groups received 80 percent fresh and 20 percent dry feed.

Maturation test with *P. vannamei*

The feed was evaluated under commercial conditions as a partial substitute for fresh feeds during a 39-day maturation trial with *P. vannamei* at the Quirola Group's Quimasaru hatchery in Ecuador. The experiment used 16 maturation tanks (eight per treatment), each stocked with approximately 80 wild-caught shrimp at a sex ratio 1:1. Shrimp of 40 grams average body weight were acclimated and ablated prior to stocking. Natural photoperiod was used, with a 50 percent daily water exchange and temperature of 29 to 30 degrees-C.




Fig. 2: Mating frequency during a 39-day maturation trial with *Penaeus vannamei* broodstock fed fresh feeds or a mixture of fresh and dry feeds. During the last two weeks, both groups received 80% fresh and 20% dry feed. Observations were discontinued during periods between production runs.

During Period I, the first 19 days of the experiment, shrimp in the control tanks were fed solely on a mixture of fresh feeds that included enriched artemia biomass, mussels, squid, bloodworms, and shrimp heads. Contrary to current practice, the shrimp heads and other potentially risky feed elements were included because there was no white spot syndrome virus alert at the time of the trial.

The other group was fed a reduced ration of fresh feeds in combination with the dry pellets. Overall, fresh feed replacement approximated 50 percent on a dry-matter basis, but was different for each type of fresh feed fed to the control tanks.

During the experiment, the hatchery operator wanted to benefit from the reduced mortality due to the supplementation of the broodstock feed. Therefore, the two groups were switched to the same feeding regime in the final period. Both groups received a mixture of about 80 percent fresh feeds and 20 percent dry pellets during Period II, the last two weeks of the experiment.

Mortality

High mortality after ablation of wild broodstock, with higher mortality observed in females, was an important problem during production runs. During Period I, animals fed the mixed diet had lower mortalities than animals fed just the fresh diet (Fig. 1). During Period II, overall mortality rate was reduced, and the effect of the dry feed supplement during the previous period remained important.

Mating



Fig. 3: Average fecundity from group spawnings during a 39-day maturation trial with *Penaeus vannamei* broodstock fed fresh feeds or a mixture of fresh and dry feeds.

Mating frequency (daily percentage of females fertilized, through artificial insemination or natural mating) was considerably higher during Period I for broodstock fed the fresh feed (Fig. 2). The proportion of natural insemination was 40 to 55 percent. However, during Period II the percentage of successfully fertilized females dropped in the shrimp fed fresh feeds. Overall, a more consistent mating throughout the production run was obtained in the group that received the supplemented broodstock feed.

During Period I, average fecundity of 245,000 eggs and 135,000 nauplii/female for the broodstock fed the mixed diet compared to 260,000 eggs and 152,000 nauplii/female for animals fed solely fresh feed (Fig. 3). This reflected 9 percent lower egg production and 14 percent fewer nauplii (no significant differences) in the group fed the mixed diet.

Fertilization percentages, which averaged 58 to 60 percent, were similar in both treatments. During Period II, average fecundity remained at an intermediate level compared to Period I (243,000 eggs and 144,000 nauplii/female).

Maturation test with *p. stylirostris*

Three trials were run with *P. stylirostris* at La Station d’Aquaculture de Saint-Vincent, an IFREMER facility in New Caledonia. Females were held at three animals per square meter in round, 4-meter-diameter tanks under typical maturation conditions of 29 to 30 degrees-C, with 50 percent water exchange/day and reversed photoperiod. Eyestalk ablation was performed.

The control feeding regime consisted of a mixture of fresh feeds (squid, mussels, and shrimp) and a small ration of dry commercial pellets. In the experimental regime, 50 percent of the fresh feed was replaced by the dry broodstock feed INVE Breed-Shrimp (Table 2).

Coutteau, Dietary regimes tested for maturation, Table 2

Treatment Squid Mussels Shrimp Dry Pellet Broodstock Feed

Fresh feed	7.3%	6.1%	2.5%	0.5%	0%
50/50 diet	3.7%	3.1%	1.2%	0.5%	2%

Table 2. Dietary regimes tested for maturation of *P. stylirostris* (% shrimp body weight).

No major difference in overall performance between the two dietary regimes was noticed over 18 days during the trials (Table 3). However, some interesting effects were observed in successive spawns. Fertilization rate and number of nauplii per spawn, which did not show global differences between

treatments in any of the trials, decreased in trial 1 with successive spawns in the fresh feed regime, whereas these parameters improved with successive spawns in the mixed dietary regime (Fig. 4).

Coutteau, Average female performance and nauplii quality, Table 3

Parameter	Fresh Feed 50/50 Mix	
Female Performance		
Female survival %	60 ± 33	59 ± 18
Maturations/month/female	3.2 ± 1.1	3.4 ± 1.4
Eggs/female/spawn (x 1,000)	251 ± 57	241 ± 50
Fertilization %	55 ± 11	53 ± 14
Hatching %	74 ± 16	73 ± 19
Nauplii/female/spawn (x 1,000)	112 ± 19	99 ± 20
Nauplii Quality		
Deformity index (0-5)	4.7 ± 0.2	4.8 ± 0.1
Activity index (0-5)	4.7 ± 0.1	4.8 ± 0.1
Survival to zoea I in culture (%)	65 ± 8	75 ± 11

Table 3. Average female performance and nauplii quality for *P. stylirostris* broodstock fed fresh feeds or a 50% ration of fresh feeds combined with a broodstock pellet.

Also, although no significant differences could be detected in overall survival rate from nauplii to zoea I, survival decreased drastically in the last week of the maturation trial in the fresh feed treatment. This was not the case for the 50/50 mixed regime (Fig 5). These results showed that the nutritional boost provided by the broodstock pellet supported more sustainable reproductive activity and goodquality offspring through repetitive spawnings.

Conclusion

A maturation diet consisting of a mixture of fresh feeds supplemented with a dry broodstock pellet produced lower broodstock mortality for male and female *P. vannamei* following ablation and more consistent mating throughout the production cycle than a diet of fresh feeds. Comparable fecundity and fertilization rates were recorded for both feed types in trials.

The replacement of 50 percent of fresh feed with dry feed for *P. stylirostris* broodstock resulted in similar female reproductive response rates and similar larval quality (deformities, activity, and survival) when compared to animals fed the control diet. The mixed diet also improved fecundity and larval quality over repeated spawnings.

(Editor’s Note: This article was originally published in the June 2001 print edition of the Global Aquaculture Advocate.)

Now that you've finished reading the article ...

... we hope you’ll consider supporting our mission to document the evolution of the global aquaculture industry and share our vast network of contributors’ expansive knowledge every week.

By becoming a Global Seafood Alliance member, you’re ensuring that all of the pre-competitive work we do through member benefits, resources and events can continue. Individual membership costs just \$50 a year.

Not a GSA member? Join us.

[Support GSA and Become a Member](#)

Authors

-  P. Coutteau

P. Coutteau

INVE Technologies N.V.
Oeverstraat, Belgium
-  E. Dencece

E. Dencece

IFREMER – Station d’Aquaculture de Saint-Vincent
Boulouparis, New Caledonia
-  D. Pham

D. Pham

IFREMER – Station d’Aquaculture de Saint-Vincent
Boulouparis, New Caledonia

-  E. Pinon

E. Pinon

Guayaquil, Ecuador

-  Y. Balcazar

Y. Balcazar

Grupo Quirola
Division Laboratories – Guayaquil, Ecuador

Share

-  [Share via Email](#)
-  [Share on Twitter](#)
-  [Share on Facebook](#)
-  [Share on LinkedIn](#)

Tagged With

[P. Coutteau](#) [Shrimp broodstock](#) [Maturation diets](#) [E. Dencece](#) [D. Pham](#) [E. Pinon](#) [Y. Balcazar](#)

Related Posts

Health & Welfare

[Advances in fish hatchery management](#)

Advances in fish hatchery management – particularly in the areas of brood management and induced spawning – have helped establish aquaculture for multiple species.

Health & Welfare

[10 paths to low productivity and profitability with tilapia in sub-Saharan Africa](#)

Tilapia culture in sub-Saharan Africa suffers from low productivity and profitability. A comprehensive management approach is needed to address the root causes.

Aquafeeds

[Fishmeal, fish oil replacements in sea bream, sea bass diets need nutritional compensation](#)

Fishmeal and fish oil replacements present a challenge for the aquafeed industry. The range of alternative protein and fat sources is limited, given multiple considerations.

Health & Welfare

[Alfalfa concentrate: natural shrimp color enhancer](#)

Adding alfalfa concentrate containing natural carotenoids and pigments to the finishing diets of shrimp can enhance coloration of shrimp after cooking.

About The Advocate

The Responsible Seafood Advocate supports the Global Seafood Alliance’s (GSA) mission to advance responsible seafood practices through education, advocacy and third-party assurances.

[Learn More](#)

Search Responsible Seafood Advocate



Advertising Opportunities

[2022 Media & Events Kit](#)

Categories

[Aquafeeds](#) [Aquafeeds](#) > [Health & Welfare](#) > [From Our Sponsors](#) > [Innovation & Investment](#) > [Intelligence](#) > [Responsibility](#) > [Fisheries](#) > [Artículos en Español](#) >

Don't Miss an Article

Featured

- [Health & Welfare](#) [An update on vibriosis, the major bacterial disease shrimp farmers face](#)
- [Uncategorized](#) [A seat at the table: Fed By Blue team says aquaculture needs a stronger voice](#)
- [Responsibility](#) [Quantifying habitat provisioning at macroalgae cultivation locations](#)

Popular Tags

All Tags ▼

Recent

- [Fisheries](#) [Second Test: Another filler for the fisheries category](#)
- [Fisheries](#) [Test: This is filler for the fisheries Category](#)
- [Aquafeeds](#) [Test Article](#)
- [Responsibility](#) [Study: Climate change will shuffle marine ecosystems in unexpected ways as ocean temperature warms](#)
- [Health & Welfare](#) [Indian shrimp researchers earn a patent for WSSV diagnostic tool](#)



- [About](#)
- [Membership](#)
- [Resources](#)
- [Best Aquaculture Practices \(BAP\)](#)
- [Best Seafood Practices \(BSP\)](#)
- [GOAL Events](#)
- [Advocate Magazine](#)
- [Aquademia Podcast](#)
- [Blog](#)
- [Contact](#)

Stay up to date with GSA

- 
- 
- 
- 
- 

Copyright © 2024 Global Seafood Alliance
All rights reserved.

[Privacy](#)

[Terms of Use](#)

[Glossary](#)