



ALLIANCE™

(<https://gsa.rakadev.com>).



**Responsible
Seafood**
ADVOCATE



Health &
Welfare

Multiple characteristics considered in selection of probiotics for marine shrimp

1 November 2011

By Dr. Felipe do Nascimento Vieira , Dr. José Luis Mouriño , Bruno Correa da Silva, M.S. , Dr. Walter Quadros Seiffert and Dr. Luis Alejandro Vinatea

Shrimp fed diet supplemented with probiotics had higher survival, lower FCR



The use of *L. plantarum* was evaluated under intensive commercial pond conditions at the Yakult Experimental Farm-USFC. After 75 days, shrimp whose diet was supplemented with probiotics had higher survival and lower feed-conversion than control shrimp, with no alteration in growth rate.

Most people on this vast planet do not realize that microorganisms represent the majority of the Earth's inhabitants in number, biomass and genetic diversity. Among this immense world of microorganisms, the development of a specific probiotic for use in aquaculture is an arduous task that includes the isolation of microorganisms, selection according to in vitro characteristics and trials on pilot and commercial scales.

With financial support from the Brazilian Ministry of Fishing and Aquaculture (MPA), Financing Agency for Studies and Projects (FINEP), and Scientific and Technological Research Support Foundation of Santa Catarina (FAPESC), the authors conducted studies at the Universidade Federal de Santa Catarina in Brazil to develop a probiotic for marine shrimp cultivation.

Isolation of bacteria strains

The isolation of bacteria from the intestinal tract of the animal in study is the first step toward successful development of a probiotic. Selective and differentiating culture mediums are used to isolate specific groups of beneficial bacteria. In this case, the authors sought to isolate lactic acid bacteria because they have beneficial characteristics such as easy multiplication, the production of compounds that inhibit pathogenic bacteria (hydrogen peroxide, organic acids and bacteriocins), and production of exogenous enzymes and immunostimulant compounds.

To isolate the strains, a de Man, Rogosa and Sharpe medium with aniline blue indicator was used, so the lactic acid bacteria colonies were colored blue. In this step, 78 strains of lactic acid bacteria were isolated from the intestinal tracts of *Litopenaeus vannamei* shrimp cultivated in a super-intensive biofloc system.

In vitro strain selection

The high number of the bacterial strains isolated made direct evaluation of the effects of each strain on the animals in cultivation difficult. Various in vitro laboratory selection tests were thus conducted to select a few strains to test with the animals. For the selection used in this study, the following characteristics were considered.

Inhibition of pathogens.

Evaluation of the capacity of the isolated strains to inhibit the growth of pathogenic bacteria for marine shrimp by means of the diameter of the inhibition zone (Kirby-Bauer technique in diffusion in agar).

Growth kinetics. Growth velocity is an important characteristic, as high growth speed permits more efficient

commercial production processes and can indicate greater competitiveness of the strain in the environment.

Resistance to different salinities. In marine shrimp cultivation, variations in salinity depend on the region, time of year and rainfall, and can range from close to zero to higher than that of the ocean in very saline regions. It is therefore important for commercial probiotic bacteria to survive wide variations in salinity.

Resistance to different pH values. Since the pH of water used for cultivation is not constant and can vary from close to 6 in super-intensive shrimp cultivation with biofloc up to 9 in eutrophic ponds, probiotic strains must handle a broad pH range.

Resistance to biliary salts. Biliary salts have an emulsifying function, increasing the solubility of fats and fat-soluble vitamins to help in their absorption. This detergent effect is microbicidal because it can affect the phospholipids and fatty acids on the walls of the microorganisms. For a probiotic to efficiently colonize an animal's digestive tract, it is important for it to resist the action of biliary salts.



Lactic acid bacteria grown in agar with aniline blue indicator appeared blue.

From these in vitro tests, a theoretically ideal strain based on the best results obtained from the evaluated strains was defined. The ideotype had the highest growth velocity, greatest capacity to inhibit pathogens and greater capacity to resist biliary salts, different pH values and salinities. *Lactobacillus plantarum* was selected for later studies.

Probiotic in shrimp hatchery

The colonization of the digestive tract of shrimp larvae by beneficial bacteria can be an important strategy for the prevention of infirmities. The larvae are only totally colonized in the digestive tract when there is a transformation to the protozoa state with the initiation of exogenous feeding. Thus, the supply of a diet supplement with probiotics in this stage can lead to the establishment of a favorable intestinal bacterial microbiota from the beginning.

In an experimental hatchery, it was observed that from the nauplius V stage to postlarva 1, diet supplementation with *L. plantarum* increased the survival of the larva, with survival rates of 51 percent for those in the probiotic treatment and 21 percent for the control group. There was also a modification in the bacterial microbiota of the larvae, with an increase in the concentration of beneficial bacteria (lactic acid) and decrease in the population of *Vibrio* species (Fig. 1). Larvae fed with *L. plantarum* had even greater resistance to an experimental challenge with *V. harveyi*, with 80 percent survival in relation to 50 percent survival in the control group.




Fig. 1: Bacterial counts in shrimp larvae fed a control diet or a diet supplemented with *L. plantarum*.

Probiotic in experimental cultivation

In this study, experimental cultivation was conducted in clearwater for 60 days, beginning with the shrimp in the post-larvae 20 stage. A modification in the intestinal bacterial microbiota was observed, with a decrease in the population of *Vibrio* species and an increase of lactic acid bacteria (Fig. 2) in the shrimp fed with probiotics.




Fig. 2: Bacterial counts in the intestines of shrimp fed a control diet or a diet supplemented with *L. plantarum*.

However, there was no alteration in the survival rate (81 percent) and final average weight (6 grams) in both treatments. These results were expected, given that the shrimp were maintained under excellent cultivation conditions with 70 percent daily water exchange and a temperature of 30 degrees-C during the experiment.

At the end of cultivation, the shrimp were challenged with *V. harveyi* by intramuscular injection. After 12 hours, it was observed that the shrimp fed with probiotics had a 66 percent survival rate compared to the control animals' 40 percent survival.

Probiotic at shrimp farm

Studies conducted in controlled laboratory conditions are important, but their results are not always replicated on a commercial scale. Thus, in the final step of this work, the use of *L. plantarum* was evaluated under commercial conditions at the Universidade Federal de Santa Catarina Yakult Experimental Farm. Six, 12,000-m² ponds stocked at 14 shrimp/m² were used. The shrimp in three ponds received a diet supplemented with probiotics and three were maintained as a control.

After 75 days of cultivation, the shrimp whose diet was supplemented with probiotics had a higher survival rate and lower feed-conversion rate (Table 1), with no alteration in their growth rate. This result should be associated with a change in the intestinal bacterial microbiota of the shrimp fed with probiotics. The decrease in the *Vibrio* populations in the animals' intestines led to the increased survival of the shrimp in the ponds fed with probiotics.

Meanwhile, the lower feed conversion can be related to the capacity of the probiotic bacteria to stimulate the production of digestive enzymes in the host, improving the digestibility of the rations. A study conducted for a master's dissertation by Celso Buglione-Neto demonstrated that the addition of *L. plantarum* to the diet of shrimp improves the apparent digestibility of the protein in the ration, corroborating the authors' hypothesis.

(Editor's Note: This article was originally published in the November/December 2011 print edition of the Global Aquaculture Advocate.)

Authors

**DR. FELIPE DO NASCIMENTO VIEIRA**

Universidade Federal de Santa Catarina
Departamento de Aquicultura
Laboratório de Camarões Marinhos
Ademar Gonzaga Street, 1346
Florianópolis, Santa Catarina, Brazil

vieirafn@lcm.ufsc.br (<mailto:vieirafn@lcm.ufsc.br>).

**DR. JOSÉ LUIS MOURIÑO**

Universidade Federal de Santa Catarina
Departamento de Aquicultura
Laboratório de Camarões Marinhos
Ademar Gonzaga Street, 1346
Florianópolis, Santa Catarina, Brazil

**BRUNO CORREA DA SILVA, M.S.**

Universidade Federal de Santa Catarina
Departamento de Aquicultura
Laboratório de Camarões Marinhos
Ademar Gonzaga Street, 1346
Florianópolis, Santa Catarina, Brazil

**DR. WALTER QUADROS SEIFFERT**

Universidade Federal de Santa Catarina
Departamento de Aquicultura
Laboratório de Camarões Marinhos

Ademar Gonzaga Street, 1346
Florianópolis, Santa Catarina, Brazil



DR. LUIS ALEJANDRO VINATEA

Universidade Federal de Santa Catarina
Departamento de Aquicultura
Laboratório de Camarões Marinhos
Ademar Gonzaga Street, 1346
Florianópolis, Santa Catarina, Brazil

Copyright © 2023 Global Seafood Alliance

All rights reserved.