



- [GOAL Events](#)
- [Advocate Magazine](#)
- [Aquademia Podcast](#)
- [Blog](#)
- [Contact](#)

-
-
-
-
-

- [Log In](#)



- [About](#)
 - [Who We Are](#)
 - [Our History](#)
 - [Our Team](#)
 - [Sustainable Development Goals](#)
 - [Careers](#)
- [Membership](#)
 - [Overview](#)
 - [Our Members](#)
 - [Corporate Membership](#)
- [Resources](#)
- [Certification](#)
 - [Best Aquaculture Practices](#)
 - [Best Seafood Practices](#)

[Log In](#)

- [About](#)
 - [Who We Are](#)
 - [Our History](#)
 - [Our Team](#)
 - [Sustainable Development Goals](#)
 - [Careers](#)
- [Membership](#)
 - [Overview](#)
 - [Our Members](#)
 - [Corporate Membership](#)
- [Resources](#)
- [Certification](#)
 - [Best Aquaculture Practices](#)
 - [Best Seafood Practices](#)
- [GOAL Events](#)
- [Advocate Magazine](#)

- [Aquademia Podcast](#)
- [Blog](#)
- [Contact](#)



 Aquafeeds
Aquafeeds


Nutritional disorders and aquatic animal health

 [Responsible Seafood Advocate logo](#)

1 December 2000 Albert G.J. Tacon, Ph.D.



Nutritional characteristics and quality of most commercial aquafeeds is not always satisfactory



“Black death,” or ascorbic acid deficiency syndrome (AADS), is a nutritional deficiency disease of penaeid shrimp. Gross signs of AADS include large melanized lesions in the subcuticular tissues.
Photo by Don Lightner.

Dietary nutritional disorders in farmed aquatic animals can be broadly defined as diet-related imbalances due to “under-” or “over-” nutrition. These disorders are more readily apparent and prevalent as farming systems intensify and animals become less dependent upon endogenously produced natural food organisms and more dependent upon

artificially compounded diets. To date, the science of nutritional pathology – often incorrectly termed nutritional “diseases” – remains one of the least studied areas of finfish and crustacean pathology.

Diets for optimum health and disease resistance

As in humans, the nutritional intake and status of an animal has a profound effect upon its overall health and well-being, including its subsequent growth and resistance to environmental stressors and disease-causing agents. In general, the poorer the diet and nutritional status of an animal, the less able it is to combat disease agents and withstand overt environmental changes and stress. It is important to emphasize that all animals have dietary nutrient requirements for optimum health over and above those normally required for optimum growth and feed efficiency.

Sadly, most nutrient requirement studies to date were conducted under indoor laboratory conditions, where strict environmental controls are placed on test animals. They are not subjected to the normally varied outdoor rearing conditions and stresses, including varying water quality and husbandry conditions, and water-borne disease agents.

Nutrients that enhance health

To date, nutrients and dietary components reported to exert a positive response on the health and/or disease resistance (including immune response) of farmed finfish and crustaceans, have included specific essential amino acids (lysine, methionine, tryptophan, arginine, histidine, leucine, isoleucine), nucleotides, polysaccharides (peptidoglycans, beta 1,3, and 1,6 glucans, lipopolysaccharides), essential fatty acids (18:2 omega-6, 18:3 omega-3; 20:4 omega-6, 20:5 omega-3, 22:6 omega-3 depending upon species), sterols, phospholipids, and essential minerals (P, K, Mg), trace elements (Fe, Zn, Mn, Cu, Se, I), vitamins (B1, B2, B6, B12, pantothenic acid, niacin, biotin, folic acid, inositol, choline, D3, A, K3, E, and C) and carotenoids.

Causes of nutritional disorders

Under practical farming conditions, nutritional disorders may arise from a variety of causes, including:

- Deficiencies and imbalances due to poor feed formulation, excessive heat treatment during the feed manufacturing process (resulting in the loss or destruction of heat-sensitive nutrients), poor/prolonged feed storage (resulting in the loss of essential nutrients through oxidation or spoilage), or through nutrient loss on immersion in water through leaching or poor feed stability.
- Anti-nutrients and contaminants, including toxic polyamines, oxidized polyunsaturated fatty acids and toxic amino acids, heavy metal contaminants, anti-vitamin factors, specific enzyme inhibitors, toxic glycosides, toxic phenols, food allergens, microbial toxins, and specific synthetic contaminants (including pesticide residues and organochlorine compounds) and residues arising from ingredient/feed processing.

Common deficiency signs resulting from nutritional disorders

Examples of some major morphological signs and nutritional disorders reported in finfish fed diets with specific nutrient deficiencies and/or toxicities under experimental conditions are shown in Table 1. Nutritional diseases reported for penaeid shrimp include Ascorbic Acid Deficiency Syndrome (“Black Death Disease”), Cramped Muscle Syndrome, Chronic Soft Shell Syndrome, and Blue Disease.

Table 1. Major nutritional pathology conditions and reported causes in finfish.

SCOLIOSIS/LORDOSIS (bent/curved spine)	
Deficiency of:	Tryptophan Magnesium Phosphorus Vitamin C Essential fatty acids (lordosis)
Toxicity of:	Lead Cadmium Leucine Vitamin A Oxidized fish oil
CATARACT (white/cloudy eye)	
Deficiency of:	Methionine Tryptophan Zinc Magnesium Copper Selenium Manganese Vitamin A Riboflavin
Toxicity of:	Choline Oxidized fish oil
FIN EROSION	
Deficiency of:	Lysine Tryptophan Zinc Riboflavin Inositol Niacin Vitamin C
Toxicity of:	Lead Vitamin A

FATTY LIVER (pale/ceroid)	
Deficiency of:	Choline Essential fatty acids
Toxicity of:	Oxidized fish oil
EXOPHTHALMIA (popeye)	
Deficiency of:	Pantothenic acid Niacin Folic acid Vitamin A Vitamin E
Toxicity of:	Oxidized fish oil
FIN/SKIN HEMORRHAGE	
Deficiency of:	Riboflavin Pantothenic acid Niacin Thiamine Inositol Vitamin C Vitamin A Vitamin K
Toxicity of:	Oxidized fish oil

Conclusion

Pale coloration can be caused by inadequate dietary carotenoids. The top shrimp received no dietary carotenoids, while the darker animal below did receive carotenoids. Photo by Don Lightner.

Although it is generally believed that the nutritional characteristics and quality of most commercial aquafeeds is satisfactory, this is not always the case. It is imperative that farmers keep good records on feed use and species health/mortality (including the occurrence of morphological deficiency signs). Moreover, whenever economically feasible, it is strongly recommended that farmers undertake periodic spot checks on new batches of feed to monitor feed quality, including label declarations.

However, equally important as the aquafeed itself, is the need for farmers to employ good on-farm feed management protocols concerning feed storage and feed use on the farm. All too often it is believed that a “high-quality” feed works by itself. It does not. It must be properly managed. There is no substitute for good on-farm feed and water management to ensure that all the nutritional attributes of feed are transferred in their entirety to the cultured target species.

(Editor's Note: This article was originally published in the December 2000 print edition of the Global Aquaculture Advocate.)

Now that you've finished reading the article ...

... we hope you'll consider supporting our mission to document the evolution of the global aquaculture industry and share our vast network of contributors' expansive knowledge every week.

By becoming a Global Seafood Alliance member, you're ensuring that all of the pre-competitive work we do through member benefits, resources and events can continue. Individual membership costs just \$50 a year.

Not a GSA member? Join us.

[Support GSA and Become a Member](#)

Author





-  Albert G.J. Tacon, Ph.D.

Albert G.J. Tacon, Ph.D.

Aquatic Feeds and Nutrition
(AQUAFAN) Program
The Oceanic Institute
Makapuu Point
Waimanalo, Hawaii 96795 USA

[103,114,111,46,101,116,117,116,105,116,115,110,105,99,105,110,97,101,99,111,64,110,111,99,97,116,97]

Share

-  [Share via Email](#)
-  [Share on Twitter](#)
-  [Share on Facebook](#)
-  [Share on LinkedIn](#)

Tagged With

[Albert G. J. Tacon](#) [Nutritional disorders](#) [The Oceanic Institute](#)

Related Posts

Aquafeeds

[Finishing feeds tailor fatty acid composition in tilapia fillets](#)

By feeding different oils at different points in a fish's life cycle, such as finishing feeds, farmers can tailor the composition of fillets to maximize value and minimize fish oil use.

Intelligence

[Increasing seafood consumption, from strategy to action](#)

Despite global strategies and political declarations that strive to support healthy diets with increased seafood consumption and regular physical activity, the world is seeing a rapid upsurge in obesity and under-nutrition, particularly in urban settings.

Intelligence

[Making the case for a seafood-based economy](#)

The “Towards a Seaweed Based Economy” report made the case that poor nutrition is the major cause for the pandemics of obesity and several chronic diseases. It recommended increased use of food-production systems like integrated multi-trophic aquaculture.

Health & Welfare

[Protein hydrolysates in larval fish nutrition](#)

Protein hydrolysates are currently used as partial substitutes for fishmeal in high-quality artificial diets since their inclusion in feeds promotes growth and survival.

About The Advocate

The Responsible Seafood Advocate supports the Global Seafood Alliance's (GSA) mission to advance responsible seafood practices through education, advocacy and third-party assurances.

[Learn More](#)

Search Responsible Seafood Advocate



Advertising Opportunities

[2022 Media & Events Kit](#)

Categories

[Aquafeeds](#) [Aquafeeds](#) > [Health & Welfare](#) > [From Our Sponsors](#) > [Innovation & Investment](#) > [Intelligence](#) > [Responsibility](#) > [Fisheries](#) > [Artículos en Español](#) >

Don't Miss an Article

Featured

- [Health & Welfare](#) [An update on vibriosis, the major bacterial disease shrimp farmers face](#)
- [Intelligence](#) [A seat at the table: Fed By Blue team says aquaculture needs a stronger voice](#)
- [Responsibility](#) [Quantifying habitat provisioning at macroalgae cultivation locations](#)

Popular Tags

All Tags








Recent

- [Fisheries Second Test: Another filler for the fisheries category.](#)
- [Fisheries Test: This is filler for the fisheries Category.](#)
- [Aquafeeds Test Article](#)
- [Responsibility Study: Climate change will shuffle marine ecosystems in unexpected ways as ocean temperature warms](#)
- [Health & Welfare Indian shrimp researchers earn a patent for WSSV diagnostic tool](#)



- [About](#)
- [Membership](#)
- [Resources](#)
- [Best Aquaculture Practices \(BAP\)](#)
- [Best Seafood Practices \(BSP\)](#)
- [GOAL Events](#)
- [Advocate Magazine](#)
- [Aquademia Podcast](#)
- [Blog](#)
- [Contact](#)

Stay up to date with GSA

- 
- 
- 
- 
- 

Copyright © 2024 Global Seafood Alliance
All rights reserved.

[Privacy](#)

[Terms of Use](#)

[Glossary](#)