### Global Seafood Alliance Logo

- GOAL Events
- Advocate Magazine
- Aquademia Podcast
- Blog
- Contact
- 0
- . 4
- X
- in
- 🔼
- Log In

- About
  - Who We Are
  - Our History
  - o Our Team
  - Sustainable Development Goals
  - Careers
- Membership
  - Overview
  - Our Members
  - Corporate Membership
- Resources
- Certification
  - Best Aquaculture Practices
  - Best Seafood Practices

Search...



# <u>Log In</u>

- About
  - Who We Are
  - Our History
  - o Our Team
  - Sustainable Development Goals
  - Careers
- Membership
  - o <u>Overview</u>
  - Our Members
  - o Corporate Membership
- Resources
- Certification
  - Best Aquaculture Practices
  - Best Seafood Practices
- GOAL Events
- Advocate Magazine

- Aquademia Podcast
- Blog
- Contact



## Paddy culture: Chinese mitten-handed crabs

Responsible Seafood Advocate logo

1 December 2002 Qigen Liu Liqiao Chen



# Environmentally friendly production approach benefits both the crabs and rice



Increasing production of Chinese mitten-handed crabs has led to price drops and rising demand.

The Chinese mitten-handed crab (*Eriocheirn sinensis*) has been the fastest growing aquaculture species in China during the last decade. This migratory species spends most of its 15-month lifetime in freshwater and migrates to brackish water for spawning. Its consumption in the country dates back at least a millennium.

Mitten-handed crabs have been cultured in China for about 50 years. Their production increased over 30 times between 1990 and 1999, and reached 286,000 metric tons (MT) in 2001. The crabs are highly regarded by farmers

and in domestic and international markets. Culture of this species is one of the most profitable aquaculture activities in China today.

# **Culture stages**

The culture of *E. sinensis* could begin from the megalopa larval stage, when it can adapt to live in freshwater. However, the survival rate for megalopa larvae in large bodies of water is usually very low, so production is generally divided into the larval culture stage and the grow-out stage.

In the first stage, megalopa larvae are reared for five to six months in a nursery until they become juveniles of 5 to 20 grams called "button-sized" crabs by the Chinese crab farmers. In the second stage, which lasts seven to 10 months, the juveniles are grown to market size in ponds, pens in lakes, or rice paddies. Although production currently comes mainly from pen culture in lakes, paddy culture has also increased rapidly in recent years.

# Culture in rice paddies

Crab culture in rice paddies is an environmentally friendly production approach that has become quite popular. Paddy culture benefits both the crabs and the rice.

The rice plants provide shelter to the crabs and reduce sibling cannibalism, and improve environmental conditions by absorbing the nutrients excreted by the crabs in their waste. An additional benefit to the crabs is that the rice paddy usually attracts insects and other organisms that are natural and nutritious foods for the crabs. On the other hand, rice plants benefit from the crabs' presence by their manuring and grazing on weeds.

### **Construction and preparation**

The ditch-and-pit system necessary to a rice paddy often covers 20 to 40 percent of the paddy area. The system commonly includes a ditch around the paddy and about 1.2 meters from the dikes, cross ditches in the central part of the paddy, and pits at the intersections of the ditches.

The ditch-and-pit system is dug after the paddies are ploughed but before the rice seedlings are planted. The ditches are typically 1 meter wide and 0.6 meter deep, while pits are wider, deeper, and square-shaped to hold more water for the crabs when the rice paddy dries up. Dikes are raised to a higher level and strengthened to contain more water than an ordinary paddy. Walls to prevent crabs from escaping are built around the paddy or even the whole farm. Paddies are disinfected with quicklime or potassium permanganate before stocking with crabs.

## Seedstock source and stocking

Distinct local populations of this crab species are defined by their spawning grounds. These groups include the Liaohe River, Yanghzi River and Oujiang River populations.

Crabs from the Yanghzi River population grow to a larger average size and reportedly taste better than crabs from the other populations. Consequently, seedstock from the Yanghzi River population are normally used. Animals of about 7 grams are disinfected by dipping in 4 percent saltwater for five minutes before being stocked in rice paddies at about 4,500 to 6,000 animals per hectare.

# **Rice planting**

The rice paddies are planted with fewer rice seedlings than typical rice paddies. The distance between rows of seedlings is 20 cm and the distance between the lines is 5 cm.

# **Feeding**

In March and April, when the water temperature is still below 15 degrees-C, the crabs usually do not eat much. Consequently, it is only necessary to feed the crabs during sunny weather. Rations made of various oil plant meals, rice bran, or fishmeal are fed at around 2 percent of the crabs' body weight (BW). The foods are often made into dough to reduce leaching.

When the temperature rises above 15 degrees-C, the crabs are fed once every three to five days at about 5 percent of their body weight. At this time, various aquatic plants are also planted in the ditches. In May and June, the crabs are fed every day or two at 5 to 8 percent BW.

July to October is the main growing season for the crabs, when animals must be fed twice daily in the morning and evening. The daily food ration increases to 5 to 10 percent BW, with two-thirds of the daily ration applied in the second feeding. More animal food components – including trash fish, mud snails, or clam meat – are added to the crabs' diets. In late October and November, as water temperature decreases, so does the feeding.

## **Husbandry management**

This production system has both benefits and conflicts. For example, pesticide treatments needed for the rice must be done properly to prevent crabs from being affected or even poisoned to death. The paddies are filled with more water, which is exchanged an hour or two after the pesticide is applied. Quicklime at 40 to 50 ppm is also used once a month to disinfect the water and help the crabs molt.

It is also necessary for the plants to occasionally have dry soil, while the crabs need water. With such constraints, it is crucial to harmonize the needs of both species to have successful crops with both.

### Harvest

The crabs are harvested simply by draining the water at the end of a ditch. Average production from paddy culture typically reaches 300-450 kilograms per hectare.

### Conclusion

Cultured production of Chinese mitten- handed crabs has steadily increased during the last decade, leading to drops in prices and stimulating demand. Although production currently comes mainly from pen culture in lakes, culture in rice paddies has also increased.

Because of its environmentally friendly nature, crab culture in rice paddies will likely continue to grow. Pen culture may be scaled back due to the impact of crabs grazing on aquatic plants and organic loading from crab wastes and uneaten feeds.

(*Editor's Note: This article was originally published in the December 2002 print edition of the* Global Aquaculture Advocate.)

# Now that you've finished reading the article ...

... we hope you'll consider supporting our mission to document the evolution of the global aquaculture industry and share our vast network of contributors' expansive knowledge every week.

By becoming a Global Seafood Alliance member, you're ensuring that all of the pre-competitive work we do through member benefits, resources and events can continue. Individual membership costs just \$50 a year.

#### Not a GSA member? Join us.

Support GSA and Become a Member

#### Authors

• **Qigen** Liu

Qigen Liu

East China Normal University North Zhongshan Road Shanghai, China

[110,99,46,117,100,101,46,117,102,104,115,64,117,105,108,103,113]

• **Liqiao** Chen

Liqiao Chen

East China Normal University North Zhongshan Road Shanghai, China

#### **Share**

- Share via Email
- **Share on Twitter**
- **f** Share on Facebook
- in Share on LinkedIn

#### **Tagged With**

Qigen Liu Liqiao Chen mitten-handed crabs

#### **Related Posts**

Responsibility

### **Carp polyculture in India**

Carp polyculture with other species originally established through government projects has grown in India thanks to the application of advancing technologies and private financial investments.

#### Intelligence

### **Evolution of shrimp farming in Bangladesh**

The social and environmental improvements – in combination with job-generating foreign investment – have led to wide acceptance of shrimp farming in Bangladesh.

Health & Welfare

### Freshwater fish culture in China

About 60 percent of China's aquaculture production is freshwater fish. Carps top the list, although many other native and non-native species are also raised.

Responsibility

### Giving a lift to tilapia farmers in Myanmar

The U.S. Soybean Export Council's farm demonstrations, farmer education efforts and formulated feeds are boosting tilapia production in rural areas of Myanmar, where indigenous carp are more commonly raised.

#### **About The Advocate**

The Responsible Seafood Advocate supports the Global Seafood Alliance's (GSA) mission to advance responsible seafood practices through education, advocacy and third-party assurances.

#### Learn More

Search Responsible Seafood Advocate Search Search



#### **Advertising Opportunities**

2022 Media & Events Kit

### Categories

Aquafeeds Health & Welfare From Our Sponsors Innovation & Investment Intelligence Intelligence Nesponsibility Fisheries Artículos en Español

#### Don't Miss an Article

#### **Featured**

- Health & Welfare An update on vibriosis, the major bacterial disease shrimp farmers face
- Intelligence A seat at the table: Fed By Blue team says aquaculture needs a stronger voice
- Responsibility Quantifying habitat provisioning at macroalgae cultivation locations

#### **Popular Tags**

All Tags ✓

#### Recent

- Fisheries Second Test: Another filler for the fisheries category
- Fisheries Test: This is filler for the fisheries Category
- Aguafeeds Test Article
- Responsibility Study: Climate change will shuffle marine ecosystems in unexpected ways as ocean temperature warms
- Health & Welfare Indian shrimp researchers earn a patent for WSSV diagnostic tool



- About
- Membership
- Resources
- Best Aquaculture Practices (BAP)
- Best Seafood Practices (BSP)
- GOAL Events
- Advocate Magazine
- Aquademia Podcast
- Blog
- Contact

#### Stay up to date with GSA

- @
- . 🗴
- ^
- •

Copyright  $\ensuremath{\text{@}}$  2024 Global Seafood Alliance All rights reserved.

Privacy Terms of Use Glossary