Global Seafood Alliance Logo

- GOAL Events
- Advocate Magazine
- Aquademia Podcast
- Blog
- Contact
- 0
- •
- X
- in
- 🔼
- Log In

- About
 - Who We Are
 - Our History
 - o Our Team
 - Sustainable Development Goals
 - <u>Careers</u>
- <u>Membership</u>
 - o <u>Overview</u>
 - Our Members
 - Corporate Membership
- Resources
- Certification
 - Best Aquaculture Practices
 - Best Seafood Practices

Search...



Log In

- About
 - Who We Are
 - Our History
 - o Our Team
 - Sustainable Development Goals
 - Careers
- Membership
 - o <u>Overview</u>
 - Our Members
 - o Corporate Membership
- Resources
- Certification
 - Best Aquaculture Practices
 - Best Seafood Practices
- GOAL Events
- Advocate Magazine

- Aquademia Podcast
- Blog
- Contact



Pangasius diet trials show promise of DDGS

Responsible Seafood Advocate logo

25 January 2016 Dr. Jerry Shurson Dr. Budi Tangendjaja Chien Tran Trong Le Thi Ngoc Thuy



Corn distillers dried grains with solubles an excellent source of energy and digestible nutrients



Figure 1. Fillet color of pangasius from feeding 0% DDGS diets for 118 days.

In the quest to find suitable, cost effective, alternative ingredients to use in aquaculture feeds, results from a recent study conducted by the U.S. Grains Council have shown that corn distillers dried grains with solubles (DDGS) is an economical and excellent source of energy and digestible nutrients in pangasius diets.

The aquaculture industry in Vietnam is growing rapidly and the production of *Pangasius hypophthalmus*, also known as swai or tra — the predominant species of fresh water fish produced in the Mekong River — accounts for more than 1 million metric tons (MMT) produced annually. Vietnam is the third-largest fish exporter in the world, and most of the pangasius produced is exported to many different countries. However, high quality, nutrient dense, economical feed ingredients are needed to reduce cost and enable this rapidly growing industry to remain globally competitive. Corn and corn co-products are economical and excellent dietary energy sources for various

aquaculture species, but naturally contain xanthophylls (yellow pigment), which often causes concerns about adding these ingredients at relatively high dietary inclusion rates to diets for some fish species because of the potential "yellowing" of fillets. Achieving a desirable "white" fillet color is essential for meeting consumer preferences and to maintain these important export markets.

DDGS, co-product of ethanol

Corn DDGS is a co-product of ethanol production in the U.S., is produced in large quantities (> 35 MMT), and is used as an economical source of energy, digestible protein and amino acids, and digestible phosphorus in feeds for cattle, swine, poultry, and some aquaculture diets around the world. In 2015, about 12 MMT of DDGS were exported for use in various animal feeds. While the use of DDGS in aquaculture diets has been limited, its use has been increasing rapidly but it is still an unfamiliar feed ingredient for much of the global aquaculture industry. One of the major reasons for the limited use of DDGS until now is that a limited number of research studies have been published to evaluate DDGS use in various aquaculture diets.



Figure 2. Fillet color of pangasius from feeding 5% DDGS diets for 118 days.

Limited information available

However, despite a limited number of published, there is good research evidence indicating that traditional high-oil DDGS can be added to channel catfish (*Ictalurus punctatus*) diets at dietary levels up to 30 percent. Tidwell et al. (1990) conducted an experiment over an 11-week period where channel catfish fingerlings were fed diets containing 0, 10, 20 and 40 percent DDGS by replacing some of the corn and soybean meal.

After the 11-week feeding period, there were no significant differences in individual fish weight, percentage survival, feed conversion, or protein efficiency ratio (PER) among dietary treatments. Similarly, Webster et al. (1993) conducted a study using cage reared juvenile catfish that were fed diets containing 0, 10, 20, or 30 percent DDGS, which partially replaced corn and soybean meal in the diets. No differences in individual fish weights, survival, feed conversion, carcass composition, carcass waste (head, skin, viscera), and organoleptic properties of the fillets were observed among dietary treatments, indicating that up to 30 percent DDGS can be added to channel catfish diets with no negative effects on growth performance, carcass composition, or flavor qualities of the fillets.



Figure 3. Fillet color of pangasius from feeding 10% DDGS diets for 118 days.

However, subsequent studies conducted by Robinson and Li (2008) and Lim et al. (2009) showed that the addition of crystalline lysine to catfish diets was needed to support optimal growth performance when DDGS was added up to 30 or 40 percent of the diet. In both of these studies, body fat increased when DDGS was added to the diet compared to fish fed the control diets. Most recently, Zhou et al. (2010) replaced soybean meal and corn meals in juvenile hybrid catfish (channel catfish × blue catfish *I. furcatus*) and observed that diets containing 30 percent DDGS provided good growth, feed conversion, and protein retention.

No studies have been published that have evaluated to effects of adding up to 15 percent DDGS to pangasius diets on growth performance and fillet color. Corn naturally contains xanthophyll pigments, which are concentrated by a factor of 3 times in DDGS, resulting in concentrations of 25 to 50 mg/kg xanthophylls. The bioavailability of xanthophylls in DDGS has not been determined, but common recommendations for the use of corn and corn coproducts in American catfish diets indicate that the total diet xanthophyll content should not contain more than 12 mg/kg to achieve acceptable fillet color. As a result, one of the key questions involving DDGS use in pangasius diets is whether this amount of pigment will negatively affect fillet color when 15 percent DDGS diets are is fed to pangasius during a four- to six-month feeding period.



Figure 4. Fillet color of pangasius from feeding 15% DDGS diets for 118 days.

Studies on effect on fillet color

To answer this question, and evaluate the effects of feeding diets containing up to 15 percent DDGS to pangasius on growth performance, the U.S. Grains Council partnered with Pangafeed JSC in Dong Thap Province, Vietnam to conduct a feeding trial. Six thousand pangasius fingerlings (40 g in body weight) were randomly allotted to one of 16 floating cages (300 fish/cage) made of nylon netting located in a 0.5 ha pond that was 3 meters deep on the Hung Vuong Co. experimental farm, Mekong Delta Sadec Province. Four dietary treatments were evaluated over a 118-day feeding period and included 0, 5, 10, or 15 percent DDGS along with soybean meal, rice bran, cassava, fishmeal, wheat and wheat bran (Table 1). The DDGS source used contained 27.7 percent crude protein, 8.9 percent crude fat, 7.7 percent crude fiber, 0.79 percent lysine, 1.1 percent methionine+cysteine, 1.0 percent threonine, and 0.22 percent tryptophan. All diets were formulated to contain the same nutrient content and were manufactured into floating pellets for both the starter (3-4 mm) and grower (5-6 mm) phases.

Ingredients	0% DDGS	5% DDGS	10% DDGS	15% DDGS
Soybean meal	487	468.2	456.2	447
Rice bran, full fat	224.5	145.1	154.6	133.6
Wheat bran	0	0	0	0
Defatted rice bran	50	100.1	50	0
Cassava	120	120.1	120.1	165.1
Wheat	50	50	50	50
Fishmeal, 62%	40	40	40	47.5
Fishmeal, 55%	12	12	14.5	11
DDGS	0	50	100	150
Premix	10.5	10.5	10.5	10.5
Salt	6	4	4	4
Monosodium phosphate	0	0	0	0
Total	1000.0	1000.0	1000.0	1000.0
Ingredients	0% DDGS	5% DDGS	10% DDGS	15% DDGS
Ingredients Soybean meal	0% DDGS 447	5% DDGS 428	10% DDGS 424	15% DDGS 412
_				
Soybean meal	447	428	424	412
Soybean meal Rice bran, full fat	447 233	428 202	424 166	412 125
Soybean meal Rice bran, full fat Wheat bran	447 233 50	428 202 50	424 166 50	412 125 15
Soybean meal Rice bran, full fat Wheat bran Defatted rice bran	447 233 50 0	428 202 50 0	424 166 50 0	412 125 15 0
Soybean meal Rice bran, full fat Wheat bran Defatted rice bran Cassava	447 233 50 0 120	428 202 50 0 120	424 166 50 0 120	412 125 15 0 120
Soybean meal Rice bran, full fat Wheat bran Defatted rice bran Cassava Wheat	447 233 50 0 120 80	428 202 50 0 120 80	424 166 50 0 120 80	412 125 15 0 120 117.5
Soybean meal Rice bran, full fat Wheat bran Defatted rice bran Cassava Wheat Fishmeal, 62%	447 233 50 0 120 80 30	428 202 50 0 120 80 30	424 166 50 0 120 80 30	412 125 15 0 120 117.5
Soybean meal Rice bran, full fat Wheat bran Defatted rice bran Cassava Wheat Fishmeal, 62% Fishmeal, 55%	447 233 50 0 120 80 30 26	428 202 50 0 120 80 30 26	424 166 50 0 120 80 30 16	412 125 15 0 120 117.5 30
Soybean meal Rice bran, full fat Wheat bran Defatted rice bran Cassava Wheat Fishmeal, 62% Fishmeal, 55% DDGS	447 233 50 0 120 80 30 26	428 202 50 0 120 80 30 26 50	424 166 50 0 120 80 30 16 100	412 125 15 0 120 117.5 30 15
Soybean meal Rice bran, full fat Wheat bran Defatted rice bran Cassava Wheat Fishmeal, 62% Fishmeal, 55% DDGS Premix	447 233 50 0 120 80 30 26 0 8	428 202 50 0 120 80 30 26 50	424 166 50 0 120 80 30 16 100 8	412 125 15 0 120 117.5 30 15 150 8

At the end of the 118-day feeding period, there were no differences in growth performance among the dietary treatments. Body weights of fish fed 0, 5, 10, and 15 percent DDGS diets were 471, 472, 470, and 490 g, respectively. Feed to gain ratios were 1.59, 1.62, 1.56, and 1.53 for fish fed diets containing 0, 5, 10, and 15 percent DDGS respectively. Mortality of fish was not different among dietary treatments and ranged between 3.7 to 4.9 percent. Fillet yield was slightly improved by feeding the 15 percent DDGS diets (531 g/kg) compared with

feeding the control diet containing no DDGS (526 g/kg). Fillet color was evaluated at 3 locations (anterior, middle, and posterior positions) of fillets using L^* , a^* , and b^* in the Hunter Lab colorimetric system. The L^* indicates lightness or darkness of color, a^* indicates redness of color, and b^* indicates yellowness of color. There were no differences in each of the color score measures for fillet color among dietary treatments after feeding them for 118 days.

To evaluate the long-term effects of feeding DDGS diets on fillet color, 50 fish were retained and kept in smaller cages for an additional 66 days, and harvested at 0.9-1.0 kg body weight. There were no differences in L* or b* scores of fillets among dietary treatments, with minor differences in a* between fillets from fish fed the 10 percent DDGS diets compared with the control diets, but not for fish fed 15 percent DDGS diets.

Perspectives

Results from this feeding trial showed that the addition of up to 15 percent DDGS as an energy and protein source, to replace rice bran and partially replace soybean meal in starter and grower pangasius diets, provides excellent growth performance, with no negative effects on mortality, fillet yield or fillet color.

Authors

• Dr. Jerry Shurson

Dr. Jerry Shurson

Department of Animal Science University of Minnesota 1988 Fitch Ave. St. Paul. MN USA 55108

[117,100,101,46,110,109,117,64,49,48,48,115,114,117,104,115]

• Dr. Budi Tangendjaja

Dr. Budi Tangendjaja

U.S. Grains Council Komplek BPT No. 12 RT 1/3 Banjarwangi Ciawi, Bogor 16720 Indonesia

• Chien Tran Trong

Chien Tran Trong

U.S. Grains Council 13/F Hanoi Tower 19 Hai Ba Trunk Hanoi, Vietnam

• DLe Thi Ngoc Thuy

Le Thi Ngoc Thuy

Hung Vuong Tay Nam Pangafeed JSC Sa Dec City, Dong Thap Province, Vietnam

Share

- Share via Email
- **Share on Twitter**
- **f** Share on Facebook
- in Share on LinkedIn

Tagged With

<u>Vietnam ethanol Jerry Shurson Budi Tangendjaja Chien Tran Trong Le Thi Ngoc Thuy nutrients pangasius solubles</u>

Related Posts

Intelligence

Risk v. hazard: A dispassionate look at pangasius

Vietnam's pangasius industry captivated the global seafood industry, environmental organizations and the mass media. A scientific look at harmful substances detected in exported fillets and the reporting of the associated health risks through the media finds wide disparities.

Aquafeeds

Natural feed additives enhance rainbow trout performance

Various additives, many of plant origin, have been explored to replace the use of antibiotics in aquafeed. The authors compared the performance of rainbow trout fed diets containing oxytetracycline or mineral additives yellow loess and Song-Gang stone powder.

Aquafeeds

Soy-based feeds evaluated for production of Amazonian paiche

A cooperative feeding demonstration conducted by USSEC and Amazon Fish Products S.A. evaluated the use of soy to partially replace fishmeal in diets for paiche, or pirarucu, the largest freshwater fish species.

Intelligence

Global Farmed Finfish Production Outlook: Slower-paced growth

The GOAL 2016 survey for global production of farmed fish shows that most major finfish groups increased their farmed production in 2015. The total production of species and countries covered by the survey is forecasted to reach 42 MMT in 2016.

About The Advocate

The Responsible Seafood Advocate supports the Global Seafood Alliance's (GSA) mission to advance responsible seafood practices through education, advocacy and third-party assurances.

Learn More

Search Responsible Seafood Advocate | Search

Search **Q**



Advertising Opportunities

2022 Media & Events Kit

Categories

<u>Aquafeeds Aquafeeds</u> <u>Health & Welfare</u> <u>From Our Sponsors</u> <u>Innovation & Investment</u> <u>Intelligence</u> <u>Nesponsibility</u> <u>Fisheries</u> <u>Artículos en Español</u>

Don't Miss an Article

Featured

- Health & Welfare An update on vibriosis, the major bacterial disease shrimp farmers face
- <u>Intelligence A seat at the table: Fed By Blue team says aquaculture needs a stronger voice</u>
- Responsibility Quantifying habitat provisioning at macroalgae cultivation locations

Popular Tags

All Tags ✓

Recent

- Fisheries Second Test: Another filler for the fisheries category
- Fisheries Test: This is filler for the fisheries Category
- Aguafeeds Test Article
- Responsibility Study: Climate change will shuffle marine ecosystems in unexpected ways as ocean temperature warms
- Health & Welfare Indian shrimp researchers earn a patent for WSSV diagnostic tool



- About
- Membership
- Resources
- Best Aquaculture Practices (BAP)
- Best Seafood Practices (BSP)
- GOAL Events
- Advocate Magazine
- Aquademia Podcast
- Blog
- Contact

Stay up to date with GSA

Copyright © 2024 Global Seafood Alliance All rights reserved.

Privacy Terms of Use Glossary