<u>Global Seafood Alliance Logo</u>

- GOAL Events
- <u>Advocate Magazine</u>
- Aquademia Podcast
- <u>Blog</u>
- <u>Contact</u>
- 0
- f
- 🗙
- in
- 🕨
- <u>Log In</u>
- \square
 - <u>About</u>
 - Who We Are
 - <u>Our History</u>
 - <u>Our Team</u>
 - Sustainable Development Goals
 - <u>Careers</u>
 - <u>Membership</u>
 - <u>Overview</u>
 - <u>Our Members</u>
 - Corporate Membership
 - <u>Resources</u>
 - <u>Certification</u>
 - Best Aquaculture Practices
 - Best Seafood Practices

Search...

Q

<u>Log In</u>

- <u>About</u>
 - Who We Are
 - <u>Our History</u>
 - Our Team
 - Sustainable Development Goals
 - <u>Careers</u>
- <u>Membership</u>
 - <u>Overview</u>
 - Our Members
 - Corporate Membership
- <u>Resources</u>
- <u>Certification</u>
 - <u>Best Aquaculture Practices</u>
 - Best Seafood Practices
- GOAL Events
- <u>Advocate Magazine</u>

- Aquademia Podcast
- Blog
- Contact



Health & Welfare Health & Welfare

Polyploidy in shrimp

Responsible Seafood Advocate logo 1 December 2001 Arlo W. Fast, Ph.D. James Wyban, Ph.D.



Advanced technology for improved performance, genetic and environmental security




Fig. 1: Normal, diploid Litopenaeus vannamei embryo at blastula stage. PB1 = first polar body, PB2 = second polar body, FM = fertilization membrane around developing embryo. Photo by Ligiao Chen, East China Normal University, Shanghai, China.

Most of the exceptional production increases in cultured shrimp production experienced during the past 20 years (from 2 percent of total world shrimp production in 1980 to more than 30 percent in 2000) were achieved using wild shrimp stocks. Now, cost-effective production increases require closed-lifecycle technology.

Domestication is necessary for improved cultivars development through genetic selection, and is essential for maintaining specific pathogen-free and specific pathogen-resistant stocks. Closed-life-cycle culture is fundamental to virtually all other mass-produced foods, including chickens, pigs, cattle, and agricultural plants.

Ploidy manipulation

The authors and co-workers are developing new technology based on ploidy manipulation in shrimp to improve culture performance and protect our valuable shrimp stocks. Ploidy refers to the number of chromosome sets carried by each cell in plants or animals. Most animals are diploid, meaning they have two copies of every chromosome. Using genetic techniques, chromosome numbers can be increased to three (triploid) or four (tetraploid) chromosome sets.

Advantages

Polyploidy, the application of triploid or tetraploid chromosomes, is widely used in cultured fish, shellfish, and plants. Usually, triploids are sterile, which can result in superior culture performance. Triploidy disrupts hormonal balances, gametogenesis, and sexual maturation. Energy devoted to sexual maturation in normal diploids can be redirected to enhanced somatic growth in triploids.

Since triploid stocks are sterile, concerns about the use of exotic species outside their normal range are eliminated. Sterility also reduces concerns about escapement of genetically distinct, domesticated stocks that might otherwise breed with wild stocks and negatively impact the population genetics of wild stocks.

Creation and maintenance of improved, domesticated shrimp stocks require large financial investments. Polyploid technology also offers a way to protect these investments.

Producing triploids

Triploids can be produced by two means. First, traumatic shock can be applied soon after egg fertilization, causing reabsorption of the second polar body into the zygote (Fig. 1). The second polar body, a product of egg meiosis, is normally extruded following egg fertilization. After polar body reabsorption, all subsequent cells of that individual are triploid with three sets of chromosomes. With this method, triploidy induction if often less than 100 percent effective.

A second method to produce triploids involves breeding normal diploids with tetraploids. This type of cross results in 100 percent triploid progeny. This method first requires that tetraploids be produced. Shocking eggs soon after fertilization produces tetraploids. But shock must be applied after the diploid chromosome set has replicated in preparation for first cell division, but before cell division begins. This disruption of first cell division results in the cell having four chromosome sets, which are maintained when the cell continues dividing.

Induced polyploidy in shrimp




Fig. 2: Chromosome spreads of diploid and tetraploid *Litopenaeus vannamei* embryo cells. A = diploid cell chromosomes ($2N \cong 88$). B = tetraploid cell chromosomes ($4N \cong 176$). Photos by Liqiao Chen, East China Normal University, Shanghai, China.

Polyploidy induction has only recently been achieved with shrimp, with most of this research occurring in China. Triploid and tetraploid fleshy shrimp (*Penaeus chinensis*) were produced using both temperature and chemical shocks. Chemical shocks with cytochalasin B were effective, but often produced mosaics, which is a combination of ploidy cell types within an individual.

Tetraploid *Sicyonia ingesta* shrimp were produced using both warm- and cold-temperature shocks, and triploid *Litopenaeus vannamei* have been produced using cold-temperature shocks. Arlo Fast and his colleagues successfully produced tetraploid *L. vannamei* using cold-temperature shocks to disrupt first mitotic cell division (Fig. 2). High Health Aquaculture in Kona, Hawaii, USA is attempting to produce polyploid *P. monodon* and *L. vannamei*, supported in part by National Science Foundation funding.

Comparing diploid and triploid shrimp

One study from China reports growth comparisons between diploid and triploid shrimp. Xiang et al. (1999) conducted growth trials with triploid *P. chinensis* in small, indoor tanks. Triploid shrimp were cultured for four months at three stocking densities (88, 182, and 200 shrimp per cubic meter) and their growth compared with diploid shrimp stocked at 88 shrimp per cubic meter.

At all densities, triploid shrimp grew larger than diploids, averaging 29.4, 23.8 and 1.5 percent larger for the triploids, respectively. Xiang also found that the ovaries of triploid shrimp were "undeveloped," while no differences were observed between the testes of triploid and diploids.

Conclusion

Researchers are currently developing polyploid (*P. monodon*) and *L. vannamei*. When these shrimp are achieved, their parameters for growth, health, maturation, and other factors will be carefully evaluated against diploid performance. Technology for the mass-production of polyploids can then be established.

In the shrimp-farming future, the most advanced strains of domesticated shrimp will likely be these polyploids, with faster growth, more efficient feed conversion, and greater disease resistance. Once triploid culture performance is documented, shrimp culture based on the use of sterile postlarvae can lead to significantly improved shrimp cultivars. Genetics developers will then be able to release stocks that have been withheld until genetic security is assured.

Note: Cited references are available from the authors.

(*Editor's Note: This article was originally published in the December 2001 print edition of the* Global Aquaculture Advocate.)

Now that you've finished reading the article ...

... we hope you'll consider supporting our mission to document the evolution of the global aquaculture industry and share our vast network of contributors' expansive knowledge every week.

By becoming a Global Seafood Alliance member, you're ensuring that all of the pre-competitive work we do through member benefits, resources and events can continue. Individual membership costs just \$50 a year.

Not a GSA member? Join us.

Support GSA and Become a Member

Authors

• 🖳 Arlo W. Fast, Ph.D.

Arlo W. Fast, Ph.D.

University of Hawaii at Manoa Hawaii Institute of Marine Biology P.O. Box 1056 Kaneohe, Hawaii 96744 USA

[116,101,110,46,97,104,111,108,97,64,116,115,97,102,111,108,114,97]

• 🛃 James Wyban, Ph.D.

James Wyban, Ph.D.

High Health Aquaculture Kurtistown, Hawaii, USA

Share

- Share via Email
- **Share on Twitter**
- **F** Share on Facebook
- in <u>Share on LinkedIn</u>

Tagged With

Polyploidy in shrimp James Wyban Arlo W. Fast

Related Posts

Health & Welfare

CSIRO research evaluates technologies to produce genetically protected, all-female shrimp

In Australia, CSIRO is investigating techniques to produce reproductively sterile, all-female shrimp populations through polyploidy, irradiation, or genetic engineering.

Responsibility

A helping hand to lend: UK aquaculture seeks to broaden its horizons

Aquaculture is an essential contributor to the world food security challenge, and every stakeholder has a role to play in the sector's evolution, delegates were told at the recent Aquaculture's Global Outlook: Embracing Internationality seminar in Edinburgh, Scotland.

Health & Welfare

Improving shellfish aquaculture through triploidy

Triploid bivalves have sparked worldwide interest since the 1980s because of their reported enhanced growth, meat content, flavor and disease resistance.

Health & Welfare

Catfish genetics and breeding

Although research on catfish genetics began in the 1960s, applications of genetic improvement in aquaculture have lagged behind other animal industries.

About The Advocate

The Responsible Seafood Advocate supports the Global Seafood Alliance's (GSA) mission to advance responsible seafood practices through education, advocacy and third-party assurances.

Learn More

Search Responsible Seafood Advocate	Sooreb	Search Q	
Search Responsible Searoou Auvocale	Search	Search 🔍	



Listen to the seafood industry's top podcast

Advertising Opportunities

2022 Media & Events Kit

Categories

Aquafeeds > Health & Welfare Health & Welfare > From Our Sponsors > Innovation & Investment > Intelligence > Responsibility > Fisheries > Artículos en Español >

Don't Miss an Article

Featured

- Health & Welfare <u>An update on vibriosis, the major bacterial disease shrimp farmers face</u>
- Uncategorized <u>A seat at the table: Fed By Blue team says aquaculture needs a stronger voice</u>
- <u>Responsibility Quantifying habitat provisioning at macroalgae cultivation locations</u>

Popular Tags

6/13/2024

All Tags

Recent

- Fisheries Second Test: Another filler for the fisheries category
- Fisheries Test: This is filler for the fisheries Category
- <u>Aquafeeds Test Article</u>

×

- <u>Responsibility Study: Climate change will shuffle marine ecosystems in unexpected ways as ocean</u> <u>temperature warms</u>
- Health & Welfare Indian shrimp researchers earn a patent for WSSV diagnostic tool



Listen to the seafood industry's top podcast

- <u>About</u>
- <u>Membership</u>
- <u>Resources</u>
- <u>Best Aquaculture Practices (BAP)</u>
- <u>Best Seafood Practices (BSP)</u>
- GOAL Events
- <u>Advocate Magazine</u>
- <u>Aquademia Podcast</u>
- <u>Blog</u>
- <u>Contact</u>

Stay up to date with GSA

- Ø f
- 🕅
- in

Copyright © 2024 Global Seafood Alliance All rights reserved. <u>Privacy</u> Terms of Use <u>Glossary</u>