



- [GOAL Events](#)
- [Advocate Magazine](#)
- [Aquademia Podcast](#)
- [Blog](#)
- [Contact](#)
-
-
-
-
-
- [Log In](#)



- [About](#)
 - [Who We Are](#)
 - [Our History](#)
 - [Our Team](#)
 - [Sustainable Development Goals](#)
 - [Careers](#)
- [Membership](#)
 - [Overview](#)
 - [Our Members](#)
 - [Corporate Membership](#)
- [Resources](#)
- [Certification](#)
 - [Best Aquaculture Practices](#)
 - [Best Seafood Practices](#)

Search...



[Log In](#)

- [About](#)
 - [Who We Are](#)
 - [Our History](#)
 - [Our Team](#)
 - [Sustainable Development Goals](#)
 - [Careers](#)
- [Membership](#)
 - [Overview](#)
 - [Our Members](#)
 - [Corporate Membership](#)
- [Resources](#)
- [Certification](#)
 - [Best Aquaculture Practices](#)
 - [Best Seafood Practices](#)
- [GOAL Events](#)
- [Advocate Magazine](#)
- [Aquademia Podcast](#)
- [Blog](#)
- [Contact](#)



Post-harvest treatments defeat Vibrio vulnificus



1 May 2007 George J. Flick, Jr., Ph.D.



Bacteria account for 95 percent of all U.S. seafood-related deaths



Vibrio contamination of mussels and other shellfish can be effectively eliminated through several varied treatments.

Vibrio vulnificus is currently considered the most infectious and lethal of all human pathogenic vibrios. In the United States, it accounts for 95 percent of all seafood-related deaths. The organism is naturally ubiquitous in marine coastal waters and shellfish, and requires at least 0.5 percent sodium chloride for growth.

With fatality rates as high as 60 percent, septicemic infections caused by the organism usually result from the consumption of raw shellfish. The incubation period varies from several hours to several days, with a median of 26 hours. The most frequent symptoms include fever, chills, nausea, hypotension, and endotoxic shock. Liver damage and cirrhosis arising from chronic alcoholism, chronic renal disease, diabetes, and immunocompromising diseases are considered major factors in susceptibility. Secondary necrotic lesions of the extremities often necessitate surgical debridement or limb amputation.

Levels of *V. vulnificus* decrease in oysters stored at temperatures ranging 0.5 to 22 degrees-C for 10 days, which suggests that temperature abuse of oysters may not be a factor in increasing the public health risk of *V. vulnificus* through raw oyster consumption. However, the data also suggests that even with proper temperature control, indigenous levels of *V. vulnificus* can remain sufficiently high in oysters to produce infection in compromised individuals.

Presence in shellfish

Although *V. vulnificus* is widely distributed, its density can vary widely. In a 1998-1999 study, for example, shell oysters (*Crassostrea virginica*) were sampled at 275 restaurants or seafood markets throughout the United States. The oysters were harvested from the Gulf of Mexico, Pacific, Mid-Atlantic, and North Atlantic coasts of the U.S. and Canada.

V. vulnificus densities in oysters harvested during summer months from the Gulf coast and mid-Atlantic states sometimes exceeded 10,000 most probable number (MPN) per gram. The majority (78 percent) of lots harvested on the North Atlantic, Pacific and Canadian coasts contained densities below the detectable level of 0.2 MPN per gram. None exceeded 100.0 MPN per gram.

In analyses of edible bivalves at sites on the coast of Valencia, Spain, only two samples were confirmed as *V. vulnificus* because the organism can not grow in the high salinity of the Mediterranean Sea or compete with other predominant *Vibrio* species. In Italy, a survey of 600 mussel samples obtained from retailers in the Puglia region found *V. vulnificus* in 2.8 percent of the samples.

Along the southern coast of Brazil, *Crassostrea gigas* oysters were found to contain *V. vulnificus* organisms ranging 3 to 30 MPN per gram in 12 percent of the samples, with 25 isolates pathogenic. Seasonal sampling of oysters at two sites in the tropical waters of India detected the vibrio in 85 percent of the samples. Counts ranged from 10 per gram during the summer months to 1,000 per gram in the monsoon season. The density was controlled more by salinity than the relatively constant water temperature above 25 degrees-C.

Oysters from the U.S. Gulf of Mexico exhibited a similar seasonal distribution of *V. vulnificus*. The median concentration of 2,300 MPN/g of oyster meat from May through October was followed by a gradual reduction during November and December to below 10 per gram, where it remained from January through mid-March. A sharp increase to much higher summer levels in late March and April resulted with water temperatures up to 26 degrees-C.

In a year-long study of shrimp imported to Denmark, the microorganism was detected in three of 46 frozen raw shrimp samples. However, the organism was not recovered from any of 61 frozen cooked products tested, suggesting that the products were adequately heat treated and cross-contamination did not occur during processing. Imported shrimp products should not constitute a hazard to public health, provided the shrimp are consumed with adequate cooking and headed with sufficient care to prevent puncture wounds that could provide an opportunity for infection.

Processing effects

Tests have shown that postharvest treatments such as low-temperature pasteurization, rapid chilling, freezing, high-pressure processing, irradiation, and heat shock are capable of achieving reductions of *V. vulnificus*. However, most of these processes require either significant monetary investments or major efforts in personnel training. In addition, oysters are often killed during the process.

High pressure

A 6-log (99.9999 percent) reduction of *V. vulnificus* in phosphate-buffered saline (PBS) at 241 megapascals (mpa) required five minutes, which included a three-minute pressure come-up time. More than a 5.4-log reduction in *C. virginica* oysters at 345 mpa occurred during a six-minute pressure come-up time. The organism in both PBS and oysters was reduced to nondetectable levels at 586 mpa during a seven-minute pressure come-up time. In addition to reducing the pathogen, the oysters had an extended shelf life and required reduced hand-shucking time.

Electrolyzed oxidizing water

In another study, Pacific oysters (*Crassostrea gigas*) were inoculated with a five-strain cocktail of *V. vulnificus* at levels of 10^6 and treated with electrolyzed oxidizing water containing 1 percent sodium chloride at room temperature. Reductions were determined at two, four, six, and eight hours. A significant 1.05-log reduction of the organism occurred between four and six hours. Treatments exceeding six hours should be avoided to prevent the oysters' death.

Ice immersion

Unwashed *C. virginica* oysters were rapidly chilled by immersion in ice for three hours and examined for their total plate count, fecal coliform, and *V. vulnificus* contents several times during a seven-day post-treatment, refrigerated storage period at 7.2 degrees-C. The numbers of *V. vulnificus* generally declined when compared to the controls, but some increases in total heterotrophic bacteria and fecal coliforms were observed in the treated samples. The study did not support the use of ice immersion as a postharvest treatment because of the relatively small declines in the pathogen, usually less than 1 log (90 percent).

Irradiation

Live oysters, *C. virginica*, with naturally and artificially inoculated pathogenic *V. vulnificus* were exposed to 0-3 kGy of gamma irradiation. The organism was reduced from 10^6 colony-forming units per gram to nondetectable levels below 3 MPN per gram with 0.75-1.0 kGy irradiation exposure. Sensory panels indicated the participants were unable to distinguish the irradiated oysters from nonirradiated oysters.

Heat shock process

Further work with *C. virginica* oysters harvested from the Gulf of Mexico containing 10^2 - 10^4 MPN per gram of *V. vulnificus* were subjected to a heat-shock process of one to four minutes at internal meat temperatures ranging 45 to 53 degrees-C. *V. vulnificus* and total bacterial levels were significantly reduced 1-4 log in the finished product.

If the temperature was elevated to 50 degrees-C for 10 minutes, the levels were reduced to non-detectable, with the greatest reductions in oysters with the highest initial levels. The process, however, was not able to eliminate the organism in oysters with MPNs greater than 10^3 . The time required to reduce the microbial population by one decimal at 47 degrees-C for 52 strains averaged 78 seconds.

Cold storage

The time required for normal levels of *V. vulnificus* in *C. virginica* oysters from the Gulf of Mexico stored at 0 and 4 degrees C to reach undetectable levels normally exceeded the usual storage life of 14 days for shucked oyster meats and 21 days for shellstock oysters. Oysters stored at abuse temperatures of 7, 13, and 21 degrees-C over a 10-day period had *V. vulnificus* populations reaching 10^5 - 10^6 /g, a 2- to 4-log increase.

Freezing and storing the oysters for 12 weeks at minus-20 degrees-C did not result in complete destruction of the organism. However, other studies have shown that storage periods extended past 12 weeks result in a continuing decrease in *V. vulnificus* populations.

(Editor's Note: This article was originally published in the May/June 2007 print edition of the Global Aquaculture Advocate.)

Now that you've finished reading the article ...

... we hope you'll consider supporting our mission to document the evolution of the global aquaculture industry and share our vast network of contributors' expansive knowledge every week.

By becoming a Global Seafood Alliance member, you're ensuring that all of the pre-competitive work we do through member benefits, resources and events can continue. Individual membership costs just \$50 a year.

Not a GSA member? Join us.

[Support GSA and Become a Member](#)

Author

George J. Flick, Jr., Ph.D.

Food Science and Technology Department
Virginia Tech/Virginia Sea Grant (0418)
Blacksburg, Virginia 24061 USA

[117,100,101,46,116,118,64,103,107,99,105,108,102]

Share

- [✉ Share via Email](#)
- [🐦 Share on Twitter](#)
- [f Share on Facebook](#)
- [in Share on LinkedIn](#)

Tagged With

[shellfish](#) [Vibrio vulnificus](#) [George J. Flick](#)

Related Posts**Intelligence****[As ocean temperatures rise, so too will vibrio outbreaks](#)**

A study using a half-century of data has linked climate change and warming sea temperatures with an increase in illnesses from the common vibrio bacteria. Shellfish growers, fighting a particularly virulent strain of Vibrio parahaemolyticus, are changing their harvest protocols.

Health & Welfare

[Bacterial, chemical residues impact tilapia quality](#)

Both bacterial pathogens and chemical residues can affect the quality of farmed tilapia products. In production systems, Vibrio bacteria quickly proliferate following disease outbreaks.

Health & Welfare

[AHPN inferences based on behavior of vibrio bacteria](#)

Vibrio parahaemolyticus, a strain of which is the cause of acute hepatopancreatic necrosis (AHPN), has both virulent and benign strains. This strain colonizes the stomachs of shrimp by the formation of a biofilm, which protects it from antibiotics and other potential treatments.

Health & Welfare

[Beneficial microbes and pathogen control](#)

A range of alternative products is available to improve animal health and water quality, and control pathogen loads. Beneficial microbes produce antimicrobial compounds and suppress pathogen proliferation.


About The Advocate

The Responsible Seafood Advocate supports the Global Seafood Alliance's (GSA) mission to advance responsible seafood practices through education, advocacy and third-party assurances.

[Learn More](#)

Search Responsible Seafood Advocate

Search

Search 

Advertising Opportunities

[2022 Media & Events Kit](#)

Categories

[Aquafeeds](#) > [Health & Welfare](#) > [From Our Sponsors](#) > [Innovation & Investment](#) > [Intelligence Intelligence](#) > [Responsibility](#) > [Fisheries](#) > [Artículos en Español](#) >

Don't Miss an Article

Featured

- [Health & Welfare](#) [An update on vibriosis, the major bacterial disease shrimp farmers face](#)
- [Intelligence](#) [A seat at the table: Fed By Blue team says aquaculture needs a stronger voice](#)
- [Responsibility](#) [Quantifying habitat provisioning at macroalgae cultivation locations](#)

Popular Tags

All Tags








Recent

- [Fisheries](#) [Second Test: Another filler for the fisheries category.](#)
- [Fisheries](#) [Test: This is filler for the fisheries Category.](#)
- [Aquafeeds](#) [Test Article](#)
- [Responsibility](#) [Study: Climate change will shuffle marine ecosystems in unexpected ways as ocean temperature warms](#)
- [Health & Welfare](#) [Indian shrimp researchers earn a patent for WSSV diagnostic tool](#)



- [About](#)
- [Membership](#)
- [Resources](#)
- [Best Aquaculture Practices \(BAP\)](#)
- [Best Seafood Practices \(BSP\)](#)
- [GOAL Events](#)
- [Advocate Magazine](#)
- [Aquademia Podcast](#)
- [Blog](#)
- [Contact](#)

Stay up to date with GSA

- 
- 
- 
- 
- 

Copyright © 2024 Global Seafood Alliance
All rights reserved.

[Privacy](#)

[Terms of Use](#)

[Glossary](#)