



- [GOAL Events](#)
- [Advocate Magazine](#)
- [Aquademia Podcast](#)
- [Blog](#)
- [Contact](#)
-
-
-
-
-
- [Log In](#)



- [About](#)
 - [Who We Are](#)
 - [Our History](#)
 - [Our Team](#)
 - [Sustainable Development Goals](#)
 - [Careers](#)
- [Membership](#)
 - [Overview](#)
 - [Our Members](#)
 - [Corporate Membership](#)
- [Resources](#)
- [Certification](#)
 - [Best Aquaculture Practices](#)
 - [Best Seafood Practices](#)

Search...



[Log In](#)

- [About](#)
 - [Who We Are](#)
 - [Our History](#)
 - [Our Team](#)
 - [Sustainable Development Goals](#)
 - [Careers](#)
- [Membership](#)
 - [Overview](#)
 - [Our Members](#)
 - [Corporate Membership](#)
- [Resources](#)
- [Certification](#)
 - [Best Aquaculture Practices](#)
 - [Best Seafood Practices](#)
- [GOAL Events](#)
- [Advocate Magazine](#)
- [Aquademia Podcast](#)
- [Blog](#)
- [Contact](#)



Health & Welfare
Health & Welfare


Postlarvae evaluation key to controlling shrimp diseases



2 September 2013 Dr. Chalor Limsuwan Dr. Carlos A. Ching



Producers must adhere to standard criteria for stocking postlarvae



The three rostrum spines and small spine bud in the left photo identify a P.L.₁₀ shrimp. Each completely formed spine in the rostrum of white shrimp postlarvae represents three larval stages (right).

Serious economic losses reaching billions of dollars have been caused by viral and bacterial diseases in shrimp farming during the last few years. The quality of postlarvae has become one of the most important aspects in controlling these diseases. Shrimp farmers should apply a series of standard criteria in evaluating postlarvae to be stocked in their ponds.

Stocking size

The recommended size of shrimp for direct stocking into grow-out ponds is postlarvae that have been in the postlarvae stage for 10 days (P.L.₁₀), because complete gill development is attained at this stage, making the shrimp capable of withstanding transportation, acclimation and stocking at the farm. However, at farms with salinity below 5 ppt, P.L.₁₂ is the appropriate stage for stocking.

A practical way to recognize white shrimp postlarval stages is by looking at the number of spines in their rostrums. The P.L.₁₀ stage has three spines completely developed and a bud of the fourth spine developing in the rostrum, while P.L.₁₂ shrimp have four completely developed spines. On the other hand, weights and ages of shrimp postlarvae directly relate to their stages. So in the hatchery, normally developed larvae should have the following characteristics:

- Day 14 (P.L.₄₋₅) body weight $\leq 1,000$ P.L./1 g
- Day 16 (P.L.₇₋₈) body weight ≤ 700 P.L./1 g
- Day 18 (P.L.₁₀) body weight ≤ 300 P.L./1 g.

Another important factor for evaluating shrimp postlarvae in the hatchery is the nauplii stocking density, which should not exceed 100 nauplii/L. It is also important that the water temperature during hatchery culture is maintained at 30 ± 1 degrees-C for optimal larval development and quality.

Microbiological tests


The first criteria to take into account for evaluating shrimp postlarvae quality is to run a series of microbiological tests that will ensure pathogens are absent from the animals. Usually, the following tests are recommended during stage P.L.₆ to approve the postlarvae for stocking:

- Negative polymerase chain reaction (PCR) for infectious hypodermal and hematopoietic necrosis virus, infectious myonecrosis virus, Taura syndrome virus, yellow head virus and white spot syndrome virus.
- Maximum total bacteria count of 1.0×10^3 CFU/g of larval macerate in agar, of which more than 90 percent of the colonies should be yellow.
- Negative presence of *Vibrio harveyi* (bioluminescent bacteria), which can be detected in agar.

If the postlarvae fail to pass any of the microbiological tests, they should be rejected for farm stocking.

Direct visual evaluation

After the postlarvae pass the microbiological tests and before transporting them to the farm, direct visual evaluation in the tanks and observations under a light microscope should be done at the hatchery. The following aspects for direct evaluation should be considered.



Healthy shrimp postlarvae have a large, dark hepatopancreas containing many lipid droplets.

Larvae activity. To make sure larval activity is normal, suspend aeration in the larvae tank for a couple of minutes to look for positive rheotaxis – swimming against the water current by the animals.


Hepatopancreas condition. A large, dark-colored hepatopancreas with a lot of lipid droplets should be observed under the light microscope. A small, whitish hepatopancreas with only a few lipid droplets indicates an infection is occurring.

Digestive tract. Peristaltic movement in the intestine and a muscle:gut ratio of 4:1 in the last abdominal segment are indications of good health.

Fouling and necrosis. Direct observation of postlarvae under the microscope may detect the presence of *Lagenidium* species fungi and/or ciliate protozoa (*Zoothamnium*, *Epistylis*, *Vorticella* species), which usually cause fouling of the postlarvae gills. The presence of these parasites might indicate poor water quality in the larval tank. Eventually, if this situation is not controlled, *Leucotrix* species filamentous bacteria may invade other tissues, causing tissue damage and mortality.

Transport conditions

Transportation of larvae is an important issue to keep in mind for keeping good larvae quality before stocking. Appropriate oxygen, temperature and food requirements relate to the time required for transportation from the hatchery to the farm. The following are recommended conditions for transportation:



A muscle:gut ratio of 4:1 (arrows) is an indication of healthy shrimp.

- Less than four hours of travel: maintain ambient temperature.
- Four to 12 hours of travel: maintain 24 to 28 degrees-C.
- More than 12 hours of travel: maintain 18 to 23 degrees-C.
- At all distances, a minimum dissolved-oxygen concentration of 5.0 mg/L must be kept.
- For each shrimp postlarva, 15 to 20 artemia nauplii should be available every four hours.

If the conditions above are not met during transportation, stress of the animals can occur, with an increment in the concentration of ammonia in the water and bacteria within the postlarvae. Eventually, this situation can cause mortality before arrival at the farm, in which case stocking of postlarvae should be suspended.

(Editor's Note: This article was originally published in the September/October 2013 print edition of the Global Aquaculture Advocate.)

Authors

-  Dr. Chalor Limsuwan

Dr. Chalor Limsuwan

Professor
Department of Fishery Biology
Kasetsart University
50 Phaholyothin Road Chatuchak
Bangkok 10900 Thailand

[104,116,46,99,97,46,117,107,64,99,116,110,115,105,102,102]

-  Dr. Carlos A. Ching

Dr. Carlos A. Ching

Aquaculture Manager
Nicovita – Alicorp SAA
Callao, Lima, Peru

Share

-  [Share via Email](#)
-  [Share on Twitter](#)
-  [Share on Facebook](#)
-  [Share on LinkedIn](#)

Tagged With

[PCR postlarvae](#) [shrimp diseases](#) [Carlos A. Ching](#) [Chalor Limsuwan](#) [stocking size](#)

Related Posts

Health & Welfare

[Acclimating shrimp postlarvae before pond stocking](#)

Shrimp postlarvae acclimation before stocking into the various growout systems (ponds, raceways, tanks) is a critical – and often overlooked, sometimes taken for granted – step in the shrimp culture process. Various water quality parameters should be changed slowly so that the young shrimp have the time to gradually adapt to the new conditions.

Aquafeeds

[Biofloc consumption by Pacific white shrimp postlarvae](#)

The stable isotopes technique with $\delta^{13}\text{C}$ and $\delta^{15}\text{N}$ can be used to determine the relevance of different food sources to shrimp feeding during the pre-nursery phase of *Litopenaeus vannamei* culture. During this trial, different types of commercial feed, microalgae, *Artemia* sp. nauplii and bioflocs were used as food sources.

Health & Welfare

[A comprehensive look at the Proficiency Test for farmed shrimp](#)

The University of Arizona Aquaculture Pathology Laboratory has carried out the Proficiency Test (PT) since 2005, with 300-plus diagnostic laboratories participating while improving their capabilities in the diagnosis of several shrimp pathogens.

Responsibility

[A look at various intensive shrimp farming systems in Asia](#)

The impact of diseases led some Asian shrimp farming countries to develop biofloc and recirculation aquaculture system (RAS) production technologies. Treating incoming water for culture operations and wastewater treatment are biosecurity measures for disease prevention and control.

About The Advocate

The Responsible Seafood Advocate supports the Global Seafood Alliance's (GSA) mission to advance responsible seafood practices through education, advocacy and third-party assurances.

[Learn More](#)

Search Responsible Seafood Advocate



Advertising Opportunities

[2022 Media & Events Kit](#)

Categories

[Aquafeeds](#) > [Health & Welfare](#) [Health & Welfare](#) > [From Our Sponsors](#) > [Innovation & Investment](#) > [Intelligence](#) > [Responsibility](#) > [Fisheries](#) > [Artículos en Español](#) >

Don't Miss an Article

Featured

- [Health & Welfare](#) [An update on vibriosis, the major bacterial disease shrimp farmers face](#)
- [Intelligence](#) [A seat at the table: Fed By Blue team says aquaculture needs a stronger voice](#)
- [Responsibility](#) [Quantifying habitat provisioning at macroalgae cultivation locations](#)

Popular Tags

All Tags ▼

Recent

- [Fisheries](#) [Second Test: Another filler for the fisheries category](#)
- [Fisheries](#) [Test: This is filler for the fisheries Category](#)
- [Aquafeeds](#) [Test Article](#)
- [Responsibility](#) [Study: Climate change will shuffle marine ecosystems in unexpected ways as ocean temperature warms](#)
- [Health & Welfare](#) [Indian shrimp researchers earn a patent for WSSV diagnostic tool](#)



- [About](#)
- [Membership](#)
- [Resources](#)
- [Best Aquaculture Practices \(BAP\)](#)
- [Best Seafood Practices \(BSP\)](#)
- [GOAL Events](#)
- [Advocate Magazine](#)
- [Aquademia Podcast](#)
- [Blog](#)
- [Contact](#)

Stay up to date with GSA

- 
- 
- 
- 
- 

Copyright © 2024 Global Seafood Alliance
All rights reserved.

[Privacy](#)

[Terms of Use](#)

[Glossary](#)