Global Seafood Alliance Logo

- GOAL Events
- Advocate Magazine
- Aquademia Podcast
- <u>Blog</u>
- Contact
- 0
- X
- in
- Log In

- About
 - Who We Are
 - Our History
 - o Our Team
 - Sustainable Development Goals
 - Careers
- <u>Membership</u>
 - o <u>Overview</u>
 - Our Members
 - Corporate Membership
- Resources
- Certification
 - Best Aquaculture Practices
 - Best Seafood Practices

Search...

Q

Log In

- About
 - Who We Are
 - o Our History
 - o Our Team
 - Sustainable Development Goals
 - Careers
- Membership
 - o <u>Overview</u>
 - o Our Members
 - Corporate Membership
- Resources
- Certification
 - Best Aquaculture Practices
 - Best Seafood Practices
- GOAL Events
- Advocate Magazine
- Aquademia Podcast
- **Blog**
- Contact



Prebiotics, probiotics provide alternatives to antibiotics

Responsible Seafood Advocate logo

1 July 2010 Brian C. Peterson, Ph.D. Natha J. Booth, Ph.D. Lester Khoo, VMD, Ph.D. Bruce B. Manning, Ph.D.



Some prebiotics show protection when injected but not when fed



Oligosaccharides fed to European sea bass and other species have shown varying degrees of success in improving growth efficiency and resistance to

As aquaculture continues to expand, pressures to find alternatives to medicinal disease treatments are also rising. Concerns over antibiotic use in aquaculture include the development and transfer of antibiotic resistance to fish and human pathogens, as well as the potential for accumulation of residual antibiotics in aquaculture products.

As the demand for aquaculture products increases, so does the search for environmentally friendly alternatives to antibiotics. Alternatives to antibiotics include dietary prebiotics, probiotics and synbiotics.

Prebiotics and probiotics have been shown to enhance both growth efficiency and disease resistance in a variety of aquatic species, while synbiotics (mixtures of pre- and probiotics) have yet to be extensively evaluated. Research using prebiotics and probiotics in aquaculture is growing, but the results are often inconsistent, and the mechanisms through which they function are poorly understood.

Prebiotics

A prebiotic is a non-digestible substance that is thought to provide a beneficial effect on the host by stimulating growth or activity of naturally occurring bacteria in the gastrointestinal tract. Typically, prebiotics are non-digestible carbohydrates (fructo-oligosaccharides, galacto-oligosaccharides, mannanoligosaccharides), but can also include indigestible dietary fibers (dextrins, inulin, lignin, waxes, beta glucans).

Many forms of prebiotics have been fed to a wide variety of fish species with varying responses. For example, inulin has recently been shown to alter the intestinal microbial communities of turbot, Arctic char, Atlantic salmon and hybrid striped bass, but did not yield any benefits toward improving growth or feed efficiency.

Prebiotics like glucans have shown protection against enteric septicemia of catfish in channel catfish when the products are injected but not when fed. Furthermore, a prebiotic composed of brewer's yeast, dairy components and dried fermentation products was found to significantly increase feed efficiency and reduce the mortality of hybrid striped bass challenged with bacterial pathogens.

Non-digestible oligosaccharides fed to rainbow trout, European sea bass, channel catfish and hybrid striped bass have shown varying degrees of success in improving growth efficiency and resistance to disease. The results of these studies suggested that prebiotics have a role in aquaculture as a feed additive for improving growth and resistance to disease.




Fig. 1: The lower intestine of catfish fed a control diet showed microvilli lengths similar to those in the intestine of fish that received feed supplemented with the prebiotic mannanoligosaccharide.

Probiotics

Probiotics are live microorganisms thought to have a beneficial health effect on the host by adding "good" bacteria to the gut. Lactobacillus, Bacillus, Enterococcus, Carnobacterium, Saccharomyces and Candida are probiotic organisms widely investigated in aquaculture. These organisms are added directly to the water environment of the fish or administered orally in feed.

Probiotics have been shown to protect against bacterial infections by Vibrio, Aeromonas, Yersinia and Ichthyophthirius in rainbow trout. In African catfish, Lactobacillus enhanced fish health, survival and growth performance. In channel catfish, Pediococcus and Entebacteria had no growth-promoting or immune-stimulating effect, while Bacillus bacteria were shown to increase the survival of pond-raised catfish.

Synbiotics

Another class of compounds called synbiotics includes mixtures of prebiotics and probiotics. Synbiotics may work by stimulating the growth of beneficial bacteria within the gastrointestinal tract of the host. The authors are not aware of any studies in aquatic animals that have evaluated the efficacy of these products, but synbiotics have potential for manipulating the microflora of the gut and improving growth and disease resistance.

MOS research

At the Thad Cochran National Warmwater Aquaculture Center in Stoneville, Miss., USA, the authors examined a commercial mannan-oligosaccharide (MOS) available from Alltech, Inc. derived from the cell wall of yeast as a potential feed additive in channel catfish.

In one study, the catfish were fed the prebiotic at 2 grams per kg diet for six weeks, then challenged with *Edwardsiella ictaluri*, the bacterium that causes enteric septicemia of catfish. All fish were bled before the challenge as well as on day 14 of the challenge. In a second study, catfish received the prebiotic at 4 grams per kg diet for nine weeks, followed by challenge with E. ictaluri. Mortalities were collected for 21 days.

Gut morphology was examined in fish fed diets with mannan-oligosaccharide, as well as controls prior to challenge. Sections of the upper, middle and lower parts of the intestine were taken for analysis.

Results

Feeding MOS at 2 or 4 grams per kg diet for six or nine weeks did not improve weight gain, specific growth rate or feed conversion in the fish (Table 1). However, survival was significantly increased after challenge with *E. ictaluri* in both studies.

Peterson, Culture performance of channel catfish, Table 1

Treatment	Weight Gain (g)	* Specific Growth Rat	e Feed-Conversion Ratio	Survival (%)
Study I				
Control	59.5	2.00	1.40	52.5 ^y
MOS	54.9	1.90	1.49	90.0 ^z
Standard error	r 4.0	0.10	0.10	5.8
Study II				
Control	40.6	2.60	1.32	45.0 ^y
MOS	44.4	2.90	1.33	70.0 ^z
Standard error	r 1.4	0.01	0.10	1.7

Table 1. Culture performance of channel catfish fed an MOS -supplemented diet or control diet. Values with different letters within columns are significantly different (P < 0.05).

Plasma levels of lysozyme, an indicator of a non-specific immune response, were similar in fish fed MOS feed before and after challenge. Fig. 1 shows representative lower intestine sections of fish fed MOS compared to controls. Microvilli lengths in the lower, middle and proximal intestine were similar after nine weeks of feeding.

(Editor's Note: This article was originally published in the July/August 2010 print edition of the Global Aquaculture Advocate.)

Now that you've finished reading the article ...

... we hope you'll consider supporting our mission to document the evolution of the global aquaculture industry and share our yast network of contributors' expansive knowledge every week.

By becoming a Global Seafood Alliance member, you're ensuring that all of the pre-competitive work we do through member benefits, resources and events can continue. Individual membership costs just \$50 a year.

Not a GSA member? Join us.

Support GSA and Become a Member

Authors

• Brian C. Peterson, Ph.D.

Brian C. Peterson, Ph.D.

Thad Cochran National Warmwater Aquaculture Center United States Department of Agriculture Agricultural Research Service Stoneville, Mississippi 38776 USA

[118,111,103,46,97,100,115,117,46,115,114,97,64,110,111,115,114,101,116,101,112,46,110,97,105,114,98]

Natha J. Booth, Ph.D.

Natha J. Booth, Ph.D.

Thad Cochran National Warmwater Aquaculture Center United States Department of Agriculture Agricultural Research Service Stoneville, Mississippi 38776 USA

Lester Khoo, VMD, Ph.D.

Lester Khoo, VMD, Ph.D.

Thad Cochran National Warmwater Aquaculture Center United States Department of Agriculture Agricultural Research Service Stoneville, Mississippi 38776 USA

• Bruce B. Manning, Ph.D.

Bruce B. Manning, Ph.D.

Thad Cochran National Warmwater Aquaculture Center United States Department of Agriculture Agricultural Research Service Stoneville, Mississippi 38776 USA

Share

- Share via Email
- **Share on Twitter**
- f Share on Facebook
- in Share on LinkedIn

Tagged With

antibiotics probiotics prebiotics Brian C. Peterson Natha J. Booth Lester Khoo Bruce B. Manning

Related Posts

Health & Welfare

Prebiotics and their role in sustainable aquaculture

Results of a study to evaluate a commercial prebiotic on growth of European sea bass juveniles indicates this product can improve fish growth performance and allowing for the fish to reach the marketable size earlier, boosting their immune status, and reducing potential fish losses due to pathogens.

Health & Welfare

Probiotics, prebiotics in aquatic animals

Research has shown that probiotics and prebiotics can help mediate stress responses and improve disease resistance, growth performance, feed utilization, carcass composition and other traits by stimulating animals' innate immune systems.

Aquafeeds

Biosecurity protocols needed for shrimp feeds, feeding practices

Shrimp aquafeeds – live, fresh or formulated – should not be an entry point of potential pathogens to the shrimp and/or to their culture systems.

Aquafeeds

Evaluating commercial, dietary pre-, pro- and synbiotics in hybrid catfish

This study evaluated a commercially available prebiotics and probiotics under conditions simulating commercial production in hybrid catfish. Results showed that the various dietary treatments did not have any significant effects on processing characteristics or composition.

About The Advocate

The Responsible Seafood Advocate supports the Global Seafood Alliance's (GSA) mission to advance responsible seafood practices through education, advocacy and third-party assurances.

Learn More

Search Responsible Seafood Advocate Search Search Search



Advertising Opportunities

2022 Media & Events Kit

Categories

Aquafeeds > Health & Welfare Health & Welfare > From Our Sponsors > Innovation & Investment > Intelligence > Responsibility > Fisheries > Artículos en Español >

Don't Miss an Article

Featured

- Health & Welfare An update on vibriosis, the major bacterial disease shrimp farmers face
- Uncategorized A seat at the table: Fed By Blue team says aquaculture needs a stronger voice
- Responsibility Quantifying habitat provisioning at macroalgae cultivation locations

Popular Tags



Recent

- Fisheries Second Test: Another filler for the fisheries category
- Fisheries Test: This is filler for the fisheries Category
- Aquafeeds Test Article
- · Responsibility Study: Climate change will shuffle marine ecosystems in unexpected ways as ocean temperature warms
- Health & Welfare Indian shrimp researchers earn a patent for WSSV diagnostic tool



- **About**
- <u>Membership</u>
- Resources
- Best Aquaculture Practices (BAP)
- Best Seafood Practices (BSP)
- GOAL Events
- Advocate Magazine
- Aquademia Podcast
- Blog
- Contact

Stay up to date with GSA

- 0

Copyright © 2024 Global Seafood Alliance All rights reserved.

Privacy Terms of Use <u>Glossary</u>