#### Global Seafood Alliance Logo

- GOAL Events
- <u>Advocate Magazine</u>
- Aquademia Podcast
- <u>Blog</u>
- <u>Contact</u>
- 0
- f
- 🗙
- in
- 🕨
- <u>Log In</u>
- $\square$ 
  - <u>About</u>
    - Who We Are
    - <u>Our History</u>
    - <u>Our Team</u>
    - Sustainable Development Goals
    - <u>Careers</u>
  - <u>Membership</u>
    - <u>Overview</u>
    - <u>Our Members</u>
    - Corporate Membership
  - <u>Resources</u>
  - <u>Certification</u>
    - Best Aquaculture Practices
    - Best Seafood Practices

Search...

Q

<u>Log In</u>

- <u>About</u>
  - Who We Are
  - <u>Our History</u>
  - <u>Our Team</u>
  - Sustainable Development Goals
  - <u>Careers</u>
- <u>Membership</u>
  - <u>Overview</u>
  - Our Members
  - Corporate Membership
- <u>Resources</u>
- <u>Certification</u>
  - Best Aquaculture Practices
  - Best Seafood Practices
- GOAL Events
- <u>Advocate Magazine</u>

- <u>Aquademia Podcast</u>
- <u>Blog</u>
- Contact



Health & Welfare Health & Welfare

## **Red abalone morphology**

Responsible Seafood Advocate logo 1 November 2007 Miguel Á. del Río-Portilla Alberto Miranda-Velasco

## Implications for breeding programs



Red abalone breeding programs need to consider total weight and size at first maturity in broodstock selection.

In Mexico, it takes three to four years of culture for an abalone to reach market size of 80-cm shell length and 80gram total weight. Commercial breeding and selection programs that seek to reduce this grow-out time must consider exploiting the biological potential of the species, including whether abalone morphology changes as abalone grow. The use of allometry has been of great value in understanding the way animal shape changes during growth in many species.

## Allometry

Allometry is the study of size and its implications. Within a species, individuals differ in such features as body size, age at first reproduction, and the effort they put into growth and reproduction. When two variables in an organism increase proportionally, they are considered isometric, but when the increase in one of the variables is smaller (negative) or higher (positive) than the other, they are allometric. To determine whether a pair of variables is isometric or allometric, the allometric equation  $Y = a X^b$  is used, where X and Y are size-related measures, and a and b are empirical constants.

To determine whether a two-variable relationship – like shell length and width, for example – is isometric or allometric, it is necessary to test the value of b and find out whether it agrees with the theoretical isometric value: a) one for linear vs. linear units (cm vs. cm) or b) three for volume vs. linear units (g vs. cm).

# Abalone morphology study

At two commercial abalone farms in Baja California, Mexico, the authors determined morphological relationships in red abalone (*Haliotis rufescens*) that ranged 12.0 to 84.4 cm in shell length. All variables were plotted against shell length and constants A and B were statistically tested to determine the isometric per allometric relationships between the variables and shell length. The animals' sex (male, female, or undifferentiated) was noted to determine the size at which abalones matured. Table 1 shows the allometric equation parameter estimates determined.


## del Rio-Portilla, Allometric equation parameter estimation against shell length for red abalones, Table 1

	Width	Height	Meat Weigh	t Shell Weigh	t Total Weight
Abulones Cultivados Farm					
n	65	65	65	65	65
a	0.7015				
b	1.0056				
r <sup>2</sup>	0.9847				
	Isometrie	c - Allometri	c Isometric	- Allometric	- Allometric
Productores Marinos Baja Farn	ı				
n	52	52	52	52	52
a	0.7193	0.4379	0.0001	0.0001	0.0002
b	0.9913	0.7889	3.0482	2.7868	2.9286
r <sup>2</sup>	0.9918	0.9744	0.9909	0.9947	0.9948
	Isometric - Allometric Isometric			- Allometric	- Allometric

Table 1. Allometric equation parameter estimation against shell length for red abalones grown at two farms in Mexico.

Growing conditions, including the quality and quantity of available food, affect abalone growth and meat content. Under the good growing conditions at the farms, meat weight was isometric.

# **Practical implications**



Natural negative allometry in red abalone shell length could be useful for abalone growers, because abalones may have a heavier shell when small that then becomes lighter as the animals grow, thus increasing meat yield. This also represents an advantage for abalone handling, because small abalones are proportionally less sensitive to handling than larger abalones.

Because it is not possible to measure abalone meat weight without sacrificing the animals, the authors suggested, based on the study, that red abalone breeding programs should use either family-sibling selection to estimate meat weight, or consider those abalones with a total weight higher than the expected value obtained with the equation Total Weight = 0.0002 Shell Length<sup>2.947</sup>. This equation was obtained when abalones from both farms were pooled together. The authors also found no difference in shape and trend among undifferentiated male and female abalones, which is the case in other gastropods.

## Size at maturity

Another consideration for abalone-breeding programs is size at maturity. Red abalones mature at a relatively small size of 50- to 55-mm shell length and 20- to 25-gram total weight, while commercial size is around 80-mm shell length. This has important implications for abalone culture, because growth rates decrease when maturity is reached. Therefore, abalone-breeding programs should also consider size at first maturity for breeder selection.

The use of triploids has been proposed as an alternative for abalone production, because triploid animals, in general, do not use energy in gamete production and thus have more energy for growth. It would, however, be necessary to determine whether the weight per shell length relationship for these animals remains isometric or allometric.

## Outlook

Due to high abalone fecundity, it is a common hatchery practice to use small numbers of broodstock. However, any abalone selective-breeding program should consider a large number of families. Thus, more studies on heritability and correlation between traits need to be carried out. The authors believe red abalone-breeding programs need to consider total weight rather than shell length when broodstock animals are selected, because of their allometric characteristics.

(*Editor's Note: This article was originally published in the November/December 2007 print edition of the* Global Aquaculture Advocate.)

## Now that you've finished reading the article ...

... we hope you'll consider supporting our mission to document the evolution of the global aquaculture industry and share our vast network of contributors' expansive knowledge every week.

By becoming a Global Seafood Alliance member, you're ensuring that all of the pre-competitive work we do through member benefits, resources and events can continue. Individual membership costs just \$50 a year.

#### Not a GSA member? Join us.

Support GSA and Become a Member

#### Authors

• 📄 Miguel Á. del Río-Portilla

Miguel Á. del Río-Portilla

Laboratorio de Genética Departamento de Acuicultura CICESE Km 107 Carretera Tijuana-Ensenada Ensenada, Baja California, Mexico 22800

[120, 109, 46, 101, 115, 101, 99, 105, 99, 64, 111, 105, 114, 108, 101, 100, 109]

Alberto Miranda-Velasco

Alberto Miranda-Velasco

Laboratorio de Genética Departamento de Acuicultura CICESE Km 107 Carretera Tijuana-Ensenada Ensenada, Baja California, Mexico 22800

#### Share

- Share via Email
- **J** Share on Twitter
- **f** Share on Facebook
- in <u>Share on LinkedIn</u>

#### **Tagged With**

Red abalone Miguel Á. del Río-Portilla Alberto Miranda-Velasco

#### **Related Posts**

Health & Welfare

## <u>Algae shows promise as alternative DHA source in rainbow trout diets</u>

A growth trial in Canada evaluated the use of algae biomass to increase the concentration of long-chain polyunsaturated fatty acids in the tissues of rainbow trout.

#### Intelligence

## An emerging shellfish farming industry in Namibia

For shellfish farming in Namibia to continue expanding, industry must better comply with approved sanitation standards. The Namibian Shellfish Monitoring and Sanitation Program, currently in development, will help.

#### Innovation & Investment

### **Aquaculture Exchange: George S. Lockwood**

With his book, "Aquaculture: Will it Rise to Its Potential to Feed the World?" hot off the presses, the pioneer abalone farmer vents on U.S. aquaculture regulations but remains deeply optimistic about fish farming.

#### Aquafeeds

## Aquaculture Exchange: Giovanni Turchini, Deakin University, part 1

One of the world's leading fish nutrition experts talks about how aquaculture can learn to survive, and even thrive without depending on fishmeal and fish oil. It'll take a lot of innovation, but Giovanni Turchini is confident that the industry is on the right path.

#### About The Advocate

The Responsible Seafood Advocate supports the Global Seafood Alliance's (GSA) mission to advance responsible seafood practices through education, advocacy and third-party assurances.

#### Learn More

Search Responsible Seafood Advocate	Search	Search <b>Q</b>	
Search Responsible Searoou Auvocale	Search	Search	



# Listen to the seafood industry's top podcast

#### **Advertising Opportunities**

#### 2022 Media & Events Kit

#### Categories

Aquafeeds > Health & Welfare Health & Welfare > From Our Sponsors > Innovation & Investment > Intelligence > Responsibility > Fisheries > Artículos en Español >

#### Don't Miss an Article

#### Featured

- Health & Welfare An update on vibriosis, the major bacterial disease shrimp farmers face
- Uncategorized <u>A seat at the table: Fed By Blue team says aquaculture needs a stronger voice</u>
- <u>Responsibility Quantifying habitat provisioning at macroalgae cultivation locations</u>

#### **Popular Tags**

All Tags

#### Recent

- Fisheries Second Test: Another filler for the fisheries category
- Fisheries Test: This is filler for the fisheries Category
- <u>Aquafeeds Test Article</u>

×

- <u>Responsibility Study: Climate change will shuffle marine ecosystems in unexpected ways as ocean</u> <u>temperature warms</u>
- Health & Welfare Indian shrimp researchers earn a patent for WSSV diagnostic tool



# Listen to the seafood industry's top podcast

- <u>About</u>
- <u>Membership</u>
- <u>Resources</u>
- <u>Best Aquaculture Practices (BAP)</u>
- <u>Best Seafood Practices (BSP)</u>
- GOAL Events
- <u>Advocate Magazine</u>
- <u>Aquademia Podcast</u>
- <u>Blog</u>
- <u>Contact</u>

#### Stay up to date with GSA

- 0 f
- X •
- in

Copyright © 2024 Global Seafood Alliance All rights reserved. Privacy Terms of Use <u>Glossary</u>