



- [GOAL Events](#)
- [Advocate Magazine](#)
- [Aquademia Podcast](#)
- [Blog](#)
- [Contact](#)
-
-
-
-
-
- [Log In](#)



- [About](#)
 - [Who We Are](#)
 - [Our History](#)
 - [Our Team](#)
 - [Sustainable Development Goals](#)
 - [Careers](#)
- [Membership](#)
 - [Overview](#)
 - [Our Members](#)
 - [Corporate Membership](#)
- [Resources](#)
- [Certification](#)
 - [Best Aquaculture Practices](#)
 - [Best Seafood Practices](#)

Search...



[Log In](#)

- [About](#)
 - [Who We Are](#)
 - [Our History](#)
 - [Our Team](#)
 - [Sustainable Development Goals](#)
 - [Careers](#)
- [Membership](#)
 - [Overview](#)
 - [Our Members](#)
 - [Corporate Membership](#)
- [Resources](#)
- [Certification](#)
 - [Best Aquaculture Practices](#)
 - [Best Seafood Practices](#)
- [GOAL Events](#)
- [Advocate Magazine](#)
- [Aquademia Podcast](#)
- [Blog](#)
- [Contact](#)



Health & Welfare

Health & Welfare


Rotifer enrichment: Concept simplifies larval fish feeding



2 September 2013 Eamonn O'Brien Philippe Dhert



Standalone culture diet significantly reduces requirement to heat water



Establishing a standalone culture diet makes rotifers immediately available at any time during the culture cycle.

For the last quarter of a century, hatcheries have tended to apply the same labor-intensive batch culture system of rotifer production. This process, which usually involves three- or four-day batches, follows a protracted sequence of events in which rotifers are harvested, rinsed and put into an enrichment tank for a number of hours. After this, they are harvested again, then cleaned and finally fed to larval fish.

The more rotifers are handled, the greater the risk of damaging them becomes. Furthermore, fish are normally fed rotifers three to four times a day, which often requires repetitive separate rotifer harvesting and enrichment processes.

Rotifer enrichment trials

In order to verify the hypothesis that a well-formulated live food diet could avoid the use of an enrichment step, the authors conducted trials in which an existing algae-based rotifer culture diet was modified to slowly release the enrichment components, such as protein, fat, vitamins and minerals. The presence of nutrients at metabolizing concentrations allowed not only gut tissue filling but also gradually enriched the tissue of the rotifers without consuming excessive oxygen or eutrophication, which is a well-known problem under conventional enrichment conditions.

The experiment was conducted over several culture cycles to ensure the diet did not compromise the health or reproduction of the rotifers. Similar growth and reproduction figures were registered with the conventional rotifer diet and the standalone diet for a period of nine weeks, which corresponds with 15 rotifer cycles (Table 1). Feed-conversion ratios averaged 0.9 to 1.0, while egg ratios were 25 to 40 percent – similar to these for traditional diets.

O'Brien, Average growth, Table 1

Day	Tank Volume (L)	Density (rotifers/mL)	Rotifers (million)	Egg %	Feed (g/million)	Feed (g)
0	900	466	419.4	24	0.50	209.70
1	900	696	626.4	30	0.40	250.56
2	900	1,016	914.4	43	0.40	365.76
3	900	1,288	1,159.2	35	0.31	359.35
4	900	1,890	1,701.0	23	–	–

Table 1. Average growth of a batch rotifer culture.

With a slow enrichment over one or two days, the standalone diet was naturally enriched to 80 to 90 percent, about the same as following a conventional enrichment process. After one three- to four-day cycle, the enrichment seen with the standalone diet was 100 percent that of the conventional procedure.

Analyses of the protein, fat, vitamin and mineral compositions of the rotifers were compared to those obtained after an enrichment treatment. The kinetics of the enrichment during the standalone culture trial and traditional culture and enrichment procedures are presented in Figures 1 and 2. The natural algae rotifer diet gave a basic enrichment between 10-15 mg/g (dry weight) during the four-day culture cycle. An enrichment diet with added proteins, phospholipids and marine oils enriched rotifers in two to four hours, while the standalone diet provided immediate enrichment.




Fig. 1: Highly unsaturated fatty acids enrichment kinetics of rotifers.




Fig. 2: Long-term n-3 highly unsaturated fatty acids enrichment.

Varied advantages

The trials confirmed many gains can be had from establishing a standalone culture diet. Such a product makes rotifers immediately available to hatchery staff at any time during the culture cycle, and can result in significant labor and cost savings. Rotifers are enriched at 26 degrees-C, which requires hatcheries to heat a lot of water for both the enrichment and subsequent rinsing steps. A standalone culture diet significantly reduces that requirement and also simplifies the processes for hatchery managers so they can focus more on the quality aspects of cultivating rotifers.

Another benefit of using a standalone diet is it results in more stable rotifers. Since the rotifers incorporate the algae feed into their own biomass, the enrichment stays in the rotifers longer. Following the traditional enrichment process, rotifers are put into a very eutrophic environment filled with emulsions or fats. Removing the enrichment step all but eliminates the emulsion discharges into the environment during post-enrichment rinsing.

Final product

These findings and others led Skretting's Marine Hatchery Feeds division to develop ORI-ONE, a commercial algae-based powder for rotifer production. No yeast or emulsion is used, and a wide particle size range is available, which results in more efficient uptake. The particles also come in a wide range of shapes, which gives better adhesion and retention within the rotifers.

(Editor's Note: This article was originally published in the September/October 2013 print edition of the Global Aquaculture Advocate.)

Authors

-  Eamonn O'Brien

Eamonn O'Brien

Skretting
Akkerhage 4
9000 Gent, Belgium





[109,111,99,46,103,110,105,116,116,101,114,107,115,64,110,101,105,114,98,111,46,110,110,111,109,97,101]

-  Philippe Dhert

Philippe Dhert

Skretting
Akkerhage 4
9000 Gent, Belgium

Share

-  [Share via Email](#)
-  [Share on Twitter](#)
-  [Share on Facebook](#)
-  [Share on LinkedIn](#)

Tagged With

[algae rotifer](#) [Philippe Dhert](#) [Eamonn O'Brien](#) [enrichment](#) [rotifer harvesting](#)

Related Posts

[Aquafeeds](#)

[Rotifers and diatoms aid shrimp biofloc nurseries](#)

Larval shrimp performance (feed conversion and final weight) in a biofloc nursery system improved with both added diatoms (*Navicula* sp.) and rotifers (*Brachionus plicatilis*).

[Health & Welfare](#)

[New-generation diets omit rotifer enrichment, improve water quality](#)

An innovative new diet supports the mass culture of rotifers at the same time it transforms their nutritional composition to meet the requirements of first-feeding fish larvae. As in nature, the diet is based on a mixture of algae-based nutrients to which rotifers respond.

[Health & Welfare](#)

[Advances in intensive copepod production technology](#)

Research at the Oceanic Institute has been successful in overcoming bottlenecks associated with rearing small-mouthed fish larvae by finding a suitable first feed. Early work on the calanoid copepod *Parvocalanus crassirostris* focused on parameters necessary for successful maintenance of stock cultures.

[Responsibility](#)

[Assessing culture potential of red emperor snapper in New Caledonia](#)

The red emperor snapper, known as “pouatte” in New Caledonia, is valuable throughout its broad geographic range and a highly valued food fish locally. Declining wild catches and market demand have provided the incentive to carry out technical feasibility studies to determine its commercial aquaculture potential.

About The Advocate

The Responsible Seafood Advocate supports the Global Seafood Alliance’s (GSA) mission to advance responsible seafood practices through education, advocacy and third-party assurances.

[Learn More](#)

Search Responsible Seafood Advocate



Advertising Opportunities

[2022 Media & Events Kit](#)

Categories

[Aquafeeds](#) > [Health & Welfare](#) [Health & Welfare](#) > [From Our Sponsors](#) > [Innovation & Investment](#) > [Intelligence](#) > [Responsibility](#) > [Fisheries](#) > [Artículos en Español](#) >

Don't Miss an Article

Featured

- [Health & Welfare](#) [An update on vibriosis, the major bacterial disease shrimp farmers face](#)
- [Intelligence](#) [A seat at the table: Fed By Blue team says aquaculture needs a stronger voice](#)
- [Responsibility](#) [Quantifying habitat provisioning at macroalgae cultivation locations](#)

Popular Tags

All Tags

Recent

- Fisheries [Second Test: Another filler for the fisheries category.](#)
- Fisheries [Test: This is filler for the fisheries Category.](#)
- [Aquafeeds Test Article](#)
- [Responsibility Study: Climate change will shuffle marine ecosystems in unexpected ways as ocean temperature warms](#)
- Health & Welfare [Indian shrimp researchers earn a patent for WSSV diagnostic tool](#)



- [About](#)
- [Membership](#)
- [Resources](#)
- [Best Aquaculture Practices \(BAP\)](#)
- [Best Seafood Practices \(BSP\)](#)
- [GOAL Events](#)
- [Advocate Magazine](#)
- [Aquademia Podcast](#)
- [Blog](#)
- [Contact](#)

Stay up to date with GSA

-
-
-
-
-

Copyright © 2024 Global Seafood Alliance
All rights reserved.
[Privacy](#)
[Terms of Use](#)
[Glossary](#)