

 [Global Seafood Alliance Logo](#)

- [GOAL Events](#)
- [Advocate Magazine](#)
- [Aquademia Podcast](#)
- [Blog](#)
- [Contact](#)

- 
- 
- 
- 
- 

- [Log In](#)

—

- [About](#)
 - [Who We Are](#)
 - [Our History](#)
 - [Our Team](#)
 - [Sustainable Development Goals](#)
 - [Careers](#)
- [Membership](#)
 - [Overview](#)
 - [Our Members](#)
 - [Corporate Membership](#)
- [Resources](#)
- [Certification](#)
 - [Best Aquaculture Practices](#)
 - [Best Seafood Practices](#)

Search...



[Log In](#)

- [About](#)
 - [Who We Are](#)
 - [Our History](#)
 - [Our Team](#)
 - [Sustainable Development Goals](#)
 - [Careers](#)
- [Membership](#)
 - [Overview](#)
 - [Our Members](#)
 - [Corporate Membership](#)
- [Resources](#)
- [Certification](#)
 - [Best Aquaculture Practices](#)
 - [Best Seafood Practices](#)
- [GOAL Events](#)
- [Advocate Magazine](#)
- [Aquademia Podcast](#)
- [Blog](#)
- [Contact](#)



 Responsibility
Responsibility

Salinity in aquaculture, Part 2

 [Responsible Seafood Advocate logo](#)

4 November 2019 Claude E. Boyd, Ph.D.



Different methods to measure important environmental factor



In aquaculture settings, salinity can be measured adequately with a handheld salinity refractometer assuming the refractometer is performing correctly. Photo by Kandschwar [CC BY-SA 2.0 de (<https://creativecommons.org/licenses/by-sa/2.0/de/deed.en>)]

There are several ways of measuring common salinity to include electrical conductivity (specific conductance), density, chlorinity and refractive index. The purpose here is to describe the different methods of measuring the common salinity of water.

Electrical conductivity

The ability of water to conduct electricity increases with greater salinity because electricity is conducted through water by free ions. A conductivity meter (Fig. 1) is basically a Wheatstone bridge – traditionally used to measure resistivity – modified to measure the reciprocal of resistivity or conductivity. The unit of resistance traditionally was the ohm, but to avoid using 1/ohm for conductivity, the unit mho (ohm spelled backwards) was adopted. The unit siemen also is used as the unit for conductivity, and 1 micromho/cm is the same as 1 microsiemen/cm. Conductivity increases with greater temperature, but most modern conductivity meters are temperature compensated to read out for 25 degrees-C.




Fig. 1: A conductivity meter.

There is a linear relationship between conductivity and salinity. Seawater has a conductivity of about 50,000 mmhos/cm, and half-strength seawater (about 17.25 ppt) has a conductivity of around 25,000 mmhos/cm. Most portable conductivity meters have an option for reading out salinity directly.

[Salinity in aquaculture, Part 1](#)

Chlorinity

The salinity of seawater was traditionally estimated from Cl^- concentration by the Knudsen equation:

$$\text{Salinity} = 1.80655 \text{ Cl}^-$$

where Cl^- (chloride concentration) is in grams per liter. Kits that allow chloride concentration to be measured by titration of water samples with mercuric nitrate are available. Salinity can be estimated accurately from chlorinity in the ocean and in estuaries, but in freshwaters and inland saline waters, the ratio of Cl^- to total dissolved substances often differs greatly from that of seawater.

Density




Fig. 2: A density hydrometer.

The density of freshwater is about 1 g/mL. Dissolved ions are denser than water and 1 gram of ions displaces less than 1 mL water. As a result, density increases with greater salinity (Table 1). The density of water can be measured with a hydrometer (Fig. 2). A traditional hydrometer is a cylindrical, air-filled bulb, cone-shaped at its bottom with a graduated stem protruding from its top. The bulb contains ballast causing the hydrometer to float upright. The distance that the stem extends above the surface depends upon water density, and the greater the density, the higher the stem rises above the surface. Hydrometers for determining salinity have stems calibrated in salinity units.

Boyd, salinity pt. 2, Table 1

Degrees-C	Salinity (0 g/L)	Salinity (10 g/L)	Salinity (20 g/L)	Salinity (30 g/L)	Salinity (40 g/L)
0	0.99984	1.0080	1.0160	1.0241	1.0321
5	0.99997	1.0079	1.0158	1.0237	1.0316
10	0.00070	1.0075	1.0153	1.0231	1.0309
15	0.99910	1.0068	1.0144	1.0221	1.0298
20	0.99821	1.0058	1.0134	1.0210	1.0286
25	0.99705	1.0046	1.0121	1.0196	1.0271
30	0.99565	1.0031	1.0105	1.0180	1.0255
35	0.99403	1.0014	1.0088	1.0162	1.0237
40	0.99222	0.9996	1.0069	1.0143	1.0217

Table 1. The density of water (g/cm3) of different salinities at selected temperatures between 0 and 40 degrees-C.

Refractive index

Light travels faster through some media than through others. According to Snell’s law, if the first medium is less dense than the second, light decreases in velocity upon entering the second medium causing it to refract towards the normal. The opposite occurs when light travels faster in the second medium than in the first. The index of refraction is the ratio of the speed of light in a vacuum to the speed of light in a second medium. The refraction of light by water is evident when one views from the side a drinking straw placed in a clear container of water (Fig. 3).

Fig. 3: Visual evidence that light is refracted by water.

The refractive index of water increases as a function of density, and is influenced also by wavelength of measurement, atmospheric pressure, and temperature. A good quality, handheld, salinity refractometers (Fig. 4) measure salinities of 1 to 60 mg/L to one decimal place. They are widely used to measure salinity at aquaculture facilities supplied with inland, saline water, estuarine water, or seawater.

Salinity and aquaculture

Some aquaculture species such as ictalurid catfish, pangasius and common carp grow best at salinities of <5 g/L; species such as Atlantic salmon, tilapia and rainbow trout grow well up to 20 g/L salinity; estuarine species such as penaeid shrimp grow well at salinities of 2 to 40 g/L. Marine and estuarine species can be farmed in inland saline water, but they may not survive and grow well in spite of adequate salinity. This results from ionic imbalance caused by low concentrations of K⁺, Mg²⁺, and Ca²⁺ or a combination of these cations. Mineral supplements are applied to increase concentrations of major ions.



Fig. 4. A handheld, salinity refractometer.

Boyd, salinity Pt. 2, Table 2

Salinity Food energy recovered as fish growth (%)	
0.5	33.4
2.5	31.8
4.5	22.2
6.5	20.1
8.5	10.4
10.5	-1.0

Table 2. Effect of salinity on recovery of food energy as growth in the common carp. Source: Wang et al. (1997).

Some freshwaters have very low concentrations of dissolved ions (low salinity), but ion concentration can be increased by liming and adding certain mineral salts. The only practical way of reducing salinity is by adding water of lower salinity to culture systems. This is sometimes done in ponds in arid regions or during prolonged drought. In fish and shrimp hatchery vessels, it is possible to regulate salinity by adding commercially available sea salt mixes of specific salts. Concentrated brine solutions from coastal, seawater evaporation ponds have been added to freshwater to allow inland culture of marine species.

References available from author.

Now that you've finished reading the article ...

... we hope you'll consider supporting our mission to document the evolution of the global aquaculture industry and share our vast network of contributors' expansive knowledge every week.

By becoming a Global Seafood Alliance member, you're ensuring that all of the pre-competitive work we do through member benefits, resources and events can continue. Individual membership costs just \$50 a year.

Not a GSA member? Join us.

[Support GSA and Become a Member](#)

Author



Claude E. Boyd, Ph.D.

School of Fisheries, Aquaculture and Aquatic Sciences
Auburn University
Auburn, Alabama 36849 USA

[117,100,101,46,110,114,117,98,117,97,64,49,101,99,100,121,111,98]

Share

- [✉ Share via Email](#)
- [🐦 Share on Twitter](#)
- [f Share on Facebook](#)
- [in Share on LinkedIn](#)

Tagged With

[density](#) [chlorinity](#) [refractive index](#) [Knudsen equation](#) [salinity](#) [electrical conductivity](#) [Claude E. Boyd](#)

Related Posts

Health & Welfare

[Dissolved oxygen dynamics](#)

Dissolved oxygen management is the most important requirement of aquaculture pond water quality. DO concentration below 3 mg/L is stressful to shrimp.

Responsibility

[Accuracy of custom water analyses varies](#)

The reliability of trace element analyses reported by custom laboratories cannot be checked by simple techniques, and results may not always be accurate. One should check the reliability of major ion analyses by determining the charge balance and comparing the measured total ion concentration with the total ion concentration estimated from conductivity.

Responsibility

[Drying, liming, other treatments disinfect pond bottoms](#)

The traditional way of destroying organisms in pond bottoms is thorough dry-out for a week or longer. Fish toxicants and liming can kill unwanted parasites.

Responsibility

[Light penetration in water](#)

Light penetrating water is scattered and absorbed exponentially as it passes downward. The presence of dissolved organic matter and suspended solids further impedes light penetration, and different types of solids absorb different wavelengths.

About The Advocate

The Responsible Seafood Advocate supports the Global Seafood Alliance's (GSA) mission to advance responsible seafood practices through education, advocacy and third-party assurances.

[Learn More](#)

Search Responsible Seafood Advocate



Advertising Opportunities

[2022 Media & Events Kit](#)

Categories

[Aquafeeds](#) > [Health & Welfare](#) > [From Our Sponsors](#) > [Innovation & Investment](#) > [Intelligence](#) > [Responsibility](#) [Responsibility](#) > [Fisheries](#) > [Artículos en Español](#) >

Don't Miss an Article

Featured

- [Health & Welfare](#) [An update on vibriosis, the major bacterial disease shrimp farmers face](#)
- [Intelligence](#) [A seat at the table: Fed By Blue team says aquaculture needs a stronger voice](#)
- [Responsibility](#) [Quantifying habitat provisioning at macroalgae cultivation locations](#)

Popular Tags

All Tags ▼

Recent

- [Fisheries](#) [Second Test: Another filler for the fisheries category.](#)
- [Fisheries](#) [Test: This is filler for the fisheries Category.](#)
- [Aquafeeds](#) [Test Article](#)
- [Responsibility](#) [Study: Climate change will shuffle marine ecosystems in unexpected ways as ocean temperature warms](#)
- [Health & Welfare](#) [Indian shrimp researchers earn a patent for WSSV diagnostic tool](#)



- [About](#)
- [Membership](#)
- [Resources](#)
- [Best Aquaculture Practices \(BAP\)](#)
- [Best Seafood Practices \(BSP\)](#)
- [GOAL Events](#)
- [Advocate Magazine](#)
- [Aquademia Podcast](#)
- [Blog](#)
- [Contact](#)

Stay up to date with GSA

- 
- 
- 
- 
- 

Copyright © 2024 Global Seafood Alliance
All rights reserved.

[Privacy](#)
[Terms of Use](#)
[Glossary](#)