



- [GOAL Events](#)
- [Advocate Magazine](#)
- [Aquademia Podcast](#)
- [Blog](#)
- [Contact](#)

-
-
-
-
-

- [Log In](#)



- [About](#)
 - [Who We Are](#)
 - [Our History](#)
 - [Our Team](#)
 - [Sustainable Development Goals](#)
 - [Careers](#)
- [Membership](#)
 - [Overview](#)
 - [Our Members](#)
 - [Corporate Membership](#)
- [Resources](#)
- [Certification](#)
 - [Best Aquaculture Practices](#)
 - [Best Seafood Practices](#)

Search...



[Log In](#)

- [About](#)
 - [Who We Are](#)
 - [Our History](#)
 - [Our Team](#)
 - [Sustainable Development Goals](#)
 - [Careers](#)
- [Membership](#)
 - [Overview](#)
 - [Our Members](#)
 - [Corporate Membership](#)
- [Resources](#)
- [Certification](#)
 - [Best Aquaculture Practices](#)
 - [Best Seafood Practices](#)
- [GOAL Events](#)
- [Advocate Magazine](#)
- [Aquademia Podcast](#)
- [Blog](#)
- [Contact](#)



Health & Welfare
Health & Welfare

Scombrottoxins, part 2



1 August 2005 George J. Flick, Jr., Ph.D.



Prevalence in seafood



Amberjack and other scombroid fish can potentially cause scombroid poisoning. Amberjack broodstock photo from the University of Miami Florida Keys Rosenstiel School of Marine and Atmospheric Science, courtesy of NOAA.

Biogenic amines from fish and shellfish are among the foremost causes of seafood-related morbidity worldwide. Seafood harvesters and processors should be keenly aware of the conditions that exacerbate the formation of these compounds in their products, the tendency of different areas of a fish carcass to support formation of these amines, and the effects of bacterial flora on the formation of these toxins.

Biogenic amines in fish

The prevalence of biogenic amines in fish depends on several factors. In general, concentrations in newly caught fish are low. For example, research found cadaverine values ranged 1.16 to 10.36 ppm in high-quality rockfish, salmon steaks, and shrimp, and putrescine levels ranged 1.36 to 6.30 ppm in high-quality lobster tails, salmon steaks, and shrimp. Another study reported that high-quality tuna had cadaverine and putrescine values ranging 0.24 to 5.32 and 0 to 1.84 ppm, respectively.

Biogenic amine prevalence also varies year to year. For example, a three-year study focused on biogenic amines in 102 samples of albacore tuna harvested off the northwest U.S. coast from 1994 to 1996. There were significant differences of amine levels in these fish depending on the year.

Total levels of spermine, spermidine, putrescine, cadaverine, histamine, and tyramine varied 5.9 to 56.5 ppm. These levels were probably lower due to the fact that the samples were frozen on board or chilled on board and immediately frozen after reaching the dock and kept at minus-40 degrees-C until analysis. Spermine was present at higher levels, followed by spermidine, histamine, putrescine, cadaverine, and tyramine.

Muscle type

Researchers have observed no difference in amine levels in upper and lower loin light muscles. However, dark muscles contained higher concentrations of spermidine (Table 1). Samples from intestine walls contained high amine levels.

One particularly broad study examined the amounts of histidine and histamine formed in 21 aquatic species during spoilage. The conclusions were consistent with those of other researchers: more histamine is produced in the red muscle fishes, such as tuna and mackerel, than in white muscle species such as rockfish.

Table 1. Levels of biogenic amines in varied tissues of albacore tuna. Source: Gloria et al., 1999

Samples	Spermine	Spermidine	Histamine	Putrescine	Cadaverine	Serotonin	Total
Light muscle upper loin	0.68 ± 0.12 ^b	0.26 ± 0.07 ^c	0.00 ^b	0.22 ± 0.07 ^{ab}	0.13 ± 0.02 ^b	0.00 ^b	1.29 ± 0.17 ^c
Light muscle lower loin	1.21 ± 0.26 ^b	0.25 ± 0.05 ^c	0.00 ^b	0.14 ± 0.05 ^b	0.11 ± 0.06 ^b	0.00 ^b	1.77 ± 0.37 ^c
Dark muscle	2.50 ± 0.97 ^{ab}	0.79 ± 0.18 ^b	0.00 ^b	0.06 ± 0.03 ^b	0.07 ± 0.05 ^b	0.00 ^b	3.42 ± 0.72 ^b
Intestine wall	5.35 ± 2.46 ^a	3.63 ± 1.18 ^a	0.52 ± 0.25 ^a	0.43 ± 0.16 ^a	1.96 ± 0.59 ^a	4.38 ± 1.33 ^a	16.3 ± 4.59 ^a

Microflora

A variety of microorganisms is able to produce biogenic amines. The production of cadaverine and putrescine is not surprising, since studies have shown that the covalent linking of cadaverine and putrescine to peptidoglycan is necessary for normal microbial growth. As such, production of these amines supports the continued expansion of microbial colonies on the surface of fish.

Several inoculation studies on both culture media and fish have shown that the microorganisms *Morganella* sp., *Proteus morganii*, *Proteus* sp., *Hafnia alvei*, and *Klebsiella* sp. can produce histamines and other biogenic amines. The majority of the studies also concurred that the potential of these microorganisms to produce toxic levels of biogenic amines is potentiated at abusive temperatures.

Tables 2 and 3, which summarize research on biogenic amines, show study results on the production of biogenic amines by bacterial isolates inoculated on different culture media and fish likely to be cold smoked, respectively. In addition, data from studies where isolates from fish were incubated and histamine production monitored are shown in Table 4.

Table 2. Production of biogenic amine by bacteria growing on media culture.

Histamine Producer	Histamine Concentration	Temperature and Time
<i>Morganella</i> sp.	1,000 ppm 1,000 ppm 0 ppm	5° C for 24 hours 25° C for 19 hours, followed by 5° C for 100 hours 5° C for 100 hours
<i>Proteus</i> sp.	large	
<i>Proteus morganii</i>	More than 200 nM/ml	15, 30, 37° C for less than 24 hours
<i>Enterobacter aerogenes</i>	More than 200 nM/ml	
<i>Klebsiella pneumoniae</i>	large	15, 30, 37° C for less than 24 hours
<i>Hafnia alvei</i>	large	30, 37° C for more than 48 hours
<i>Citrobacter freundii</i>	large	30, 37° C for more than 48 hours
<i>Escherichia coli</i>	large	30, 37° C for more than 48 hours

Similarly, when the relationship between microflora on horse mackerel and dominant spoilage bacteria was investigated, results showed that *Pseudomonas I/II*, *Pseudomonas III/IV-NH*, *Vibrio*, and *Photobacterium* were dominant when high levels of putrescine, cadaverine, and histamine were detected.

The activity of decarboxylase enzymes produced by bacteria can be an indirect measurement of potential for biogenic amine formation. A study showed that 14 bacterial isolates from mackerel tissue were capable of exhibiting decarboxylase activity (production of histamine, cadaverine, and putrescine) when incubated in Spanish mackerel at 0, 15 and 30 degrees-C. Many other bacteria also have strong histidine decarboxylase activities.

Best practices required

Since many microorganisms can cause scombrotxin illness, it is imperative that fishermen, processors, distributors and retailers employ high-level sanitary practices for products under their control. The production of histamine and other biogenic amines can be effectively controlled by cooling fish as quickly as possible after harvest, maintaining fish temperatures constantly under 4 degrees-C, and implementing effective sanitation programs at

processing and distribution facilities. Scombrottoxin poisoning can easily be avoided through adherence to good manufacturing practices from harvest through sale to consumers.

(Editor's Note: This article was originally published in the October 2005 print edition of the Global Aquaculture Advocate.)

Now that you've finished reading the article ...

... we hope you'll consider supporting our mission to document the evolution of the global aquaculture industry and share our vast network of contributors' expansive knowledge every week.

By becoming a Global Seafood Alliance member, you're ensuring that all of the pre-competitive work we do through member benefits, resources and events can continue. Individual membership costs just \$50 a year.

Not a GSA member? Join us.

[Support GSA and Become a Member](#)

Author



George J. Flick, Jr., Ph.D.

Food Science and Technology Department
Virginia Tech/Virginia Sea Grant (0418)
Blacksburg, Virginia 24061 USA

[117,100,101,46,116,118,64,103,107,99,105,108,102]

Share

- [✉ Share via Email](#)
- [🐦 Share on Twitter](#)
- [f Share on Facebook](#)
- [in Share on LinkedIn](#)

Tagged With

[George J. Flick](#) [Scombrottoxins](#)

Related Posts

[Health & Welfare](#)

[Scombrottoxins, part 1](#)

Without proper control of seafood handling and processing, high amino acid content and bacterial activity can rapidly elevate concentrations of biogenic amines like scombrottoxins.

[Health & Welfare](#)

[Scombrottoxins, part 3](#)

While freezing, salting, smoking and irradiation can minimize growth biogenic amine formation, special care from harvest through consumption can prevent scombrottoxin poisoning.

[Intelligence](#)

[An examination of seafood packaging](#)

Some substances can migrate from plastics and other seafood packaging materials into the product. Even if the substances are not harmful, they can affect the flavor and acceptability of the food.

[Aquafeeds](#)

[Animal byproduct concentrates useful tools in formulation](#)

With the market volatility of fishmeal, as well as rising sustainability concerns, the aquaculture industry is seeking sources of protein, such as animal byproduct concentrates, to substitute for fishmeal.

About The Advocate

The Responsible Seafood Advocate supports the Global Seafood Alliance's (GSA) mission to advance responsible seafood practices through education, advocacy and third-party assurances.

[Learn More](#)

Search Responsible Seafood Advocate



Advertising Opportunities

[2022 Media & Events Kit](#)

Categories

[Aquafeeds](#) > [Health & Welfare](#) [Health & Welfare](#) > [From Our Sponsors](#) > [Innovation & Investment](#) > [Intelligence](#) > [Responsibility](#) > [Fisheries](#) > [Artículos en Español](#) >

Don't Miss an Article

Featured

- [Health & Welfare](#) [An update on vibriosis, the major bacterial disease shrimp farmers face](#)
- [Uncategorized](#) [A seat at the table: Fed By Blue team says aquaculture needs a stronger voice](#)
- [Responsibility](#) [Quantifying habitat provisioning at macroalgae cultivation locations](#)

Popular Tags

All Tags

Recent

- [Fisheries Second Test: Another filler for the fisheries category.](#)
- [Fisheries Test: This is filler for the fisheries Category.](#)
- [Aquafeeds Test Article](#)
- [Responsibility Study: Climate change will shuffle marine ecosystems in unexpected ways as ocean temperature warms](#)
- [Health & Welfare Indian shrimp researchers earn a patent for WSSV diagnostic tool](#)



- [About](#)
- [Membership](#)
- [Resources](#)
- [Best Aquaculture Practices \(BAP\)](#)
- [Best Seafood Practices \(BSP\)](#)
- [GOAL Events](#)
- [Advocate Magazine](#)
- [Aquademia Podcast](#)
- [Blog](#)
- [Contact](#)

Stay up to date with GSA

-
-
-
-
-

Copyright © 2024 Global Seafood Alliance
All rights reserved.
[Privacy](#)
[Terms of Use](#)
[Glossary](#)