Global Seafood Alliance Logo

- GOAL Events
- Advocate Magazine
- Aquademia Podcast
- Blog
- Contact
- (O)

- Log In

- About
 - Who We Are
 - Our History
 - o Our Team
 - Sustainable Development Goals
 - <u>Careers</u>
- <u>Membership</u>
 - Overview
 - Our Members
 - Corporate Membership
- Resources
- **Certification**
 - Best Aquaculture Practices
 - Best Seafood Practices

Search...



- About
 - Who We Are
 - Our History
 - o Our Team
 - Sustainable Development Goals
 - Careers
- <u>Membership</u>
 - o <u>Overview</u>
 - Our Members
 - o Corporate Membership
- Resources
- **Certification**
 - Best Aquaculture Practices
 - Best Seafood Practices
- GOAL Events
- Advocate Magazine

- Aquademia Podcast
- Blog
- Contact



Selectively bred shrimp survive varied TSV exposure

Responsible Seafood Advocate logo

1 August 2003 Shaun M. Moss, Ph.D. Steve M. Arcel Dustin R. Moss Yu-Sen Hwang Brenda White-Noble Donald V. Lightner, Ph.D.



14-day study conducted at the University of Arizona



Selectively bred shrimp were maintained in 1,000-liter fiberglass tanks containing artificial seawater.

In 1999, significant mortalities of a Taura Syndrome Virus (TSV)-tolerant strain of the Western blue shrimp (Litopenaeus stylirostris) occurred at shrimp farms in Mexico. These mortalities resulted from TSV epizootics accompanied by a slight deviation in the gross signs typical of TSV infection, and there were concerns a new TSV strain had emerged.

Since 2000, researchers from the University of Arizona (UAZ) in Tucson, Arizona, USA, have analyzed TSV geographic and year isolates in relation to a reference isolate from Hawaii, USA, to determine possible differences in virulence and host range. Using selected Office International des Epizooties (OIE) diagnostic methods and sequence analysis of nucleotides and amino acids in the viral genome, the researchers compared the reference isolate with three TSV isolates from Mexico and found similar but distinct isolates of TSV.

New serotype from Belize

In 2001, significant mortalities of *L. vannamei* from TSV epizootics at shrimp farms in Belize raised concerns that another TSV strain had emerged. UAZ researchers purified the Belize 2002 isolate of TSV from cultured *L*. *vannamei* and found the virions were slightly but significantly larger than those from Hawaii and Mexico.

Again, using OIE diagnostic methods and sequence analysis, UAZ researchers concluded that the Belize isolate represented a new serotype. Importantly, broodstock suppliers to Belize reported that shrimp bred for resistance to the "old" Taura strain, the Hawaii isolate, succumbed to the "new" Belize strain.

Research to develop tolerance

In response to these concerns, researchers from the Oceanic Institute explored the possibility of developing selectively bred families of *L. vannamei* that exhibited tolerance to both the Hawaii and Belize TSV serotypes. Since 1995, researchers at Oceanic Institute have developed families of shrimp that exhibit high survival to the Hawaii TSV serotype.

Offspring from these unique families were produced and distributed to several U.S. broodstock suppliers and subsequently challenged with both TSV serotypes. Results from these challenge tests are summarized below.

Experimental challenge

Juvenile shrimp of about 2 grams from Rainbow Hawaii Farms of Mililani, Hawaii, USA, were shipped to UAZ and exposed to the TSV serotypes by feeding them infected shrimp tissue at 10 percent of their body weight once per day for three consecutive days. Shrimp survival was monitored daily, and final survival was determined after 14 days (Table 1).

Moss, Survival of two populations of L. vannamei, Table 1

| TSV Serotype | Shrimp Strain | Survival (%) |
|--------------|--|--------------|
| Hawaii TSV | O.I. "Kona" shrimp (Positive control) | 20 |
| Hawaii TSV | Rainbow Hawaii Farms shrimp | 95 |
| Belize TSV | O.I. "Kona" shrimp (Positive control) | 0 |
| Belize TSV | Rainbow Hawaii Farms shrimp | 63 |
| None | Rainbow Hawaii Farms shrimp (Negative control) | 100 |

Table 1. Survival of two populations of *L. vannamei* after 14 days when exposed to two TSV serotypes.

Positive and negative control groups were also evaluated. The negative control group, which consisted of selectively bred shrimp fed a commercial diet for the 14-day study, evaluated shrimp mortalities unrelated to TSV exposure. Positive control groups consisted of the Oceanic Institute "Kona" reference strain of L. vannamei exposed to both TSV serotypes. Kona shrimp are susceptible to TSV and exhibit consistently low survival of 0 to 40 percent after TSV exposure.

Positive controls were evaluated to ensure the virus-infected tissue used in the challenges was infectious. All selectively bred shrimp were maintained in 1,000-liter fiberglass tanks, whereas the Kona shrimp were maintained in 90-liter glass aquariums. The tanks and aquariums contained artificial seawater at 33 ppt salinity and 29 degrees-C and were equipped with biological filters.

Survival to TSV exposure

The selectively bred shrimp exhibited 95 percent survival after exposure to the Hawaii TSV serotype, while the Kona reference strain experienced only 20 percent survival. The selectively bred shrimp also had 63 percent survival after exposure to the Belize TSV serotype, whereas all the Kona shrimp died by day 4 of the challenge test.

These results indicated the Belize TSV serotype was more lethal than the Hawaii serotype, especially for the reference shrimp. However, through artificial selection or other genetic improvement techniques, shrimp breeders may be able to compete in the "arms race" against mutating viral pathogens.

(Editor's Note: This article was originally published in the August 2003 print edition of the Global Aquaculture Advocate.)

Now that you've finished reading the article ...

... we hope you'll consider supporting our mission to document the evolution of the global aquaculture industry and share our vast network of contributors' expansive knowledge every week.

By becoming a Global Seafood Alliance member, you're ensuring that all of the pre-competitive work we do through member benefits, resources and events can continue. Individual membership costs just \$50 a year.

Not a GSA member? Join us.

Support GSA and Become a Member

Authors

Shaun M. Moss, Ph.D.

Shaun M. Moss, Ph.D.

The Oceanic Institute 41-202 Kalanianaole Highway Waimanalo, Hawaii 96795 USA

[103,114,111,46,101,116,117,116,105,116,115,110,105,99,105,110,97,101,99,111,64,115,115,111,109,115]

Steve M. Arcel

Steve M. Arcel

The Oceanic Institute 41-202 Kalanianaole Highway Waimanalo, Hawaii 96795 USA

Dustin R. Moss

Dustin R. Moss

The Oceanic Institute 41-202 Kalanianaole Highway Waimanalo, Hawaii 96795 USA

Yu-Sen Hwang

Yu-Sen Hwang

Rainbow Hawaii Farms Mililani, Hawaii, USA

Brenda White-Noble

Brenda White-Noble

University of Arizona, Department of Veterinary Science Tucson, Arizona, USA



Donald V. Lightner, Ph.D.

University of Arizona, Department of Veterinary Science Tucson, Arizona, USA

Share

- Share via Email
- **Y** Share on Twitter
- **f** Share on Facebook
- in Share on LinkedIn

Tagged With

Donald V. Lightner Dustin R. Moss Shaun M. Moss Brenda White-Noble TSV exposure Steve M. Arcel Yu-Sen **Hwang**

Related Posts

Health & Welfare

Study: TSV exposure may lessen YHV effects in white shrimp

Since Taura syndrome virus was in the Americas when yellow head virus become pandemic in Asia, it is possible TSV prevented shrimp in the Western Hemisphere from infection by materials from Asia containing YHV. To test this idea, the authors evaluated the viral interactions in shrimp preinfected with TSV and then challenged with YHV.

Health & Welfare

A study of Zoea-2 Syndrome in hatcheries in India, part 1

Indian shrimp hatcheries have experienced larval mortality in the zoea-2 stage, with molt deterioration and resulting in heavy mortality. Authors investigated the problem holistically.

Health & Welfare

A study of Zoea-2 Syndrome in hatcheries in India, part 2

Indian shrimp hatcheries have experienced larval mortality in the zoea-2 stage, with molt deterioration and resulting in heavy mortality. Authors considered biotic and abiotic factors. Part 2 describes results of their study.

Health & Welfare

Challenging Pacific white postlarvae with AHPND


Study results indicate that P. vannamei challenged with AHPND in biofloc had higher survival rates than shrimp challenged in clear water.

About The Advocate

The Responsible Seafood Advocate supports the Global Seafood Alliance's (GSA) mission to advance responsible seafood practices through education, advocacy and third-party assurances.

Search Q

Learn More



Advertising Opportunities

2022 Media & Events Kit

Categories

Aquafeeds > Health & Welfare Health & Welfare > From Our Sponsors > Innovation & Investment > Intelligence > Responsibility > Fisheries > Artículos en Español >

Don't Miss an Article

Featured

- Health & Welfare An update on vibriosis, the major bacterial disease shrimp farmers face
- Intelligence A seat at the table: Fed By Blue team says aquaculture needs a stronger voice
- Responsibility Quantifying habitat provisioning at macroalgae cultivation locations

Popular Tags



Recent

- Fisheries Second Test: Another filler for the fisheries category
- Fisheries Test: This is filler for the fisheries Category
- Aquafeeds Test Article
- Responsibility Study: Climate change will shuffle marine ecosystems in unexpected ways as ocean temperature warms
- Health & Welfare Indian shrimp researchers earn a patent for WSSV diagnostic tool



- **About**
- <u>Membership</u>
- Resources
- Best Aquaculture Practices (BAP)
- Best Seafood Practices (BSP)
- GOAL Events
- Advocate Magazine
- Aquademia Podcast
- Blog
- **Contact**

Stay up to date with GSA

Copyright © 2024 Global Seafood Alliance All rights reserved.

Privacy Terms of Use <u>Glossary</u>