



- [GOAL Events](#)
- [Advocate Magazine](#)
- [Aquademia Podcast](#)
- [Blog](#)
- [Contact](#)
-
-
-
-
-
- [Log In](#)



- [About](#)
 - [Who We Are](#)
 - [Our History](#)
 - [Our Team](#)
 - [Sustainable Development Goals](#)
 - [Careers](#)
- [Membership](#)
 - [Overview](#)
 - [Our Members](#)
 - [Corporate Membership](#)
- [Resources](#)
- [Certification](#)
 - [Best Aquaculture Practices](#)
 - [Best Seafood Practices](#)

Search...



[Log In](#)

- [About](#)
 - [Who We Are](#)
 - [Our History](#)
 - [Our Team](#)
 - [Sustainable Development Goals](#)
 - [Careers](#)
- [Membership](#)
 - [Overview](#)
 - [Our Members](#)
 - [Corporate Membership](#)
- [Resources](#)
- [Certification](#)
 - [Best Aquaculture Practices](#)
 - [Best Seafood Practices](#)
- [GOAL Events](#)
- [Advocate Magazine](#)
- [Aquademia Podcast](#)
- [Blog](#)
- [Contact](#)



Health & Welfare

Health & Welfare

Shrimp hybrids outperform parents in Brazil studies



1 May 2009 Joao L. Rocha, Ph.D. Ana Carolina Guerrelhas, B.S. Ana Karina Teixeira, B.S. Flávio Augusto Farias, B.S. Ana Paula Teixeira, B.S.



Potential product underwent commercial trials at three shrimp farms



Technicians weighed shrimp during tests using electronic balances connected to computers that recorded data automatically.

Genearch Aquacultura, a new shrimp genetics company in Brazil, and Aquatec, the country's leading commercial hatchery, have been working for several years to develop local specific pathogen-free (SPF) lines of shrimp.

Given the generally poor pond results obtained with the first generation of KY30 and KY40, two non-locally adapted SPF lines, and their crosses, the authors more recently sought to evaluate the hybrid vigor that might result from crosses between the imported SPF lines and the non-SPF, genetically improved and locally adapted Speed Line. Hybrid crosses between the SPF lines and the Speed Line were designated SPKY30 and SPKY40.

Test design, analysis

Multiple spawns from each of the five shrimp lines were reared under normal larviculture protocols. Performance tests then involved two concurrent experiments. The first used five side-by-side 1,000-square-meter experimental ponds, each stocked with 50,000, 11- to 13-day-old postlarvae from only one line at 50 shrimp per square meter. Stocking, feeding and routine pond management followed standard commercial procedures for the region. The tests targeted an average harvest weight of 12-13 g from a grow-out cycle of about 80 days.

The second experiment involved animals that reached 2 to 3 grams in the above ponds. About 750 animals from each line were tagged with elastomers and stocked into each of the other ponds at a rate of 150 animals/line/pond. A total of 750 tagged animals from all five lines were placed in each pond. The grow-out time for the tagged animals was about 50 days and also targeted animals of 12 to 13 grams.

An overnight power failure that caused a drastic drop in oxygen levels and significant mortalities led to shortening of the test period and earlier pond harvests. All dead animals were collected, counted, weighed and identified by their tags. The five ponds were harvested in the next few days, which provided the opportunity to evaluate animals for their tolerance to low oxygen levels.

The presence of tagged animals from all lines in all five ponds allowed the quantification of the environmental effects for each pond and the ability to correct for them within a statistical framework. Analyses were first conducted separately for each experimental testing block for non-tagged and tagged animals, and subsequently a global statistical analysis was performed.

A third set of tests to confirm the results from the studies above followed similar experimental design and statistical approaches in the winter of 2008. After excellent pond results were obtained for the SPKY40 hybrid, this potential commercial product underwent commercial trials in 26 ponds at three local shrimp farms.

Significant results

Nearly all statistical effects tested were highly significant ($P < .0001$). Stocking weight was highly correlated with low oxygen mortality, indicating that animals which started the test at higher weights were more susceptible to low oxygen stress. Analyses conducted for each experiment with tagged and untagged animals revealed fairly similar results, with a few exceptions due to some pond effects that were statistically corrected. Hence, Figs. 1-4 show results for the different traits evaluated under the global statistical analysis framework. To facilitate interpretation, the lines are arranged by decreasing magnitude of their harvest biomass yield effects.




Fig. 1: Adjusted harvested biomass of nonlocal SPF lines, Speed Line and crosses.




Fig. 2: Adjusted harvest weight of individual shrimp in tested lines.

Fig. 3: Adjusted weekly growth of individual shrimp in tested lines.

Fig. 4: Adjusted mortality from low-oxygen stress in tested lines.

The figures show the performance advantage of the hybrid cross SPKY40 (Speed Line x SPFKY40) for all traits evaluated, with a harvest biomass advantage over the Speed Line of over 800 kg/ha and notable low-oxygen stress tolerance. This was the first time a line superior to the genetically improved and locally adapted Speed Line was identified. The “second place” ratings of the Speed Line showed the results of many years of genetic improvement work.

The hybrid crosses SPKY40 and SPKY30 clearly performed better than the two SPF “purebred” parents and even the two parents in the case of SPKY40, suggesting a potential for further exploration of hybrid vigor. Also, as shown in Figure 1, the two nonlocal SPF lines had lower pond performance.

At about 70 percent, pond survival was highest for SPKY40, followed by 55 percent survival for the Speed Line, 53 percent for SPKY30, 52 percent for KY40 and 43 percent for KY30. As shown in Figure 4, low-oxygen stress caused high mortality rates for both the KY30 line and its derived hybrid cross, indicating they may be very susceptible to stresses present in commercial production settings. However, the tolerance to low oxygen shown by KY40 was remarkable for a nonlocal line never subjected to genetic improvement in the country.

Multiple spawns from five shrimp lines were reared under normal larviculture protocols.

Hybrid superiority

Indicating the superiority of the hybrid cross over the average of the two parents, heterosis values for SPKY40 were 9.1 percent for harvest weight, 11.0 percent for weekly growth, 28.8 percent for growout survival, 40.4 percent for harvest biomass yield and 17.1 percent for low-oxygen mortality. In the same order, values for SPKY30 were 6.3 percent, 3.1 percent, 7.9 percent, 14.3 percent and -17.9 percent. The considerable heterosis values for SPKY40, especially for survival and harvest yield, warrant further work on a hybrid cross.

The results obtained in the third set of tests confirmed those from the second tests. More importantly, the commercial farm trials run throughout 2008 also showed the superiority of SPKY40 over its commercial parent, the Speed Line, with a higher harvest biomass of around 700 kg/ha.

(Editor's Note: This article was originally published in the May/June 2009 print edition of the Global Aquaculture Advocate.)

Now that you've finished reading the article ...

... we hope you'll consider supporting our mission to document the evolution of the global aquaculture industry and share our vast network of contributors' expansive knowledge every week.

By becoming a Global Seafood Alliance member, you're ensuring that all of the pre-competitive work we do through member benefits, resources and events can continue. Individual membership costs just \$50 a year.

Not a GSA member? Join us.

[Support GSA and Become a Member](#)

Authors

-  Joao L. Rocha, Ph.D.

Joao L. Rocha, Ph.D.

Genearch Aquacultura and Aquatec
Barra do Cunhaú, Rio Grande do Norte
CEP 59190, Brazil

[114,98,46,109,111,99,46,104,99,114,97,101,110,101,103,64,97,104,99,111,114,110,104,111,106]

-  Ana Carolina Guerrelhas, B.S.

Ana Carolina Guerrelhas, B.S.

Genearch Aquacultura and Aquatec
Barra do Cunhaú, Rio Grande do Norte
CEP 59190, Brazil

-  Ana Karina Teixeira, B.S.

Ana Karina Teixeira, B.S.

Genearch Aquacultura and Aquatec
Barra do Cunhaú, Rio Grande do Norte
CEP 59190, Brazil

-  Flávio Augusto Farias, B.S.

Flávio Augusto Farias, B.S.

Genearch Aquacultura and Aquatec
Barra do Cunhaú, Rio Grande do Norte
CEP 59190, Brazil

-  Ana Paula Teixeira, B.S.

Ana Paula Teixeira, B.S.

Genearch Aquacultura and Aquatec
Barra do Cunhaú, Rio Grande do Norte
CEP 59190, Brazil

Share

-  [Share via Email](#)
-  [Share on Twitter](#)
-  [Share on Facebook](#)
-  [Share on LinkedIn](#)

Tagged With

[João L. Rocha](#) [Genetic](#) [Ana Carolina Guerrelhas](#) [Ana Karina Teixeira](#) [Flávio Augusto Farias](#) [Ana Paula Teixeira](#) [Shrimp](#) [Brazil](#) [mortality](#) [hybrid](#)

Related Posts

Intelligence

[Brazilian aquaculture: Constraints and challenges \(Part 1\)](#)

The Brazilian aquaculture industry has been growing steadily during the last two decades. Despite facing a number of challenges it is looking at continued growth and a larger role in the export markets.

Intelligence

[Considerations for tilapia farming in saltwater environments](#)

Tilapias are farmed in a variety of production systems, but mostly in freshwater and low-salinity waters. But tilapias are an excellent candidate for aquaculture in brackish- and seawater because they can tolerate a wide range of water salinity.

Health & Welfare

[Testing dietary potassium diformate in tilapia feed](#)

A study conducted in Huila, Colombia, tested the effects of the feed additive potassium diformate on the performance of juvenile tilapia. Results showed an enhanced fish productivity index, lower mortality and improvements in weight gain.

Intelligence

[Walleye and saugeye growing successfully indoors](#)

Past and ongoing studies at the UWSP Northern Aquaculture Demonstration Facility (UWSP-NADF) have shown that walleye and hybrid walleye can be successfully raised indoors using early advanced spawning, incubation and rearing techniques. Marketing studies indicate good aquaculture potential and high demand.

About The Advocate

The Responsible Seafood Advocate supports the Global Seafood Alliance's (GSA) mission to advance responsible seafood practices through education, advocacy and third-party assurances.

[Learn More](#)

Search Responsible Seafood Advocate



Advertising Opportunities

[2022 Media & Events Kit](#)

Categories

[Aquafeeds](#) > [Health & Welfare](#) [Health & Welfare](#) > [From Our Sponsors](#) > [Innovation & Investment](#) > [Intelligence](#) > [Responsibility](#) > [Fisheries](#) > [Artículos en Español](#) >

Don't Miss an Article

Featured

- [Health & Welfare](#) [An update on vibriosis, the major bacterial disease shrimp farmers face](#)
- [Uncategorized](#) [A seat at the table: Fed By Blue team says aquaculture needs a stronger voice](#)
- [Responsibility](#) [Quantifying habitat provisioning at macroalgae cultivation locations](#)

Popular Tags

All Tags

Recent

- [Fisheries Second Test: Another filler for the fisheries category.](#)
- [Fisheries Test: This is filler for the fisheries Category.](#)
- [Aquafeeds Test Article](#)
- [Responsibility Study: Climate change will shuffle marine ecosystems in unexpected ways as ocean temperature warms](#)
- [Health & Welfare Indian shrimp researchers earn a patent for WSSV diagnostic tool](#)



- [About](#)
- [Membership](#)
- [Resources](#)
- [Best Aquaculture Practices \(BAP\)](#)
- [Best Seafood Practices \(BSP\)](#)
- [GOAL Events](#)
- [Advocate Magazine](#)
- [Aquademia Podcast](#)
- [Blog](#)
- [Contact](#)

Stay up to date with GSA

-
-
-
-
-

Copyright © 2024 Global Seafood Alliance
All rights reserved.
[Privacy](#)
[Terms of Use](#)
[Glossary](#)