



- [GOAL Events](#)
- [Advocate Magazine](#)
- [Aquademia Podcast](#)
- [Blog](#)
- [Contact](#)
- 
- 
- 
- 
- 
- [Log In](#)



- [About](#)
  - [Who We Are](#)
  - [Our History](#)
  - [Our Team](#)
  - [Sustainable Development Goals](#)
  - [Careers](#)
- [Membership](#)
  - [Overview](#)
  - [Our Members](#)
  - [Corporate Membership](#)
- [Resources](#)
- [Certification](#)
  - [Best Aquaculture Practices](#)
  - [Best Seafood Practices](#)

Search...

[Log In](#)

- [About](#)
  - [Who We Are](#)
  - [Our History](#)
  - [Our Team](#)
  - [Sustainable Development Goals](#)
  - [Careers](#)
- [Membership](#)
  - [Overview](#)
  - [Our Members](#)
  - [Corporate Membership](#)
- [Resources](#)
- [Certification](#)
  - [Best Aquaculture Practices](#)
  - [Best Seafood Practices](#)
- [GOAL Events](#)
- [Advocate Magazine](#)
- [Aquademia Podcast](#)
- [Blog](#)
- [Contact](#)





## Tasmanian salmon farms examine net biofouling to reduce impacts



3 May 2013 Belinda Yaxley Dr. Dominic O'Brien



## Industry evaluates 'footprint' of in situ net-cleaning systems



The degree of net fouling varies considerably between summer (left) and winter (right) months.

Worldwide, the salmon aquaculture industry is addressing concerns regarding potential negative impacts on marine ecosystems by reducing its reliance on the use of copper-based anti-fouling coatings to control biofouling on nets. With the recent development of more efficient and economical in situ net-cleaning systems, the fish-farming industry in Tasmania, Australia, has committed to a strategy for the reduced use of copper-based anti-fouling coatings.

Two Tasmanian salmon aquaculture companies, Tassal Operations Pty. Ltd. and Huon Aquaculture Group Pty. Ltd., have embarked on an Australian government-funded project under the Caring for Our Country program to address potential water quality issues associated with in situ net cleaning. The results from this project will assist in developing best practice guidelines and promoting environmentally sustainable practices on farms that use in situ net cleaning within the greater salmon-farming industry in Tasmania.

## Objectives



CARING  
FOR  
OUR  
COUNTRY

Upon adopting in situ net cleaning, it became apparent to Huon Aquaculture Group and Tassal Operations that little scientific information was available on the possible environmental effects of this cleaning method, both within lease areas and in the greater marine environments surrounding the farms.

To investigate the potential impacts of in situ net cleaning on water quality, the project has three objectives:

- Define the types of fouling organisms that grow on nets and determine their seasonality.
- Characterize the net wash material expelled by in situ net cleaning.
- Use deposition models to demonstrate the footprint of the net wash.

## Fouling organisms



The authors identified *Ectopleura crocea* as a major net-fouling organism.

Identifying the types of fouling organisms, their preferences for different net types, timing of settlement and seasonality allows for the alteration of operational procedures to maximize the effectiveness of cleaning machines. For example, some fouling organisms are harder to clean off than others, so if cleaning is done more regularly at certain times of the year, difficult types may be easier to remove or kept from settling on the net altogether.

To understand seasonality and net preference, a methodology developed by Dr. Simone Dürr of Liverpool John Moores University in the United Kingdom was employed. For a period of two years, experimental frames housing different net types were deployed at the Huon and Tassal farms in southeast Tasmania. Each month, the frames were retrieved and sampled, with the selected net pieces replaced before redeployment.

The first year's data identified the dominant fouling groups on all net types as amphipod housings and the hydroid species *Plumularia setacea*, *Sarsia eximia*, *Obelia australis* and *Ectopleura crocea*, which together occupied 90 percent of the biological sample points. When looking at fouling assemblages among net types, the results indicated significant differences among net types and seasons.

## Water quality

To investigate potential impacts on water quality due to in situ net cleaning, the net wash was characterized by species composition and quantity. Analytes and size fractions were chosen based on the efficiency of previous sampling techniques and discussion with independent environmental chemists. The fates of copper, aluminum, phosphate, nitrate, nitrogen, carbon and ammonia were determined by quantifying analytes from a sample of net wash collected directly from the net cleaner.

Of particular interest to state and federal regulators was the amount of copper released from the nets during the cleaning process. To investigate the relationships between the amount of copper released to the environment and the number of net cleans, net wash was collected from cages treated with copper-based paint that had undergone differing numbers of cleanings.

While there was no statistically significant relationship between the numbers of cleanings and amounts of copper release, Tassal and Huon provided descriptive statistics that were reported to regulatory bodies. The data demonstrated the amount of copper released was thought to be within a comfortable range.

Determining the destination of net wash materials released to the environment was achieved using settlement testing. This established the particle size and velocity of the net wash. In addition, hydrodynamic data was collected at sites with different flows. Both the settlement tests and hydrodynamic information were critical in modeling the deposition of copper within the farm lease.




Fig. 1: Copper deposition ( $\text{mg}/\text{m}^2$ ) at a high-deposition farm site. The hydrodynamic regime plays a significant role in defining sediment impacts. Typically, a narrow footprint is observed in low-flow conditions, whereas the footprint widens under medium to high flow.

To understand the copper footprint from in situ net cleaning, a deposition model was employed to demonstrate the movement of the analytes in the water column for high- and low-deposition sites. At the high-deposition site (Fig. 1), interpretation of the model revealed that copper deposition reached  $10.0 \text{ mg}/\text{m}^2$  and above directly downstream of the source, but quickly reduced to concentrations of approximately  $0.3 \text{ mg}/\text{m}^2$  150 m from the release point.

*Editor's Note: Since the time this article was written, Tassal Operations has stopped using copper-treated nets.*

*(Editor's Note: This article was originally published in the May/June 2013 print edition of the Global Aquaculture Advocate.)*

## Authors

- Belinda Yaxley

Belinda Yaxley

Tassal Operations Pty. Ltd  
GPO Box 1645  
Hobart, Tasmania, Australia 7001

[117,97,46,109,111,99,46,108,97,115,115,97,116,64,121,101,108,120,97,121,46,97,100,110,105,108,101,98]

- Dr. Dominic O'Brien

Dr. Dominic O'Brien

Huon Aquaculture Group Pty. Ltd.  
Dover, Tasmania, Australia

## Share

- [Share via Email](#)
- [Share on Twitter](#)
- [Share on Facebook](#)
- [Share on LinkedIn](#)

## Tagged With

[Dominic O'Brien](#) [salmon](#) [Australia](#) [water quality](#) [biofouling](#) [tasmanian](#) [Belinda Yaxley](#).

## Related Posts

### Intelligence

#### [A land grab for salmon \(and shrimp\) in upstate New York](#)

The operators of Hudson Valley Fish Farm see their inland locale as a pilot to prove that land-based fish farming, located in close proximity to major metropolitan markets, can be successful.

Health & Welfare

#### [Sex determination in Tasmanian Atlantic salmon](#)

The maturation of Atlantic salmon prior to harvest is costly, so since male salmon are more likely to mature early, the salmon industry in Tasmania relies on all-female production.

Innovation & Investment

#### [Automatic submersible fish cage systems counter weather, surface problems](#)

The development of submersible fish cage technologies may be necessary to avoid the operational challenges of surface-based aquaculture, which can include extreme temperature and weather conditions, jellyfish infestation, oil spills and many types of biofouling.

Intelligence

#### [An engineer's design for a classroom aquaculture-aquaponics system](#)

An aquaponics teaching system was designed, built and operated by students at the University of Arizona, integrating its operation and management into the educational curriculum. This engineering design will require minimum maintenance and will last years.

## About The Advocate

The Responsible Seafood Advocate supports the Global Seafood Alliance's (GSA) mission to advance responsible seafood practices through education, advocacy and third-party assurances.

[Learn More](#)

Search Responsible Seafood Advocate





## Advertising Opportunities

[2022 Media & Events Kit](#)

## Categories

[Aquafeeds](#) > [Health & Welfare](#) > [From Our Sponsors](#) > [Innovation & Investment](#) > [Intelligence](#) > [Responsibility Responsibility](#) > [Fisheries](#) > [Artículos en Español](#) >

## Don't Miss an Article

### Featured

- [Health & Welfare](#) [An update on vibriosis, the major bacterial disease shrimp farmers face](#)
- [Intelligence](#) [A seat at the table: Fed By Blue team says aquaculture needs a stronger voice](#)
- [Responsibility](#) [Quantifying habitat provisioning at macroalgae cultivation locations](#)

## Popular Tags

All Tags ▼

### Recent

- [Fisheries](#) [Second Test: Another filler for the fisheries category](#)
- [Fisheries](#) [Test: This is filler for the fisheries Category](#)
- [Aquafeeds](#) [Test Article](#)
- [Responsibility](#) [Study: Climate change will shuffle marine ecosystems in unexpected ways as ocean temperature warms](#)
- [Health & Welfare](#) [Indian shrimp researchers earn a patent for WSSV diagnostic tool](#)



- [About](#)
- [Membership](#)
- [Resources](#)
- [Best Aquaculture Practices \(BAP\)](#)
- [Best Seafood Practices \(BSP\)](#)
- [GOAL Events](#)
- [Advocate Magazine](#)
- [Aquademia Podcast](#)

- [Blog](#)
- [Contact](#)

Stay up to date with GSA

- 
- 
- 
- 
- 

Copyright © 2024 Global Seafood Alliance

All rights reserved.

[Privacy](#)

[Terms of Use](#)

[Glossary](#)