



- [GOAL Events](#)
- [Advocate Magazine](#)
- [Aquademia Podcast](#)
- [Blog](#)
- [Contact](#)
-
-
-
-
-
- [Log In](#)



- [About](#)
 - [Who We Are](#)
 - [Our History](#)
 - [Our Team](#)
 - [Sustainable Development Goals](#)
 - [Careers](#)
- [Membership](#)
 - [Overview](#)
 - [Our Members](#)
 - [Corporate Membership](#)
- [Resources](#)
- [Certification](#)
 - [Best Aquaculture Practices](#)
 - [Best Seafood Practices](#)



[Log In](#)

- [About](#)
 - [Who We Are](#)
 - [Our History](#)
 - [Our Team](#)
 - [Sustainable Development Goals](#)
 - [Careers](#)
- [Membership](#)
 - [Overview](#)
 - [Our Members](#)
 - [Corporate Membership](#)
- [Resources](#)
- [Certification](#)
 - [Best Aquaculture Practices](#)
 - [Best Seafood Practices](#)
- [GOAL Events](#)
- [Advocate Magazine](#)
- [Aquademia Podcast](#)
- [Blog](#)
- [Contact](#)



Tips for growing bigger shrimp



1 October 2005 Shaun M. Moss, Ph.D. PingSun Leung, Ph.D.



Optimizing strategies for growing larger *L. vannamei*



The Oceanic Institute has grown large white shrimp at high density in a biosecure raceway.

Although the giant tiger prawn (*Penaeus monodon*) is the most common shrimp species cultured in Asia, its dominance has faded in recent years as many Asian farmers stocked their ponds with Pacific white shrimp (*Litopenaeus vannamei*). Now more *L. vannamei* are produced in Asia than the Western Hemisphere, contributing to a glut of white shrimp and significant drops in shrimp prices.

Alternative production strategies

Shrimp farmers in the United States and elsewhere are considering alternative production and harvesting strategies to avoid direct competition with foreign imports from Asia, including the production of larger white shrimp. However, there is little published information on growth curves and feeding rates for larger *L. vannamei*, and even less information on optimal harvesting strategies.

O.I. research

In an attempt to grow large shrimp at high densities under biosecure conditions, researchers at the Oceanic Institute stocked specific pathogen-free juvenile *L. vannamei* in a 58.4-square meter raceway (Fig. 1) at a density of 705 shrimp per cubic meter. After 86 days, shrimp survival in the phase 1

study was 70.3 percent, mean harvest weight was 17.9 grams, and mean growth rate was 1.37 grams per week (Table 1). Production was 8.9 kilograms per cubic meter, with mean water exchange throughout the trial of less than 2 percent per day.

Moss, Performance of *L. vannamei* grown in a raceway, Table 1

Parameter	Phase 1 Trial	Phase 2 Trial
Stocking density (shrimp/m ³)	705	118
Trial duration (days)	86	69
Stocking weight (g)	1.06	17.90
Harvest weight (g)	17.9	34.20
Survival (%)	70.3	87.80
Growth rate (g/week)	1.37	1.65
Production (kg/m3)	8.9	3.50
Water use (l/kg shrimp)	219	229

Table 1. Performance of *L. vannamei* grown in a raceway. About 24% of the shrimp harvested from phase 1 were restocked for phase 2.

Immediately following the first harvest, about 24 percent of the harvested shrimp were restocked into the raceway, which contained 55 percent of the original water, at 118 shrimp per cubic meter. In this phase 2 work, shrimp survival was 87.8 percent after 69 days, mean harvest weight was 34.2 grams, and mean growth rate was 1.65 grams per week.

During the second grow-out phase, shrimp exhibited sexual growth dimorphism. Females grew 1.88 grams per week and were harvested at 36.9 grams, whereas males grew 1.42 grams per week and were harvested at 31.5 grams. The trial indicated shrimp can be grown to 34.2 grams in 155 days at high densities with minimal water use.

Compensatory growth

Interestingly, weekly growth rates increased from 0.95 grams during week 12, immediately prior to the first harvest, to 2.6 grams during the first two weeks after the initial harvest (Fig. 1). This dramatic increase suggested that *L. vannamei* may exhibit compensatory growth.




Fig. 1: Shrimp growth over time in a biosecure raceway. Note the significant increase in growth immediately after the partial harvest at week 12.

Compensatory growth occurs when an organism exhibits faster than normal growth during a period of recovery from growth restriction. In some fish, compensatory growth can be so intense that starved and refed individuals exhibit higher mean growth rates than those fed continuously.

In this case, growth restriction likely resulted from the high stocking density, as growth is density-dependent in penaeid shrimp. Earlier work at the Oceanic Institute showed that *L. vannamei* exhibit a rapid and significant increase in RNA concentration and RNA/DNA ratio following the reintroduction of feeding after several days of starvation. However, this was the first indication that the species may exhibit compensatory growth when relieved of density pressure.

Optimal harvesting

From a farm management perspective, it may be possible to exploit this compensatory growth through partial harvesting strategies to increase production and profitability.

A partial budget analysis conducted by the Oceanic Institute using data from the grow-out trial indicated that partial harvesting was 8.8 percent more profitable than a single harvest if phase 2 shrimp were harvested after four weeks. In contrast, if phase 2 were harvested after 10 weeks, there would have been a 5.9 percent loss in net revenue compared to a single harvest. In this case, the increase in yield for partial harvesting did not compensate for the opportunity loss of running more regular cycles.

The partial budget analysis used a unit of analysis of a single raceway and assumed 3.87 crops per year for the single-harvest scenario, 3.01 crops for the four-week phase 2 approach, and 2.25 crops with a phase 2 of 10 weeks. Prices for head-off shrimp with 2:1 tail per head ratios were based on prices from U.S. \$2.58 per pound for 36 per 40 shrimp to \$6.50 per pound for UN per 15 shrimp. Feed cost was assumed to be \$0.70 per kilogram, with other costs considered similar among the scenarios. Partial harvesting would increase harvesting costs somewhat.

Although this analysis suggested partial harvesting can be beneficial, the strategy has not been optimized. Growth curves and feeding rates for larger shrimp under different production scenarios are needed to develop efficient harvest models. Ultimately, these models can be used by farmers to increase production and improve profitability.

(Editor's Note: This article was originally published in the October 2005 print edition of the Global Aquaculture Advocate.)

Now that you've finished reading the article ...

... we hope you'll consider supporting our mission to document the evolution of the global aquaculture industry and share our vast network of contributors' expansive knowledge every week.

By becoming a Global Seafood Alliance member, you're ensuring that all of the pre-competitive work we do through member benefits, resources and events can continue. Individual membership costs just \$50 a year.

Not a GSA member? Join us.

[Support GSA and Become a Member](#)


Authors

-  Shaun M. Moss, Ph.D.

Shaun M. Moss, Ph.D.

The Oceanic Institute
41-202 Kalanianaʻole Highway
Waimanalo, Hawaii USA

[103,114,111,46,101,116,117,116,105,116,115,110,105,99,105,110,97,101,99,111,64,115,115,111,109,115]

-  PingSun Leung, Ph.D.

PingSun Leung, Ph.D.

Department of Molecular Biosciences and Bioengineering
University of Hawaii at Manoa
Honolulu, Hawaii USA

Share

- [✉ Share via Email](#)
- [🐦 Share on Twitter](#)
- [f Share on Facebook](#)
- [in Share on LinkedIn](#)

Tagged With

[Shrimp](#) [L. vannamei](#) [PingSun Leung](#) [Shaun M. Moss](#)

Related Posts

Responsibility

[Aquaculture gives endangered totoaba a fighting chance](#)

The tenuous fate of a pint-sized porpoise, the critically endangered vaquita, is linked to a fish targeted by poachers fueling China's appetite for maws. The vaquita remains in peril, but aquaculture presents some hope for the totoaba.

Intelligence

[Bangladesh seeks more buck for its 'bagda'](#)

As more than 80 percent of Bangladeshi shrimp exports already go to EU markets, a consultation meeting involving buyers from the bloc and Bangladesh industry stakeholders and authorities was held at the end of last month in Utrecht, the Netherlands.

Health & Welfare

[Biosecurity practices on fish farms need beefing up](#)

Biosecurity measures and preventive strategies are essential in any biological production chain. Properly planned and implemented biosecurity programs will enhance animal health, production and economics.

Health & Welfare

[Building a better shrimp nursery, part 1](#)

Shrimp nursery systems offer an important opportunity to increase profits. Properly designed and operated nurseries are highly biosecure facilities to grow postlarvae at very high densities.

About The Advocate

The Responsible Seafood Advocate supports the Global Seafood Alliance's (GSA) mission to advance responsible seafood practices through education, advocacy and third-party assurances.

[Learn More](#)

Search Responsible Seafood Advocate



Advertising Opportunities

[2022 Media & Events Kit](#)

Categories

[Aquafeeds](#) > [Health & Welfare](#) > [From Our Sponsors](#) > [Innovation & Investment](#) > [Intelligence Intelligence](#) > [Responsibility](#) > [Fisheries](#) > [Artículos en Español](#) >

Don't Miss an Article

Featured

- [Health & Welfare](#) [An update on vibriosis, the major bacterial disease shrimp farmers face](#)
- [Uncategorized](#) [A seat at the table: Fed By Blue team says aquaculture needs a stronger voice](#)
- [Responsibility](#) [Quantifying habitat provisioning at macroalgae cultivation locations](#)

Popular Tags

All Tags ▼






Recent

- [Fisheries](#) [Second Test: Another filler for the fisheries category](#)
- [Fisheries](#) [Test: This is filler for the fisheries Category](#)
- [Aquafeeds](#) [Test Article](#)
- [Responsibility](#) [Study: Climate change will shuffle marine ecosystems in unexpected ways as ocean temperature warms](#)
- [Health & Welfare](#) [Indian shrimp researchers earn a patent for WSSV diagnostic tool](#)



- [About](#)
- [Membership](#)
- [Resources](#)
- [Best Aquaculture Practices \(BAP\)](#)
- [Best Seafood Practices \(BSP\)](#)
- [GOAL Events](#)
- [Advocate Magazine](#)
- [Aquademia Podcast](#)
- [Blog](#)
- [Contact](#)

Stay up to date with GSA

- 
- 
- 
- 
- 

Copyright © 2024 Global Seafood Alliance
All rights reserved.

[Privacy](#)

[Terms of Use](#)

[Glossary](#)