



- [GOAL Events](#)
- [Advocate Magazine](#)
- [Aquademia Podcast](#)
- [Blog](#)
- [Contact](#)

-
-
-
-
-

- [Log In](#)

☰

- [About](#)
 - [Who We Are](#)
 - [Our History](#)
 - [Our Team](#)
 - [Sustainable Development Goals](#)
 - [Careers](#)
- [Membership](#)
 - [Overview](#)
 - [Our Members](#)
 - [Corporate Membership](#)
- [Resources](#)
- [Certification](#)
 - [Best Aquaculture Practices](#)
 - [Best Seafood Practices](#)

Search...



[Log In](#)

- [About](#)
 - [Who We Are](#)
 - [Our History](#)
 - [Our Team](#)
 - [Sustainable Development Goals](#)
 - [Careers](#)
- [Membership](#)
 - [Overview](#)
 - [Our Members](#)
 - [Corporate Membership](#)
- [Resources](#)
- [Certification](#)
 - [Best Aquaculture Practices](#)
 - [Best Seafood Practices](#)
- [GOAL Events](#)
- [Advocate Magazine](#)
- [Aquademia Podcast](#)
- [Blog](#)
- [Contact](#)



Intelligence
Intelligence


Top eel farm upgrades effluent treatment in Netherlands

[Responsible Seafood Advocate logo](#)

1 June 2001 Andries Kamstra Ep Eding Oliver Schneider



Water consumption is roughly 500 liters per kilogram of fish produced




This storage unit at Nijvis stores concentrated sludge.

Because of the Netherlands' climate and limited available water resources, all fish farming in the country is based solely on the use of recirculation systems. Yet during the last decade, the aquaculture industry here has grown to represent roughly 60 eel farms with annual production of 3,500 metric tons (MT), and 15 farms for African catfish that produce about 2,000 MT each year.

The largest and most advanced eel farm in the Netherlands is Nijvis, with an annual production around 1,000 MT of *Anguilla anguilla*. To better deal with ongoing waste issues in the country, this farm recently invested in an expanded, state-of-the-art effluent treatment system.

Environmental constraints



A new cooling unit (right) joined the denitrification units at Nijvis.

Holland is a small country of 40,000 square km with a very strong tradition in animal husbandry. It is still the third-largest exporting country of agricultural products in the world. A large part of its animal-production business – which includes swine, poultry, and fish – is based on the use of imported feed stuffs, not farm-grown feed.

It is not difficult to imagine that environmental problems can arise under these conditions. The limited area of arable land was often used as a dumpsite for excess manure. This resulted in elevated nitrate levels in groundwater, and excessive nitrogen deposition by air in sensitive natural areas.

The government recently corrected this situation through strict legislation that limits the amount of minerals a farmer can apply. Surface and groundwater also are strongly protected, and users of these resources have to pay high levies.

Cost of wastewater

Industries and households are charged for wastewater based on the amount of oxygen needed to break down their waste loads.

The 4.57 Nkj element in the formula accounts for the oxygen demand of nitrification. At a large farm like Nijvis, wastewater is sampled daily and water consumption is measured continuously.

System design

The water treatment system at most eel farms involves a rather simple and straightforward design, with slight individual adaptations. The recirculating flow from the fish tanks is first passed through a mechanical drum or disc filter with a mesh of 40-100 μ for removal of suspended solids, then through a biological filter for nitrification. A trickling filter is usually used for this purpose, with the advantage of efficient stripping of carbon dioxide.

Some farms, including Nijvis, direct part of the flow over a submerged filter for removal of small suspended solids. In this system configuration, accumulation of nitrate is the factor that limits production. Make-up water is added to the system to keep nitrate level below 500 milligrams per liter. This results in a water consumption of roughly 500 liters per kilogram of fish produced.

Sludge



Fig. 1. Concentrated sludge accumulates on the the revolving fabric of the belt filter.




Figure 2.




Figure 3.

Before treatment, the effluent looks like Fig. 2. After treatment (Fig. 3), very little solid material remains. Phosphorus is also precipitated in the sludge process.

In the biological filter of a recirculation system, most of the soluble COD and ammonia is oxidized, which reduces the oxygen demand of the effluent by approximately 30 percent. However, the sludge separated from the mechanical filter still has a substantial COD. As long as this fraction is not retained, the levy is considerable (approximately 10,000 IE per year for Nijvis).

When the solid fraction is retained efficiently, the number of IEs can go down to around 1,000 for a farm with an annual production of 1,000 MT. One IE costs U.S. \$75, equivalent to U.S. \$.08 per kilograms fish produced.

The sludge from a mechanical filter is not very concentrated (roughly 0.3 percent DM), and concentration is necessary for efficient storage. To tackle this problem, Nijvis installed a "belt filter" together with a flocculation unit to treat the solid waste. To reduce water consumption, a denitrification unit was concurrently installed in a separate loop.

Advanced effluent treatment

All the sludge from the different mechanical filters is fed to a mixing unit, where a flocculant is added. The mixture is fed to a belt filter (Fig. 1), with a continuously revolving fabric from which the concentrated sludge is scraped. The 3 percent DM sludge is fed to a storage unit and applied as fertilizer at regular intervals.

The remaining liquid fraction is discharged to the sewer. Figs. 2 and 3 show the water before and after treatment. Little solid material remains after treatment. As a bonus, a large part of the phosphorus is precipitated in the sludge.

Denitrification is accomplished in a bypass fed directly with system water to two large tanks filled with floating beads moved by a propeller. The carbon source for denitrification is methanol, which is stored in an underground tank. The effluent of the reactors goes through a drum filter and is returned to the system.

Although sludge from the system could be also a good source to feed a denitrification reactor, methanol was chosen due to ease of operation. A newly installed cooling unit was added near the denitrification reactors. When reducing water consumption during the summer, overheating of the system becomes a problem. A farm the size of Nijvis has heat production through animal metabolism of roughly 400 kw.

Conclusion

Investment in effluent treatment has helped control waste materials at the Netherlands' largest eel farm. The equipment used is well known for other applications, and there is nothing revolutionary involved. The trick to its successful use was to make the individual components work together smoothly and continuously in the fish-farming environment.

(Editor's Note: This article was originally published in the June 2001 print edition of the Global Aquaculture Advocate.)

Now that you've finished reading the article ...

... we hope you'll consider supporting our mission to document the evolution of the global aquaculture industry and share our vast network of contributors' expansive knowledge every week.

By becoming a Global Seafood Alliance member, you're ensuring that all of the pre-competitive work we do through member benefits, resources and events can continue. Individual membership costs just \$50 a year.

Not a GSA member? Join us.

[Support GSA and Become a Member](#)

Authors

-  Andries Kamstra

Andries Kamstra

RIVO – Netherlands Institute for Fisheries Research
Wageningen-UR, The Netherlands

[108,110,46,114,117,45,103,97,119,46,111,118,105,114,64,115,101,105,114,100,110,97]

-  Ep Eding

Ep Eding

Department of Fish Culture and Fisheries
Wageningen-UR, The Netherlands

[108,110,46,114,117,45,103,97,119,46,118,110,101,118,46,103,108,97,64,103,110,105,100,101,46,112,101]

-  Oliver Schneider

Oliver Schneider

Department of Fish Culture and Fisheries
Wageningen-UR, The Netherlands

Share

-  [Share via Email](#)
-  [Share on Twitter](#)
-  [Share on Facebook](#)
-  [Share on LinkedIn](#)

Tagged With

[Ep Eding](#) [Andries Kamstra](#) [Oliver Schneider](#) [eel farm](#)

Related Posts

Innovation & Investment

[American Unagi brings eel farming back 'home'](#)

Sara Rademaker launched American Unagi to shift eel farming to American soil, where the eels are from. Why? Because of the novelty, and because she saw an opportunity to do things better.

Intelligence

[A brief look at genetically modified salmon](#)

If approved by FDA, fast-growing genetically modified salmon will provide a safe and nutritious product similar to other farmed Atlantic salmon.

Responsibility

[A helping hand to lend: UK aquaculture seeks to broaden its horizons](#)

Aquaculture is an essential contributor to the world food security challenge, and every stakeholder has a role to play in the sector's evolution, delegates were told at the recent Aquaculture's Global Outlook: Embracing Internationality seminar in Edinburgh, Scotland.

Health & Welfare

[A holistic management approach to EMS](#)

Early Mortality Syndrome has devastated farmed shrimp in Asia and Latin America. With better understanding of the pathogen and the development and improvement of novel strategies, shrimp farmers are now able to better manage the disease.

About The Advocate

The Responsible Seafood Advocate supports the Global Seafood Alliance's (GSA) mission to advance responsible seafood practices through education, advocacy and third-party assurances.

[Learn More](#)

Search Responsible Seafood Advocate



Advertising Opportunities

[2022 Media & Events Kit](#)

Categories

[Aquafeeds](#) > [Health & Welfare](#) > [From Our Sponsors](#) > [Innovation & Investment](#) > [Intelligence Intelligence](#) > [Responsibility](#) > [Fisheries](#) > [Artículos en Español](#) >

Don't Miss an Article

Featured

- [Health & Welfare](#) [An update on vibriosis, the major bacterial disease shrimp farmers face](#)
- [Intelligence](#) [A seat at the table: Fed By Blue team says aquaculture needs a stronger voice](#)
- [Responsibility](#) [Quantifying habitat provisioning at macroalgae cultivation locations](#)

Popular Tags

All Tags

Recent

- [Fisheries](#) [Second Test: Another filler for the fisheries category](#)
- [Fisheries](#) [Test: This is filler for the fisheries Category](#)
- [Aquafeeds](#) [Test Article](#)
- [Responsibility](#) [Study: Climate change will shuffle marine ecosystems in unexpected ways as ocean temperature warms](#)
- [Health & Welfare](#) [Indian shrimp researchers earn a patent for WSSV diagnostic tool](#)



- [About](#)
- [Membership](#)
- [Resources](#)
- [Best Aquaculture Practices \(BAP\)](#)
- [Best Seafood Practices \(BSP\)](#)
- [GOAL Events](#)
- [Advocate Magazine](#)
- [Aquademia Podcast](#)
- [Blog](#)
- [Contact](#)

Stay up to date with GSA

- 
- 
- 
- 
- 

Copyright © 2024 Global Seafood Alliance
All rights reserved.

- [Privacy](#)
- [Terms of Use](#)
- [Glossary](#)