## Global Seafood Alliance Logo

- GOAL Events
- Advocate Magazine
- Aquademia Podcast
- Blog
- Contact
- 0
- . 4
- X
- in
- 🔼
- <u>Log In</u>

- About
  - Who We Are
  - Our History
  - Our Team
  - Sustainable Development Goals
  - <u>Careers</u>
- <u>Membership</u>
  - o <u>Overview</u>
  - Our Members
  - Corporate Membership
- Resources
- Certification
  - Best Aquaculture Practices
  - Best Seafood Practices

Search...



## Log In

- About
  - o Who We Are
  - Our History
  - o Our Team
  - Sustainable Development Goals
  - Careers
- Membership
  - o <u>Overview</u>
  - Our Members
  - o Corporate Membership
- Resources
- Certification
  - Best Aquaculture Practices
  - Best Seafood Practices
- GOAL Events
- Advocate Magazine

- Aquademia Podcast
- Blog
- Contact



# Vibrio control in shrimp farming: Part 1

Responsible Seafood Advocate logo

11 September 2015 Stephen G. Newman, Ph.D.



# Examining all inputs into maturation and early stages of larval culture is critical



Artemia culture systems can be significant sources of bacterial contamination. Mitigants range from bacterial amendments to disinfection to kill surfaceattached Vibrios.

Perhaps the single greatest problem affecting shrimp hatcheries is high mortality in the early stages of larval culture. The term "zoeae syndrome" was coined to describe the affected stage. Typically, the animals do not molt from Z1 to Z2, and very high levels of mortality ensue. Various strains of *Vibrio* bacteria have been implicated in this process.

The challenge for hatchery managers is to identify gaps in biosecurity and how to plug them without creating a production environment that has been manipulated to the point where other problems can readily present themselves. The idea is to control the bacteria without creating niches for other potential pathogens.

# Ubiquitous bacteria

It is important to appreciate that bacteria are everywhere. Life depends on them. They are critical for ecological stability and recycling of nutrients, and play a myriad of other roles that we are just beginning to appreciate.

Most bacteria are benign, while some can negatively impact animals that have been made susceptible to them for various reasons, of which the presence of stressors is usually a critical element. A very few are obligate pathogens that kill animals merely by being present. Most of the problems in hatcheries arise from bacterial species that are not obligate pathogens.

It is also important to realize there has been an inordinate focus on *Vibrios* when many other species of bacteria could (and do) cause problems. Control efforts should not be focused on total elimination of all bacteria, but instead should target those production areas where it is possible to minimize the overall bacterial loads.

# **Production stages connected**

As with any agricultural process, aquaculture production stages are linked and overlap others. Broodstock sold in huge numbers commercially become the sources of eggs and nauplii that, in turn, are sold to become the source of postlarval shrimp stocked in production systems for growout to commercial sizes.

Broodstock come from a variety of sources. Most — especially for Pacific white shrimp (*Litopenaeus vannamei*) — are from commercial operations that sell genetically selected animals that are usually also specific-pathogen-free (SPF), typically for pathogens the World Organization for Animal Health dictates should not be present.

Black tiger shrimp, *Penaeus monodon*, are available from similar sources, although there is still significant use of wild broodstock at black tiger farms.

## **Routine procedures for maturation facilities**

The use of SPF animals has had a dramatic impact on shrimp farming globally, although it has not always proven to be the salvation that many envisioned it could be. There are many reasons for this. Suffice it to say that regardless of the source of the animals, maturation facilities should all be doing certain things routinely.

Aside from efforts to minimize the levels of bacteria entering maturation systems, the focus should be on controlling the potential for horizontal transmission throughout the production process. While mass spawning is the norm, for example, individual spawning offers greater control.

When females spawn, bacteria present on them and in their ovarian fluids and feces can readily attach to the surfaces of eggs. Females should be rinsed with a surface disinfectant such as formalin for a short period before being placed in spawning tanks to lessen the external loads of bacteria.

Females should be removed from spawning tanks as soon as possible after spawning — easier when animals are spawned individually. The eggs should be collected and washed with copious amounts of clean water between surface disinfection protocols using formalin, iodophors or other compounds that have been shown effective in reducing the levels of attached bacteria.

After the eggs hatch into nauplii, healthy nauplii are collected by their attraction to light and seeded into hatchery tanks. Prior to their addition to the tanks, they should be washed in a manner similar to that used to surface disinfect the eggs.

These procedures are aimed at significantly reducing the levels of bacteria on external surfaces. Bacteria present in eggs cannot be eliminated in this manner. If tests determine this is a problem, the broodstock need to be treated with appropriate antibiotics in a clean production system and given feeds that are not carrying potential pathogens.

## **Contamination**

Assuming the water in production systems is properly treated and that an effective tool for managing microbial loads in production tanks is used, perhaps the greatest sources of contamination from bacteria in the early life stages are *Artemia* and algal culture systems.

There are many ways to mitigate these contaminants, ranging from the use of bacterial amendments to the use of chemicals such as formalin or chloramine-T to kill any surface-attached *Vibrios* and airborne contaminants that occur as a result of where and how the *Artemia* are produced. Alternatively, *Artemia* nauplii can be collected and surface disinfected as with shrimp. The use of copious amounts of clean water is essential to dislodge weakened and weakly attached bacteria.

Algae are often heavily contaminated with bacteria, as well. This can be mitigated by the use of closed production systems. Where this is not feasible, microbial amendments can lower overall non-benign bacterial loads.

# **Perspectives**

It is critical to examine all inputs into maturation and the early stages of larval culture to ensure levels of potential pathogens are controlled. As the animals molt and various feeds are added to the production tanks, bacteria will grow. Again, the use of a bacterial bioremediation amendment can help control the levels of *Vibrios* throughout this process.

The last link in this process is the stocking of postlarvae into ponds. Some of the same tools used in disinfection of eggs and nauplii can be employed to lessen the levels of attached bacteria.

#### Author

• Stephen G. Newman, Ph.D.

Stephen G. Newman, Ph.D.

President and CEO AquaInTech Inc. 6722 162nd Place Southwest Lynnwood, Washington 98037-2716 USA

[109,111,99,46,104,99,101,116,45,110,105,45,97,117,113,97,64,109,119,101,110,103,115]

## **Share**

- Share via Email
- **Share on Twitter**
- f Share on Facebook
- in Share on LinkedIn

## **Tagged With**

bacteria biosecurity pathogens vibrio control Stephen G. Newman Shrimp Vibrio

#### **Related Posts**

Health & Welfare

## Vibrio control in shrimp farming: Part 2

Strategies to control Vibrio bacteria at aquaculture farms must revolve around limiting inputs, including those from postlarvae suppliers, those that come in with the water, those that remain in the soil between crops and those in the feed and accumulated organic matter.

Health & Welfare

## A study of Zoea-2 Syndrome in hatcheries in India, part 1

Indian shrimp hatcheries have experienced larval mortality in the zoea-2 stage, with molt deterioration and resulting in heavy mortality. Authors investigated the problem holistically.

Aquafeeds

## Global brine shrimp supply a potential bottleneck to aquaculture expansion, part 1

The brine shrimp artemia is critical for larval fish and shrimp diets. As aquaculture expands, an effective artificial replacement diet for artemia is a major consideration for the industry.

Intelligence

## As ocean temperatures rise, so too will vibrio outbreaks

A study using a half-century of data has linked climate change and warming sea temperatures with an increase in illnesses from the common vibrio bacteria. Shellfish growers, fighting a particularly virulent strain of Vibrio parahaemolyticus, are changing their harvest protocols.

#### **About The Advocate**

The Responsible Seafood Advocate supports the Global Seafood Alliance's (GSA) mission to advance responsible seafood practices through education, advocacy and third-party assurances.

ī	Γ.	_	٠,	-	. 1	١,	ť,	٠.	
		Ω,	-11	rT	1	IV/	16	ור	ኅ

Search Responsible Seafood Advocate Search Search



## **Advertising Opportunities**

2022 Media & Events Kit

## Categories

Aquafeeds > Health & Welfare Health & Welfare > From Our Sponsors > Innovation & Investment > Intelligence > Responsibility > Fisheries > Artículos en Español >

#### Don't Miss an Article

#### **Featured**

- Health & Welfare An update on vibriosis, the major bacterial disease shrimp farmers face
- Intelligence A seat at the table: Fed By Blue team says aquaculture needs a stronger voice
- Responsibility Quantifying habitat provisioning at macroalgae cultivation locations

## **Popular Tags**

All Tags ✓

#### Recent

- Fisheries Second Test: Another filler for the fisheries category
- Fisheries Test: This is filler for the fisheries Category
- Aguafeeds Test Article
- Responsibility Study: Climate change will shuffle marine ecosystems in unexpected ways as ocean temperature warms
- Health & Welfare Indian shrimp researchers earn a patent for WSSV diagnostic tool



- About
- Membership
- Resources
- Best Aquaculture Practices (BAP)
- Best Seafood Practices (BSP)
- GOAL Events
- Advocate Magazine
- Aquademia Podcast
- Blog
- Contact

## Stay up to date with GSA

- (
- \_
- 🔊
- 🔼

Copyright  $\ensuremath{^{\odot}}$  2024 Global Seafood Alliance All rights reserved.

Privacy Terms of Use Glossary