<u>Global Seafood Alliance Logo</u>

- GOAL Events
- <u>Advocate Magazine</u>
- <u>Aquademia Podcast</u>
- <u>Blog</u>
- <u>Contact</u>
- 0
- f
- 🗙
- in
- 🕨
- <u>Log In</u>
- \square
 - <u>About</u>
 - <u>Who We Are</u>
 - Our History
 - <u>Our Team</u>
 - Sustainable Development Goals
 - <u>Careers</u>
 - <u>Membership</u>
 - <u>Overview</u>
 - <u>Our Members</u>
 - Corporate Membership
 - <u>Resources</u>
 - <u>Certification</u>
 - Best Aquaculture Practices
 - Best Seafood Practices

Search...

Q

- <u>Log In</u>
 - <u>About</u>
 - Who We Are
 - <u>Our History</u>
 - Our Team
 - Sustainable Development Goals
 - <u>Careers</u>
 - <u>Membership</u>
 - <u>Overview</u>
 - Our Members
 - Corporate Membership
 - <u>Resources</u>
 - <u>Certification</u>
 - <u>Best Aquaculture Practices</u>
 - Best Seafood Practices
 - GOAL Events
 - <u>Advocate Magazine</u>

- <u>Aquademia Podcast</u>
- <u>Blog</u>
- <u>Contact</u>



Water composition and shrimp pond management

Responsible Seafood Advocate logo 1 October 2000 Claude E. Boyd, Ph.D.

Comparing seawater and brackish water



The common practice of liming pond bottoms between crops can be avoided by measuring soil pH and liming only the acidic pond soils.

Natural waters have a very complex chemical composition and may contain almost any substance found in the atmosphere, in the Earth's crust, or in living organisms. They also may contain substances synthesized by man. However, the major dissolved constituents in natural water are a few inorganic substances including calcium, magnesium, potassium, sodium, bicarbonate, sulfate and chloride.

Seawater characteristics

In addition to such substances as those listed above, seawater also has substantial concentrations of bromine, strontium, silicon and boron. The concentrations of the major dissolved inorganic substances in seawater are provided in Table 1.

The salinity of water is the concentration of all dissolved ions, and for average seawater, the sum of the ions is about 34,500 milligrams per liter or 34.5 parts per thousand (ppt). The total alkalinity of water is the concentration of bases expressed as equivalent calcium carbonate. For seawater, the primary base is bicarbonate, and the average seawater bicarbonate concentration of 142 milligrams per literis equivalent to a total alkalinity of 116 milligrams per liter.

Boyd, Average concentrations of major dissolved substances, Table 1

Substance	Seawater (mg/l)	River Water (mg/l)
Chloride (Cl)	19,000	7.8
Sodium (Na)	10,500	6.3
Sulfate (SO ₄)	2,700	11.2
Magnesium (Mg)	1,350	4.1
Calcium (Ca)	400	15.0
Potassium (K)	380	2.3
Bicarbonate (HCO ₃)	142	58.4
Bromide (Br)	65	0.02
Strontium (Sr)	8.0	0.1
Silicate (SiO ₂)	6.4	13.1
Boron (B)	4.6	0.1

Table 1. Average concentrations of major dissolved substances in seawater and river water.

The total hardness of water is the concentration of divalent cations expressed as equivalent calcium carbonate. The divalent cations in seawater are calcium, magnesium, and strontium. Average seawater has a total hardness of 6,569 milligrams per liter. The pH of seawater is normally about 8.1 to 8.3. Except for silicate and bicarbonate, seawater is two or three orders of magnitude more concentrated than river water in each major substance.

Brackish water characteristics

Brackish water results from the dilution of seawater by river flow. The average concentrations of major dissolved substances in river water also are provided in Table 1.

The average salinity of river water is about 120 milligrams per liter (0.12 ppt). Total hardness and total alkalinity average about 55 milligrams per liter and 48 milligrams per liter, respectively. The pH of river water is highly variable, but it will usually be between 6 and 8.

Concentration proportional to dilution

3/20/2024

Brackish water is more dilute than seawater in most major substances, with the decreases in substance concentration almost directly proportional to the degree of dilution with freshwater. For example, a 50 percent dilution of freshwater and seawater would provide a salinity of roughly 17.25 ppt and a sodium concentration of about 5,250 milligrams per liter.

One practical application of the dilution concept is in acclimation of postlarvae at a hatchery. In diluting salinity to match the level at ponds, hatchery managers may inadvertently dilute alkalinity to levels that would fail to buffer declines in pH during packing and transport. This stress could be avoided by measuring alkalinity in the mixed water and adjusting it to approximately 100 milligrams per liter before packing.

Some river waters have higher concentrations of bicarbonate and silicate than seawater, so brackishwater may sometimes contain more bicarbonate and silicate than seawater. This is more often the case for silicate than for bicarbonate, as average river water is lower than average seawater in bicarbonate. In general, rivers flowing from humid areas will be more dilute in major constituents than those flowing from more arid regions. Large rivers will always be more dilute than seawater with respect to most dissolved substances.

The importance of liming materials in aquaculture

Water composition and liming

Shrimp farmers like to apply liming materials to ponds. Some uses seem justifiable, but others are not.

At high application rates (approximately 1 metric tons (MT) per hectare), lime apparently acts as a water-purifying agent that can improve water quality in highly eutrophic ponds by flocculating phytoplankton, bacteria, and suspended particles. Asian shrimp farmers frequently take advantage of this property on an emergency basis to deal with excessive algal blooms, deteriorating water quality, or low-oxygen problems.

Routine additions of small quantities of lime have little appreciable effect on water quality. The equilibrium concentrations of total hardness and total alkalinity in an open system at equilibrium with atmospheric carbon dioxide and calcium carbonate are 60 to 80 milligrams per liter. Thus, application of liming materials to ponds filled with seawater or brackish water is seldom useful because the materials will not dissolve.

This is true for all types of liming materials (calcium carbonate, calcium oxide, and calcium hydroxide). Shrimp pond waters usually have a pH of 7.5 to 8.5, and application of calcium oxide or calcium hydroxide will remove carbon dioxide and cause the pH to increase. This can have undesirable consequences on productivity by limiting phytoplankton growth.

Liming and shrimp molting

Some farmers think that applying liming materials increases the calcium concentration in the water and helps shrimp molt. This is not very likely, as seawater and brackish water already contain a lot of calcium. For example, suppose that 50 kilograms per hectare of calcium oxide is added to a 1-hectare pond that is 1 meter deep. The calcium oxide dose contains 28 kg of calcium. Even if the calcium oxide dissolves completely, calcium would only increase by 2.8 milligrams per liter in the pond.

Type and size of limestone

Some farmers use dolomitic limestone because they think the magnesium in the dolomite will be beneficial. Again, shrimp pond water contains so much magnesium that the small increase resulting from dolomite application would be meaningless.

Obviously, shrimp farmers should measure total alkalinity in pond water and apply liming materials to maintain concentrations. Alkalinity should be maintained at a minimum of 70 to 80 milligrams per liter, and levels of 100 to 120 milligrams per liter are better. Where liming is needed to improve alkalinity, it makes no difference if the liming material was produced from calcitic limestone (calcium carbonate) or dolomitic limestone (calcium magnesium carbonate).

When purchasing agricultural limestone (pulverized calcium or calcium magnesium carbonate), the most finely ground material should be selected. Limestone is not very soluble, and unless the particles are very fine, they will settle to the bottom and not dissolve.

Liming and pH

Because of the high concentration of ions in seawater, pond soils often become saturated with basic ions such as calcium, magnesium, sodium, and potassium. Thus, pond soils are saline and usually have a pH of 8 or above.

Pond managers tend to lime bottom soils routinely between crops. This practice is not necessary unless pond soil pH is below 7. Much expense and effort can be avoided by simply measuring pond soil pH and liming only the acidic soils.

Silicate, potassium and phytoplankton

Some farmers use silicate fertilization to encourage diatoms. Literature suggests that if silicate concentration is above 1 milligrams per liter silicon (Si), diatoms will not be limited by a shortage of this element. Average Si concentrations in seawater and freshwater are about 3 and 6 milligrams per liter, respectively. Thus, silicate is probably not limiting in most shrimp ponds, and especially in ponds with low-salinity water.

Potassium is a key nutrient in plant growth, but phytoplankton growth is not limited by potassium at concentration levels above 1 or 2 milligrams per liter. Inclusion of potassium in shrimp pond fertilizers is usually not necessary.

Sulfate and hydrogen sulfide

Shrimp pond water has a very high sulfate concentrations. When anaerobic conditions develop in pond bottom soils, certain bacteria can use the oxygen from sulfate in respiration. The organisms produce hydrogen sulfide as a metabolic waste that is extremely toxic to shrimp. Because of the high availability of sulfate in their pond water, shrimp farmers should be especially vigilant to avoid the development of anaerobic conditions at the soil-water interface.

Conclusion

There is no need to make analyses of shrimp pond water for most of the substances shown in Table 1, for they will be at high concentration. However, measurements of total alkalinity could be useful in determining if waters need to be limed. Total alkalinity should be maintained at a minimum of 80 milligrams per liter, with levels of 100 to 120 milligrams per liter preferred. Silicate analyses will reveal if pond waters are naturally low in silicate and could possibly benefit from silicate fertilization. Pond bottom soils normally will not benefit from liming unless the pH is below 7.

(*Editor's Note: This article was originally published in the October 2000 print edition of the* Global Aquaculture Advocate.)

Now that you've finished reading the article ...

... we hope you'll consider supporting our mission to document the evolution of the global aquaculture industry and share our vast network of contributors' expansive knowledge every week.

By becoming a Global Seafood Alliance member, you're ensuring that all of the pre-competitive work we do through member benefits, resources and events can continue. Individual membership costs just \$50 a year.

Not a GSA member? Join us.

Support GSA and Become a Member

Author



Claude E. Boyd, Ph.D.

Professor, Department of Fisheries and Allied Aquacultures International Center for Aquaculture and Aquatic Environments Auburn University, Alabama 36849-5419 USA

[117,100,101,46,110,114,117,98,117,97,64,49,101,99,100,121,111,98]

Share

- <u>Share via Email</u>
- **J** Share on Twitter
- **f** Share on Facebook
- in <u>Share on LinkedIn</u>

Tagged With

water composition Claude E. Boyd shrimp ponds

Related Posts

Responsibility

<u>Appraising pond liners for shrimp culture</u>

The use of plastic-lined ponds by shrimp farmers can significantly improve production efficiency, support more production cycles per year, and higher mechanical aeration rates and stocking densities. The capital cost of lining ponds can be very significant, so a thorough feasibility analysis is recommended when considering this production tool.

Responsibility

Carbon-nitrogen ratio management

Carbon-nitrogen (C:N) ratio is an important variable related to soil fertility. Intensive ponds have lower C:N ratios, while higher ratios are found in extensive ponds and those constructed in organic soil.

Responsibility

Dissolved salts in water for inland, low-salinity shrimp culture

Low-salinity shrimp culture is an important activity in Thailand. Ponds contain water of 2 to 5 ppt salinity, mixing brine solution from coastal seawater with freshwater.

Responsibility

Calcium and magnesium use in aquaculture

Aquatic plants and animals get the essential nutrients calcium and magnesium from water and food. Calcium concentrations impact the hydration and development of eggs in a hatchery, where calcium carbonate precipitation can be troublesome.

About The Advocate

The Responsible Seafood Advocate supports the Global Seafood Alliance's (GSA) mission to advance responsible seafood practices through education, advocacy and third-party assurances.

Learn More

Search Responsible Seafood Advocate	Search	Search Q
Search Responsible Searbou Auvocale	Search	Search 🗸



Listen to the seafood industry's top podcast

Advertising Opportunities

2022 Media & Events Kit

Categories

Aquafeeds > Health & Welfare > From Our Sponsors > Innovation & Investment > Intelligence > Responsibility Responsibility > Fisheries > Artículos en Español >

Don't Miss an Article

Featured

- Health & Welfare An update on vibriosis, the major bacterial disease shrimp farmers face
- Intelligence A seat at the table: Fed By Blue team says aquaculture needs a stronger voice
- <u>Responsibility Quantifying habitat provisioning at macroalgae cultivation locations</u>

Popular Tags

3/20/2024

All Tags

Recent

- Fisheries Second Test: Another filler for the fisheries category
- <u>Fisheries Test: This is filler for the fisheries Category</u>
- <u>Aquafeeds Test Article</u>

×

- <u>Responsibility Study: Climate change will shuffle marine ecosystems in unexpected ways as ocean</u> <u>temperature warms</u>
- Health & Welfare Indian shrimp researchers earn a patent for WSSV diagnostic tool



Listen to the seafood industry's top podcast

- <u>About</u>
- <u>Membership</u>
- <u>Resources</u>
- <u>Best Aquaculture Practices (BAP)</u>
- <u>Best Seafood Practices (BSP)</u>
- GOAL Events
- <u>Advocate Magazine</u>
- <u>Aquademia Podcast</u>
- <u>Blog</u>
- <u>Contact</u>

Stay up to date with GSA

- Ø
- **f**
- 🕅
- in
- 🕨

Copyright © 2024 Global Seafood Alliance All rights reserved. <u>Privacy</u> <u>Terms of Use</u> <u>Glossary</u>