



- [GOAL Events](#)
- [Advocate Magazine](#)
- [Aquademia Podcast](#)
- [Blog](#)
- [Contact](#)

-
-
-
-
-

- [Log In](#)



- [About](#)
 - [Who We Are](#)
 - [Our History](#)
 - [Our Team](#)
 - [Sustainable Development Goals](#)
 - [Careers](#)
- [Membership](#)
 - [Overview](#)
 - [Our Members](#)
 - [Corporate Membership](#)
- [Resources](#)
- [Certification](#)
 - [Best Aquaculture Practices](#)
 - [Best Seafood Practices](#)

[Log In](#)

- [About](#)
 - [Who We Are](#)
 - [Our History](#)
 - [Our Team](#)
 - [Sustainable Development Goals](#)
 - [Careers](#)
- [Membership](#)
 - [Overview](#)
 - [Our Members](#)
 - [Corporate Membership](#)
- [Resources](#)
- [Certification](#)
 - [Best Aquaculture Practices](#)
 - [Best Seafood Practices](#)
- [GOAL Events](#)
- [Advocate Magazine](#)

- [Aquademia Podcast](#)
- [Blog](#)
- [Contact](#)



 Health & Welfare
Health & Welfare

Water quality concerns may move up food chain

 [Responsible Seafood Advocate logo](#)

1 June 2002 George J. Flick, Jr., Ph.D.



A look at brominated flame retardants and other chemicals' impact on seafood



Findings of various recent studies on water quality in U.S. waterways should be of concern to various industries, including those involved in seafood production and those that use source water.

Recently published studies have detected a variety of chemical compounds in the environment and established that they can accumulate in humans. Additional research on chemical residues in fish, shellfish, and other foods is likely in the future.

U.S. waterway study

During 1999 and 2000, the United States Geological Survey (USGS) surveyed 139 streams across 30 states to determine the types and concentrations of chemicals in U.S. waterways. The study showed that many waterways contained chemicals such as painkillers, antibiotics, caffeine, steroids, insect repellents, reproductive hormones, personal care products, antimicrobial disinfectants, fire retardants, detergent metabolites, and petroleum products. These compounds were present in 80 percent of the streams sampled – not altogether surprising, since current water treatment systems are not capable of removing them from the waste stream.

Chemicals in waterways

The surveyed chemicals are referred to as organic wastewater contaminants or pharmaceutical and personal care pollutants. Their concentrations were generally low and rarely exceeded drinking-water guidelines. However, many of the compounds do not have established guidelines.

In the study of 95 chemical compounds, those most frequently detected were coprostanol (a fecal steroid), cholesterol (an animal steroid), N,N-diethyltoluamide (an insect repellent), caffeine (a stimulant), triclosan (an antimicrobial disinfectant), tri(2-chloroethyl)phosphate (a fire retardant), and 4-nonylphenol (a nonionic detergent metabolite).

The presence of these compounds is cause for increased concern for potential adverse effects on human and ecological health. It is anticipated that many of these compounds may soon appear in the tissues of fish, if their presence has not already been detected.

Prevalent products



With increasing concern about water quality, further research on chemical residues in fish, shellfish, and other foods is likely in the future.

It is not surprising that detergent metabolites, steroids, and plasticizers comprise approximately 80 percent of the total chemical concentration, since these are derived from common household products and pharmaceuticals. Table 1 shows concentrations of organic wastewater contaminants by category.

The organic chemicals most frequently discussed by environmentalists are antibiotics, prescription and nonprescription drugs, and reproductive hormones. In this list, nonprescription drugs were found with the greatest frequency. Reproductive hormones may present the greatest environmental impact to aquatic organisms, since even low levels of these compounds produce undesirable effects.

The USGS study did not determine whether the compounds were degraded in the environment. Many compounds undergo decomposition through biodegradation, and many of the products of decomposition actually present a greater danger than the original compounds.

The possible toxicological consequences of a synergistic effect between some of the chemicals are also unknown. As always, the presence of antibiotics in the environment raises questions on the development of antibiotic-resistant microorganisms.

Flick, Total measured concentration of organic wastewater contaminants, Table 1

Contaminant	Concentration (%)
Steroids	21.5
Nonprescription drugs	9
Insect repellents	1
Detergent metabolites	36
Disinfectants	3
Plasticizers	20
Fire Retardants	1
Anitbiotics	3
Insecticides	1
Polynucleated aromatic hydrocarbons	1
Reproductive hormones	1
Other prescription drugs	1
Antioxidants	2
Fragrances	1
Solvents	1

Source: United States Geological Survey, 2002.

Table 1. Total measured concentration of organic wastewater contaminants by general use category.

Norwegian BFR study

A Norwegian study of eight selected brominated flame retardants (BFR) in human blood serum, sampled from eight groups of differing age and gender at six time periods between 1977 and 1999, was also recently published.

The serum concentrations of seven of the eight BFRs increased during the entire period. The sum of six polybrominated diphenyl ethers increased from 0.44 nanogram per gram lipids in 1977 to 3.3 nanogram per gram lipids in 1999. The BFR serum concentrations from the various age groups were similar, except in the case of very young children (ages 0-4), who exhibited concentrations that were 1.6 to 3.5 times higher. The sources of the BFRs were not suggested in the study.

BFRs are chemically similar to polychlorinated biphenyls (PCBs). They are persistent and bioaccumulative, and expected to become widely dispersed in the environment. Studies have shown some BFR compounds have biological and behavioral effects on animals. Consequently, they have the potential to pose a threat to human and animal health.

Impact on seafood

The impact of the compounds reported in the two studies on fish and shellfish is unknown. There is very little published data on the subject. One reason for the lack of data on residues in aquatic food products is that, until recently, few analytical methods were capable of detecting the compounds at low environmental concentrations.

Since wild fish and shellfish have been the subject of recent food safety concerns, aquatic foods will likely receive a high priority for investigation in further studies of chemical residues in the food chain. Problems related to scombrototoxin or histamine toxicity, methyl mercury, dioxins, polychlorinated biphenyls, natural toxicants (such as paralytic shellfish poisoning), pathogenic microorganisms (*Vibrio cholera*, *V. parahaemolyticus*, and *V. vulnificus*) and ciguatera toxin have all been reported in recent years.

Consumer response

While concerns about the safety of fish and shellfish consumption will likely continue, cultured products produced in controlled environments such as recirculating aquaculture systems may be perceived by consumers as having added value when compared with other products in the marketplace. Such consumer interest may encourage commercial aquaculturists to implement practices that further limit chemical use, such as improved biosecurity programs as an alternative to antibiotic administration for disease treatment.

(Editor's Note: This article was originally published in the June 2002 print edition of the Global Aquaculture Advocate.)

Now that you've finished reading the article ...

... we hope you'll consider supporting our mission to document the evolution of the global aquaculture industry and share our vast network of contributors' expansive knowledge every week.

By becoming a Global Seafood Alliance member, you're ensuring that all of the pre-competitive work we do through member benefits, resources and events can continue. Individual membership costs just \$50 a year.

Not a GSA member? Join us.

[Support GSA and Become a Member](#)

Author



George J. Flick, Jr., Ph.D.

Food Science and Technology Department
Virginia Tech (0418)
Blacksburg, Virginia 24061 USA

[117,100,101,46,116,118,64,103,107,99,105,108,102]

Share

- [✉ Share via Email](#)
- [🐦 Share on Twitter](#)
- [f Share on Facebook](#)
- [in Share on LinkedIn](#)

Tagged With

[water quality](#) [George J. Flick](#)

Related Posts

Responsibility

[A review of water quality improvement products](#)

Prof. Boyd examines products used by aquafarmers to improve water quality and conditions in their ponds and discusses their efficacy.

Health & Welfare

[10 paths to low productivity and profitability with tilapia in sub-Saharan Africa](#)

Tilapia culture in sub-Saharan Africa suffers from low productivity and profitability. A comprehensive management approach is needed to address the root causes.

Intelligence

[10 takeaways from GOAL 2019 in Chennai, India](#)

The Global Aquaculture Alliance held its GOAL conference in Chennai, India, and recruited a host of experts in various fields to share their expertise.

Intelligence

[A brief look at genetically modified salmon](#)

If approved by FDA, fast-growing genetically modified salmon will provide a safe and nutritious product similar to other farmed Atlantic salmon.

About The Advocate

The Responsible Seafood Advocate supports the Global Seafood Alliance's (GSA) mission to advance responsible seafood practices through education, advocacy and third-party assurances.

[Learn More](#)

Search Responsible Seafood Advocate


Search 



Advertising Opportunities

[2022 Media & Events Kit](#)

Categories

[Aquafeeds](#) >  [Health & Welfare](#) [Health & Welfare](#) > [From Our Sponsors](#) > [Innovation & Investment](#) > [Intelligence](#) > [Responsibility](#) > [Fisheries](#) > [Artículos en Español](#) >

Don't Miss an Article

Featured

- [Health & Welfare](#) [An update on vibriosis, the major bacterial disease shrimp farmers face](#)
- [Intelligence](#) [A seat at the table: Fed By Blue team says aquaculture needs a stronger voice](#)
- [Responsibility](#) [Quantifying habitat provisioning at macroalgae cultivation locations](#)

Popular Tags

All Tags ▼


Recent

- [Fisheries](#) [Second Test: Another filler for the fisheries category](#)
- [Fisheries](#) [Test: This is filler for the fisheries Category](#)
- [Aquafeeds](#) [Test Article](#)
- [Responsibility](#) [Study: Climate change will shuffle marine ecosystems in unexpected ways as ocean temperature warms](#)
- [Health & Welfare](#) [Indian shrimp researchers earn a patent for WSSV diagnostic tool](#)



- [About](#)
- [Membership](#)
- [Resources](#)
- [Best Aquaculture Practices \(BAP\)](#)
- [Best Seafood Practices \(BSP\)](#)
- [GOAL Events](#)
- [Advocate Magazine](#)
- [Aquademia Podcast](#)
- [Blog](#)
- [Contact](#)

Stay up to date with GSA

- 
- 
- 
- 
- 

Copyright © 2024 Global Seafood Alliance

All rights reserved.

[Privacy](#)

[Terms of Use](#)

[Glossary](#)

