



- [GOAL Events](#)
- [Advocate Magazine](#)
- [Aquademia Podcast](#)
- [Blog](#)
- [Contact](#)
-
-
-
-
-
- [Log In](#)



- [About](#)
 - [Who We Are](#)
 - [Our History](#)
 - [Our Team](#)
 - [Sustainable Development Goals](#)
 - [Careers](#)
- [Membership](#)
 - [Overview](#)
 - [Our Members](#)
 - [Corporate Membership](#)
- [Resources](#)
- [Certification](#)
 - [Best Aquaculture Practices](#)
 - [Best Seafood Practices](#)

Search...



[Log In](#)

- [About](#)
 - [Who We Are](#)
 - [Our History](#)
 - [Our Team](#)
 - [Sustainable Development Goals](#)
 - [Careers](#)
- [Membership](#)
 - [Overview](#)
 - [Our Members](#)
 - [Corporate Membership](#)
- [Resources](#)
- [Certification](#)
 - [Best Aquaculture Practices](#)
 - [Best Seafood Practices](#)
- [GOAL Events](#)
- [Advocate Magazine](#)
- [Aquademia Podcast](#)
- [Blog](#)
- [Contact](#)



White shrimp culture in China performance and impacts



1 October 2005 Dr. Wang Qing-Yin Dr. Li Jian Dr. Yang Cong-Hai



Species quickly becomes one of China’s most important



This intensive *L. vannamei* farm in China uses plastic-lined ponds and regular aeration.

Pacific white shrimp (*Litopenaeus vannamei*) were experimentally introduced in China in 1988, and the reproductive cycle was closed in hatcheries after 1992. In 1998, a specific pathogen-free stock was introduced from Hawaii, and spawners and postlarvae have been introduced on a commercial scale since then. Nowadays, over 1,000 hatcheries supply *L. vannamei* postlarvae to shrimp farmers in China, most of which are located in the provinces of Guangdong and Hainan in the southern part of the country.

Strong production

In 2003, China’s hatcheries produced over 241 billion shrimp postlarvae – 112,622 billion *L. vannamei* and 128,653 billion other species. The yield of *L. vannamei* from sea water farm ponds was 308,947 metric tons (MT), which accounted for 63 percent of the total marine shrimp-farming production, while 296,312 MT were reported from fresh water ponds (Figs. 1 and 2). With an estimated total value of U.S. \$1 billion, *L. vannamei* has quickly become one of the most important aquaculture species in China in terms of production scale and economic value.




Fig. 1: China’s total farmed shrimp production in 2003 was 789,373 MT.




Fig. 2: China’s shrimp-farming area in 2003 totaled 221,496 ha.

L. vannamei adapt well to higher stocking densities during culture, relative to other farmed shrimp species. In southern China, two crops can normally be harvested per year, and production in well-managed ponds usually ranges 7.5-15 metric ton per hectare per crop. In the provinces of Shandong, Hebei, and Liaoning in northern China, where water temperatures allow only about 100 days of culture annually, one good crop of *L. vannamei* can be achieved.

Greenhouses or small-scale ponds covered with plastic film are used to headstart the culture cycle of *L. vannamei* in the early spring or later autumn in the northern pro-vinces, to ensure a stable supply of fresh live shrimp for markets. In protected, well-managed farming facilities, *L. vannamei* biomass can reach 5 kilogram per square meter per crop and higher (Fig. 3).





Fig. 3: China’s average crop yield of shrimp ponds in 2003.

Impacts



Indoor shrimp farming is practiced in Shandong Province.

After the 1993 collapse of the shrimp-farming industry caused by white spot syndrome virus, the positive impacts of the *L. vannamei* introduction to China are evident. The introduction provided a new species for the industry, and new ideas such as biosecurity and healthier production practices were introduced, along with the development and implementation of new farming models and techniques.

There have also been some negative impacts, including known and new pathogens introduced along with the shrimp. Also, the use of underground water to culture *L. vannamei* in inland areas might have environmental impacts. Finally, the potential impacts of *L. vannamei* escapes to the local ecosystem should be evaluated in the long term.

(Editor's Note: This article was originally published in the October 2005 print edition of the Global Aquaculture Advocate.)

Now that you've finished reading the article ...

... we hope you'll consider supporting our mission to document the evolution of the global aquaculture industry and share our vast network of contributors' expansive knowledge every week.

By becoming a Global Seafood Alliance member, you're ensuring that all of the pre-competitive work we do through member benefits, resources and events can continue. Individual membership costs just \$50 a year.

Not a GSA member? Join us.

[Support GSA and Become a Member](#)


Authors

-  Dr. Wang Qing-Yin

Dr. Wang Qing-Yin

Yellow Sea Fisheries Research Institute
Chinese Academy of Fishery Sciences
106 Nanjing Road, Qingdao 266071
P.R. China

[110,99,46,100,115,46,100,113,46,99,105,108,98,117,112,64,103,110,97,119,121,113]

-  Dr. Li Jian

Dr. Li Jian





Yellow Sea Fisheries Research Institute
Chinese Academy of Fishery Sciences
106 Nanjing Road, Qingdao 266071
P. R. China

-  Dr. Yang Cong-Hai

Dr. Yang Cong-Hai

Yellow Sea Fisheries Research Institute
Chinese Academy of Fishery Sciences
106 Nanjing Road, Qingdao 266071
P. R. China

Share

-  [Share via Email](#)
-  [Share on Twitter](#)
-  [Share on Facebook](#)
-  [Share on LinkedIn](#)

Tagged With

[white shrimp](#) [Wang Qing-Yin](#) [Li Jian](#) [Yang Cong-Hai](#)

Related Posts

[Health & Welfare](#)

[Alphavirus replicon particles potential method for WSSV vaccination of white shrimp](#)

A study demonstrated that VP19 and VP28 white spot syndrome virus envelope proteins expressed by replicon particles provided protection against mortality due to WSSV in shrimp.

Health & Welfare

[Ammonia addition enhances microbial flocs in nursery phase for Pacific white shrimp](#)

In a study, “pre-fertilization” in the nursery phase of a biofloc system for shrimp was tested. The objective was to accelerate the biofloc formation to minimize ammonia concentrations, avoiding high peaks during culture.

Health & Welfare

[A study of Zoea-2 Syndrome in hatcheries in India, part 1](#)

Indian shrimp hatcheries have experienced larval mortality in the zoea-2 stage, with molt deterioration and resulting in heavy mortality. Authors investigated the problem holistically.

Health & Welfare

[Culture of Pacific white shrimp juveniles in super-intensive conditions](#)

Results of this study showed that *L. vannamei* juveniles can be cultured at high densities with adequate survival, growth and feeding efficiency, and with minimal water exchange or full water reuse through recirculation.

About The Advocate

The Responsible Seafood Advocate supports the Global Seafood Alliance’s (GSA) mission to advance responsible seafood practices through education, advocacy and third-party assurances.

[Learn More](#)

Search Responsible Seafood Advocate



Advertising Opportunities

[2022 Media & Events Kit](#)

Categories

[Aquafeeds](#) > [Health & Welfare](#) > [From Our Sponsors](#) > [Innovation & Investment](#) > [Intelligence Intelligence](#) > [Responsibility](#) > [Fisheries](#) > [Artículos en Español](#) >

Don't Miss an Article

Featured

- [Health & Welfare](#) [An update on vibriosis, the major bacterial disease shrimp farmers face](#)
- [Intelligence](#) [A seat at the table: Fed By Blue team says aquaculture needs a stronger voice](#)
- [Responsibility](#) [Quantifying habitat provisioning at macroalgae cultivation locations](#)

Popular Tags

All Tags ▼

Recent


- [Fisheries](#) [Second Test: Another filler for the fisheries category](#)
- [Fisheries](#) [Test: This is filler for the fisheries Category](#)
- [Aquafeeds](#) [Test Article](#)
- [Responsibility](#) [Study: Climate change will shuffle marine ecosystems in unexpected ways as ocean temperature warms](#)
- [Health & Welfare](#) [Indian shrimp researchers earn a patent for WSSV diagnostic tool](#)



- [About](#)
- [Membership](#)
- [Resources](#)
- [Best Aquaculture Practices \(BAP\)](#)
- [Best Seafood Practices \(BSP\)](#)
- [GOAL Events](#)
- [Advocate Magazine](#)
- [Aquademia Podcast](#)
- [Blog](#)
- [Contact](#)

Stay up to date with GSA

- [@](#)



Copyright © 2024 Global Seafood Alliance
All rights reserved.

[Privacy](#)

[Terms of Use](#)

[Glossary](#)